

Three structural remarks on a forced finite-time blowup construction for Navier–Stokes

Kaiyan Ren*

September 10, 2026

Abstract

Under the standing assumption that the forced finite-time blowup construction of OpenAI for the three-dimensional incompressible Navier–Stokes equations is correct as claimed, we record three structural observations: (i) in a fixed background the source’s $O(N^{-1})$ bound for the five-moment increment is non-sharp, the cumulative increment over the full modulation interval being $O(N^{-2})$; (ii) the terminal force F_0 is rigidly equivalent to the truncated-shell commutator; (iii) at $j_0 = 0$ a parity mechanism degenerates the endpoint cone test (B.19) of [S], and this degeneration cannot be avoided within the class (analytic axis profile, vanishing leading residual stress $T_0 = 0$, stress activated at X_0 , the same cone test) (in-class rigidity; see Note 3 below). Statements that a step fails are of level L0; the in-class rigidity holds only within its class definition. We make no claim about the existence or nonexistence of reflection-symmetric forced blowup, and we claim no new Navier–Stokes solution.

Contents

1	Introduction and scope	2
2	Note 1: non-sharpness of the $O(N^{-1})$ five-moment bound	3
2.1	Source statements	3
2.2	Non-sharpness: cumulative moments improve to $O(N^{-2})$	4
2.3	What the improvement does not cover	5
3	Note 2: the terminal shell force F_0, explicit formula and rigid equivalence	6
3.1	Localization and the definition of the force	6
3.2	The terminal value F_0 and the shell formula	7
3.3	Rigid equivalence	8
4	Note 3: the load-bearing role of the bias j_0 — a parity mechanism	9
4.1	Setup and the source endpoint test	9
4.2	Parity: $p_2(\cdot, 0) \equiv 0$ exactly	9
4.3	$p_1 = 2(3 - h)/\Lambda$ and the degeneration of (B.19)	10
4.4	The L0 conclusion and the in-class rigidity	11
5	Acknowledgments and statement	13

*Department of Computer Science and Engineering, The Hong Kong University of Science and Technology, Hong Kong SAR, China. krenab@connect.ust.hk

1 Introduction and scope

The OpenAI construction (one paragraph). [Source] For every positive viscosity the paper constructs, from rest, a three-dimensional incompressible Navier–Stokes solution with uniformly bounded kinetic energy and velocity unbounded in finite time, driven by a smooth compactly supported force (lines 1–6,801–832); the local field is given by Theorem 3.1 of [S] (potential decomposition (14), flat residual (15), exterior heat profile (16), axis growth (17)) (lines 745–799); after spatial–temporal truncation the field is localized to a compact set K , and the residual is extended to a C_c^∞ force (Prop. 10.1 and Lemma 10.2–10.3 of [S]) (lines 7229–7386). The axis construction takes a small bias $0 < j_0 \leq .05$ and $U^* = 4\eta + j_0$ ((28), (lines 9002–9005)); the introduction presents this bias as a deliberate breaking of the $z \mapsto -z$ reflection symmetry (lines 184–190). The parameter order chooses j_0 before Λ ((31)).

Scope of this note. [Assumption] Throughout, we read the source formulas, quantifiers and parameter order under the hypothesis that the construction is correct as claimed: we assume Theorem 3.1, the axis profile and endpoint test of Appendix B of [S], and the modulation–five-moment repair of §4 Steps 2–3 hold as stated, and we then make three observations about their internal structure. This note does not re-prove the existence of the construction, does not connect the finite-dimensional moment maps to the full fluid field, and claims no new Navier–Stokes solution or regularity criterion.

Scope discipline (L0). [Derived] Outward statements use level L0 only: *when the device is extended to the stated setting (including the external limit point $j_0 = 0$, which lies outside the source’s domain, the source allowing only $0 < j_0 \leq .05$ (lines 9002)), a step fails under the device’s own quantifiers.* The statement that there is no $j_0 = 0$ variant within one function class (L1) has been recorded as Prop. 4.9 (the class definition (H1)–(H3) is part of its hypotheses); the existence or nonexistence of reflection-symmetric forced blowup (L2) is not touched by this note, and no such statement appears below. The first cone inequality $a - b_s w > 0$ ($w = N_s/(EQ_s)$ (lines 8323)) and the second cone test $v_s > 2$ are kept separate and are not conflated into one “cone”.

Color code and line references. [Source] is verbatim readable in `navier-stokes.txt`; [Derived] is a rewriting of source formulas under the stated setting; [Assumption] is an additional hypothesis or an unproven step. Every assertion carries line numbers. The internal working record used during the writing (not distributed) contains no independent five-moment M formula or closed form of F_0 ; the formulas of Notes 1 and 2 are rewritten from the cited source lines. Paragraphs of that record marked as superseded (SUPERSEDED) are not cited at all.

Notation, line references and reproducibility. Every (lines xxx) refers to line numbers of the reference text extraction `navier-stokes.txt` of the source PDF ([1]): a plain-text extraction with 10264 lines, 507 517 bytes, each page separated by a marker of the form `===PAGE n===` (165 page markers in all) and no ASCII form-feed characters, with SHA-256

76a236df4f12ab2261f07047ae56a1dcc320b515c55c82653fea48942b10d674.

The extraction was based on `pdftotext`; the version is unknown (the existing extraction carries no version metadata), and the page-marker format is not the `pdftotext` default (which paginates with form feeds), so a fresh `pdftotext` run will not reproduce this file byte for byte. Being a full-text copy of [1], the extraction is not redistributed with this note; the fingerprint above identifies it unambiguously, and the author will supply it on request. A reader working from the source PDF directly can still locate almost every [Source] statement without it, because

this note prints the source’s own tags next to the line anchors: equation numbers such as (B.19) or (10.9), statement numbers such as Prop. B.3 of [S], and, for the extraction, the page markers.

Numbering and citation convention. The theorem-like environments of this note (Theorem/Lemma/Proposition/Corollary/Definition/Claim/Remark) are numbered consecutively by “section.counter” and displayed formulas by (1), (2), All numbers that refer to the source paper are written with “of [S]” (Theorem 3.1 of [S], Prop. B.3 of [S], Lemma B.4 of [S], Cor. B.6 of [S], Theorem 4.6(iii) of [S], etc.) to avoid confusion with the numbering of this note. Parenthesized numbers carrying an appendix letter or a section/paragraph, such as (B.19), (B.26), (10.9), (A.24), always refer to the source formulas; where this note re-typesets a source formula it uses its own number and states the source’s tag and line numbers immediately adjacent.

Status of proofs. This note records three structural observations; it is not a research paper giving complete proofs. Prop. 2.2–2.4, 3.1–3.3, 4.1–4.5 give only argument sketches: every proof block headed “Argument sketch” ends with the explicit statement “This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.” Propositions without a proof block (3.3, 4.4, 4.5) have their derivation in the steps listed in their statements, likewise not constituting a complete proof. The argument of Prop. 4.9 is its items 1–3, which depend on the sketch-only propositions above, and is valid only within the class definition (H1)–(H3). Steps tagged **[Assumption]** are unproven and must not be used as established conclusions; the evidence level is read off the color code: **[Source]** is checkable verbatim against the extracted text, **[Derived]** is a rewriting of source formulas, **[Assumption]** is unproven.

Notation: source coordinates and symbols

[Source] Similarity coordinates and weights (lines 1343–1350):

$$\begin{aligned} \tau &= 1 - t, & A &= \frac{1}{2} + h, & D &= \frac{1}{2} - h, & 0 < h < \frac{1}{2}, \\ z &= q^D \eta, & \tau &= q(1 - \eta^2), & d &= 1 - \eta^2, & L &= 1 - 2h\eta^2, & s &= r^2/2, & X &= s/q. \end{aligned} \quad (1)$$

Hence at $\eta = 0$ one has $d = L = 1$. Profiles (lines 1374–1378):

$$u_\theta^{(0)} = q^{-A} E, \quad u_z^{(0)} = q^{-A} U, \quad ru_r^{(0)} = V_0, \quad p^{(0)} = q^{-2A} \Pi, \quad E = C^{-1} \sqrt{2X} \varphi. \quad (2)$$

Stress coefficients (lines 1488–1496):

$$p_s = \left(\frac{XQ_s}{L}, \frac{XN_s}{LE} \right), \quad s = (a, -b_s), \quad a = 1 - 2D_X \log E, \quad b_s = \frac{2D_X U}{E}. \quad (3)$$

First cone direction $w = N_s/(EQ_s)$ (lines 8323); the first cone inequality $a - b_s w > 0$ (the sufficient test of Lemma 4.5 of [S] and (A.24) (lines 1778, 8324–8325)). Below, $p_1 = p_{s,1}$, $p_2 = p_{s,2}$. The shear $v_s = a(1 + t_s^2) = a + b_s^2/a$ (lines 1748–1751); in the axis region and the reference profile, where $T_0 = 0$ (i.e. $p_s = s$), $v_s = p_1 + p_2^2/p_1$ (for $p_1 > 0$; the p -form appears in (lines 9385–9389)).

2 Note 1: non-sharpness of the $O(N^{-1})$ five-moment bound

2.1 Source statements

[Source] The zero-mean primitives A, B of the periodic shear satisfy (lines 2390–2401)

$$\partial_\varphi A = -\frac{1}{2}(a_L - a_0), \quad \partial_\varphi B = \frac{1}{2}E_0(b_L - b_0), \quad \int_0^1 A d\varphi = \int_0^1 B d\varphi = 0, \quad (4)$$

and vanish near the two ends of $I = [X_-, X_+]$, where the loop is constant, with zero extension off I . The modulation (lines 2402–2408):

$$E_N = E_0 \exp\left(\frac{A(X, \eta, N \log X)}{N}\right), \quad U_N = U_0 + \frac{B(X, \eta, N \log X)}{N}. \quad (5)$$

Pointwise and shear bounds (lines 2421–2426):

$$\|U_N - U_0, E_N - E_0\|_k \leq A_k/N, \quad \|a_N - a_L, b_N - b_L\|_k \leq B_k/N. \quad (6)$$

The exact integral of the five-moment increment (fixed interval away from zero) (lines 2428–2448): put $u_N = U_N - U_0$, $e_N = E_N - E_0$, $\Delta m_N = m_N - m_0$,

$$\Delta m_N(X, \eta) = \int_{X_-}^X \begin{pmatrix} u_N \\ \sqrt{2x} e_N \\ H_0 u_N + U_0 \sqrt{2x} e_N + \sqrt{2x} u_N e_N \\ 2U_0 u_N + u_N^2 - E_0 e_N - e_N^2/2 \\ E_0 e_N/x + e_N^2/(2x) \end{pmatrix} dx. \quad (7)$$

The product rule and (6) give (lines 2449–2452)

$$\|\Delta m_N\|_k \leq D_k/N, \quad \|\Pi_N - \Pi_0^{\text{prof}}\|_k = \|\Delta C_{p,N}\|_k \leq D_k/N. \quad (8)$$

Then take $d_N = -\Delta m_N(Y_1, \eta)$, arranged as (M, J, I, S, C_p) (lines 2497–2511):

$$B(\eta)c + Q_\eta(c, c) = d_N(\eta). \quad (9)$$

and require $N \geq 8 \max_{k=0,2} \nu_k^2 q_k D_k$ so that $\|c\|_{C^2} \leq 2\nu_2 D_2/N$ (lines 2546–2552).

Remark 2.1 (Ordering of rows). [\[Source\]](#) The five integrands of (7) are arranged as (M, I, J, S, C_p) ; the target vector of (9) is arranged as (M, J, I, S, C_p) (lines 2497–2511). The two orderings must not be mixed when citing.

2.2 Non-sharpness: cumulative moments improve to $O(N^{-2})$

Proposition 2.2 (Oscillatory-integral boundary term of order $O(N^{-1})$). *[Derived] Let $[a, b] \subset (0, \infty)$ be fixed, let $P(X, \eta, \varphi)$ be smooth, 1-periodic in φ with zero mean, and let Q be its periodic primitive ($\partial_\varphi Q = P$). The phase $\varphi = N \log X$ is independent of η (lines 2409–2410, 2421). Then*

$$P(X, \eta, N \log X) = \frac{X}{N} \left(\partial_X Q(X, \eta, N \log X) - Q_X(X, \eta, N \log X) \right), \quad (10)$$

so that for a smooth weight g on a fixed interval,

$$\int_a^x g P_N dX = \frac{[XgQ_N]_a^x}{N} - \int_a^x \frac{(Xg)'Q_N + Xg(Q_X)_N}{N} dX$$

is $O(N^{-1})$ uniformly in x . The same holds for any fixed finite order of η -derivatives (constants may depend on the fixed background, the interval and the order of differentiation).

Argument sketch. The chain rule gives (10); the integration by parts is standard. The complete C_η^k estimate (only derivative bounds on a compact parameter set are needed) is not written out in this note. This note does not prove this step; it is stated only as an observation and does not constitute a complete proof. $\square$

Proposition 2.3 (Cumulative five moments $O(N^{-2})$). *[Derived] In a fixed background (U_0, E_0) and fixed k , put $u_N = B_N/N$ and $e_N = E_0(\exp(A_N/N) - 1)$. From (4) and (5),*

$$e_N = E_0 \frac{A_N}{N} + O_{C_\eta^k}(N^{-2}). \quad (11)$$

All linear increments in (7) are N^{-1} times zero-mean oscillatory integrals; the remaining terms are quadratic or higher. Hence uniformly on $J = [X_-, Y_1]$,

$$\sup_X \|\Delta m_N(X, \cdot)\|_{C_\eta^k} \leq C_k N^{-2}. \quad (12)$$

Thus the source's $O(N^{-1})$ bound (8) is non-sharp for the cumulative five moments.

Argument sketch. Substitute (11) into (7). Linear terms use Prop. 2.2; quadratic terms are already $O(N^{-2})$. The functions A, B vanish near the ends of I (lines 2401), so the boundary terms are compatible with the choice of interval. The explicit componentwise constants C_k are not given in this note. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.* $\square$

Proposition 2.4 (Second-order leading term V on the full interval). *[Derived] Because the oscillatory functions vanish near the ends, a smooth zero-mean linear term can be integrated by parts repeatedly, and the full-interval integral decays super-algebraically at any fixed order. Let $\langle \cdot \rangle$ denote the φ -average and integrate over I . Arranged as (M, I, J, S, C_p) , in C_η^k for a fixed background,*

$$\begin{aligned} N^2 \Delta m_N(Y_1, \eta) &\rightarrow V(\eta), \\ V_M &= 0, \\ V_I &= \int \sqrt{2X} E_0 \frac{\langle A^2 \rangle}{2} dX, \\ V_J &= \int \sqrt{2X} E_0 U_0 \frac{\langle A^2 \rangle}{2} dX \\ &\quad + \int \sqrt{2X} E_0 \langle AB \rangle dX, \\ V_S &= \int (\langle B^2 \rangle - E_0^2 \langle A^2 \rangle) dX, \\ V_{C_p} &= \int \frac{E_0^2 \langle A^2 \rangle}{X} dX. \end{aligned} \quad (13)$$

In particular: if $A \not\equiv 0$ for some η , then $V_{C_p} > 0$; if $A \equiv 0$ but $B \not\equiv 0$, then $V_S = \int \langle B^2 \rangle > 0$. For a nontrivial modulation the five-moment vector generally has a genuine second-order mean error, so that $O(N^{-2})$ is sharp as a vector bound for generic modulations.

Argument sketch. Expand the exponential to second order; the cubic remainder integrates to $O(N^{-3})$; the quadratic terms split into a periodic mean and a zero-mean part, the latter integrated by parts again. *Only on the full interval can the linear terms be made super-algebraically small; a partial cumulative (moving endpoint) carries boundary terms and cannot use V as is.* *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.* $\square$

2.3 What the improvement does not cover

Claim 2.5. *[Derived] Prop. 2.3 improves the integrated quantity, not the pointwise profile bound $O(N^{-1})$ or the shear error $O(N^{-1})$ of (6). Subsequent changes of p may still contain the contribution of the profile itself and cannot all be replaced by $O(N^{-2})$.*

Claim 2.6. [Assumption] If the normalization constants are also controlled, one may write $|d_2 - d_1| \leq C_k(\lambda)N^{-2}$. For fixed λ and N large enough this is indeed smaller than λ/N . This is not a two-parameter uniform statement. From this upper bound one may not conclude that “only $\lambda N \gtrsim 1$ suffices”, nor take the scaling $N \sim \lambda^{-2}$ suggested by $\nu_k = O(\lambda^{-1})$ as an established asymptotics of the full construction (the source fixes λ before increasing N (lines 2323–2326, 2542–2545)). *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.*

Remark 2.7 (The λ -factor in a normal combination: observation only). [Assumption] For fixed constants G, c_0 , the second-order leading term of the combination $\Delta J - G\Delta I - c_0\Delta M$ is

$$\int \sqrt{2X} E_0 \left((U_0 - G) \frac{\langle A^2 \rangle}{2} + \langle AB \rangle \right) dX.$$

Whether this expression contains a structural λ -factor is not proven. The source’s repair target is $-\Delta m$, of opposite sign. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.*

3 Note 2: the terminal shell force F_0 , explicit formula and rigid equivalence

3.1 Localization and the definition of the force

[Source] Potential decomposition and exterior heat profile of Theorem 3.1 of [S] (lines 745–790):

$$u = \text{curl } A + B e_\theta, \quad (14)$$

$$|\partial_x^\alpha \partial_t^b R(u, p)| \leq C_{\alpha, b, N, X_1} q^N \quad (0 \leq X \leq X_1, q \downarrow 0), \quad (15)$$

$$X \geq X_{\text{ext}} \Rightarrow A = 0, \quad B = K(r, \tau), \quad p = - \int_r^\infty \frac{K(\rho, \tau)^2}{\rho} d\rho, \quad K(r, \tau) = r^{-1-2h} H_{\text{ext}}(\tau/r^2). \quad (16)$$

Axis growth (lines 791–799):

$$u_\theta(\sqrt{2X_{\text{in}}\tau}, 0, 0, 1 - \tau) = \tau^{-A}(e_0 + O(\tau^{2h})) \quad (\tau \downarrow 0). \quad (17)$$

[Source] Localization of §3.5 of [S] (lines 801–832): take $c = \chi_x \chi_t$, with χ_x axisymmetric and supported in the compact set K , equal to 1 near $(0, 1)$, and χ_t vanishing on $[0, 1 - \tau_0]$ and equal to 1 where $1 - t \leq \tau_0/2$.

$$u = \text{curl}(cA) + cB e_\theta = c u_{\text{loc}} + \nabla c \times A, \quad p = c p_{\text{loc}}. \quad (18)$$

For $t < 1$ put $f = R(u, p)$. The difference $f - cR(u_{\text{loc}}, p_{\text{loc}})$ contains $(\partial_t c - \Delta c)u_{\text{loc}}, p_{\text{loc}} \nabla c$, derivatives and advection of $\nabla c \times A$, and $(c^2 - c)(u_{\text{loc}} \cdot \nabla)u_{\text{loc}}$, supported in the truncation transition region, away from a neighborhood of $(0, 1)$ (lines 815–819).

[Source] Prop. 10.1 of [S] (lines 7229–7258): there is a compact set $K \subset \mathbb{R}^3$ such that $\text{div } u = 0$, $u(\cdot, t) = p(\cdot, t) = 0$ for sufficiently small $t \geq 0$, and (u, p) agrees with $(u_{\text{loc}}, p_{\text{loc}})$ in a spatial neighborhood of the origin for t sufficiently close to 1. Definition of the force (lines 7259–7260):

$$f = R(u, p) = \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla p \quad (0 \leq t < 1). \quad (19)$$

3.2 The terminal value F_0 and the shell formula

[Source] Lemma 10.2 of [S] (lines 7275–7287): for every spatial multi-index α and integer $j \geq 0$, $\partial_x^\alpha \partial_t^j f$ converges uniformly on $\mathbb{R}^3$ as $t \uparrow 1$. There exist $F_j \in C_c^\infty(\mathbb{R}^3; \mathbb{R}^3)$, all supported in K , such that

$$\lim_{t \uparrow 1} \partial_x^\alpha \partial_t^j f(x, t) = \partial_x^\alpha F_j(x), \quad \partial_x^\alpha F_j(0) = 0. \quad (20)$$

Below we write

$$F_0 := F_{j=0} = \lim_{t \uparrow 1} f(\cdot, t) \in C_c^\infty(K; \mathbb{R}^3). \quad (21)$$

[Source] Exterior heat profile (lines 7312–7324): where r is bounded below and q is sufficiently small (so that $X \geq X_{\text{ext}}$),

$$K(r, \tau) = c_\infty s^{-A} H(2\tau/s), \quad s = r^2/2, \quad (22)$$

with $H_{\text{ext}}(Z) = 2^A c_\infty H(4Z)$. For every j ,

$$|\partial_\tau^j K(r, \tau)| \leq C_j r^{-1-2h-2j}, \quad \left| \partial_\tau^j \frac{K(\rho, \tau)^2}{\rho} \right| \leq C_j \rho^{-3-4h-2j}. \quad (23)$$

The untruncated exterior residual vanishes identically (lines 7343–7344).

[Source] Near the origin the truncation equals 1. We use (15) on $X \leq X_{\text{ext}}$, and the residual vanishes identically for larger X ; so for every α, j, N , on a fixed neighborhood of $(0, 1)$ (lines 7349–7356)

$$|\partial_x^\alpha \partial_t^j f(x, t)| \leq C_{\alpha, j, N} (\tau + |z|^{1/D})^N. \quad (24)$$

Hence F_0 vanishes to infinite order at the origin.

Proposition 3.1 (Shell-commutator representation). **[Derived]** Take t sufficiently close to 1 so that $\chi_t \equiv 1$ (i.e. $1 - t \leq \tau_0/2$ (lines 804, 7248–7249)); then $c = \chi_x$. Hence

$$F_0(x) = \lim_{t \uparrow 1} R(\text{curl}(\chi_x A) + \chi_x B e_\theta, \chi_x p_{\text{loc}}). \quad (25)$$

Split by support:

1. The kernel (the neighborhood where $\chi_x \equiv 1$): F_0 is the terminal value of the local residual, flat (zero to infinite order) at 0 by (24).
2. The untruncated exterior ($X \geq X_{\text{ext}}$ with $\chi_x \equiv 1$): the untruncated residual vanishes identically (lines 7343–7344), so $F_0 = 0$.
3. The truncation shell $\text{supp } |D\chi_x|$ (which may meet the time-truncation transition where q is bounded below (lines 821–825)): F_0 equals the terminal value of $f - cR(u_{\text{loc}}, p_{\text{loc}})$, i.e. the commutator listed in [S] (lines 815–818). On the exterior shell, where $A = 0$, this commutator involves only the terminal heat flow $K(\cdot, 0)$, the terminal pressure integral and the fixed cutoff χ_x ; its τ -derivative bounds are controlled by (23).

Argument sketch. For $\chi_t \equiv 1$, (18) and (19) give (25). The three-way split follows from the definition of χ_x (lines 7246–7247), the flatness (24) and the vanishing exterior residual (lines 7343–7344). The componentwise Cartesian expansion of the commutator (including the vanishing of $\nabla \chi_x \times A$ when $A = 0$) is not written out in this note. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.* $\square$

Remark 3.2 (Time extension is not the shortest transient). [\[Source\]](#) Lemma 10.3 of [S] (lines 7384–7395) uses an order-shrinking time truncation to realize

$$f(x, 1 + \sigma) = \sum_{j=0}^{\infty} \chi_0(b_j \sigma) \frac{\sigma^j}{j!} F_j(x), \quad (26)$$

supported in $K \times [0, 2]$. [\[Derived\]](#) This gives the existence of a smooth compactly supported force, not the shortest switching transient. [\[Assumption\]](#) Whether shrinking the χ_t transition width breaks the derivative compatibility of (20) currently has no anchor. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.*

3.3 Rigid equivalence

Proposition 3.3 (Rigid equivalence with the shell data). [\[Derived\]](#) Under the truncation $c = \chi_x \chi_t$ and the local field as in Theorem 3.1 of [S], the following three data determine one another (rigid equivalence):

1. the terminal force $F_0 = \lim_{t \uparrow 1} f$;
2. the full family of Taylor coefficients $\{F_j\}_{j \geq 0}$ of Lemma 10.2 of [S] (via the time compatibility (27));
3. the terminal value of the commutator on the truncation shell (flat-zero kernel + zero untruncated exterior residual + shell commutator).

The equivalence does not depend on non-flat velocity details of the axis kernel: (24) suppresses derivatives of every order near the origin.

[\[Source\]](#) Time compatibility (lines 7367–7378):

$$\partial_x^\alpha \partial_t^j f(x, t) = \partial_x^\alpha F_j(x) - \int_t^1 \partial_x^\alpha \partial_t^{j+1} f(x, s) ds. \quad (27)$$

Claim 3.4 (Whether F_0 encodes j_0 : the exterior factor carries no bias; the fingerprint question is open). [\[Derived\]](#) The bias j_0 appears in the axis velocity $U^* = 4\eta + j_0$ (lines 9003–9004), and does not appear in the exterior heat-flow factor of the shell commutator (25). Flatness (24) suppresses the contribution of the kernel velocity to F_0 at the origin to infinite order (lines 7349–7356). [\[Assumption\]](#) “ F_0 cannot by itself serve as a fingerprint of a symmetric/asymmetric variant” does not follow from the above facts: (10.9) suppresses only a neighborhood of the origin; the kernel region $X < X_{\text{ext}}$ inside $K = \text{supp } \chi_x$ (lines 7244–7247) has positive measure, and the source states only that the residual vanishes identically for $X \geq X_{\text{ext}}$ (lines 779); the dependence of the terminal residual in that kernel on j_0 is not addressed, so the fingerprint question is open. To distinguish variants by the terminal field one would need additional normalized observables on C_τ (e.g. the axis value $U(0, 0)$, the e_0 of (17), or $E(X_*, 0)$ at $\eta = 0$), and a proof that the high-frequency corrections of §5 and §9 of [S] do not erase that observable. This step is not done. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.*

Remark 3.5 (Interior points in a bounded domain). [\[Source\]](#) Placing the localized F_0 in a bounded smooth domain with 0 in the interior is the trivial restriction of Prop. 10.1 of [S] (lines 7229–7232): the domain need only contain K . This note does not regard that restriction as a nontrivial boundary blowup.

4 Note 3: the load-bearing role of the bias j_0 — a parity mechanism

4.1 Setup and the source endpoint test

This section fixes [\[Assumption\]](#)(the same test point as the second review round): $j_0 = 0$, $U^* = 4\eta$, endpoint $(X, \eta) = (4/\Lambda, 0)$, $Y = \Lambda X = 4$.

[\[Source\]](#)Axis data (lines 8998–9009):

$$U^* = 4\eta + j_0, \quad H^* = D\eta + dU^*, \quad W^* = 1 - dU_\eta^* - 2D\eta U^*, \quad (28)$$

with

$$-W^* = 3 - 8h\eta^2 + (1 - 2h)j_0\eta.$$

[\[Derived\]](#)Substituting $j_0 = 0$, $\eta = 0$ into (28) gives $W^*(0) = -3$, $H^*(0) = 0$, hence $\eta_0 = 0$ (lines 9004–9008). [\[Source\]](#) Π_0 is even(lines 8998–9000). The dichotomy (29) (lines 9014–9019):

$$\begin{aligned} &\text{first choose } \delta_* > 0 \text{ so that } \{|Z^*| \leq \delta_*\} \text{ avoids a neighborhood of } \eta_0, \\ &\text{then choose } \sigma_* \text{ so that } \chi > .99 \text{ on } \{|Z^*| \leq \delta_*\}. \end{aligned} \quad (29)$$

The amplitude C is chosen after Λ (lines 9037–9038,9215–9218); Lemma B.4 of [S] “first Λ , then C , sufficiently large”(lines 9383); §B.7 of [S]: increasing C moves X_R outward without lengthening the transition(lines 9580–9586). The absence of an upper bound is thus pinned. Azimuthal data (lines 9027–9036):

$$\zeta^* = -\frac{LH^*}{H^{*2} + \sigma_*^2}, \quad \xi_0 = \Lambda\zeta^*, \quad \varphi^* = \exp\left(\Lambda \int_0^\eta \zeta^*(w) dw\right). \quad (30)$$

Parameter order (lines 9778–9779):

$$\begin{aligned} M_d, T_d, P_*, \lambda, h &\rightarrow \text{matching tolerance, } j_0 \rightarrow \delta_*, \sigma_*, \Lambda \\ &\rightarrow (B_k), T_{\text{sh}} \rightarrow C, X_R \rightarrow \kappa_0, t_1, \text{final widths.} \end{aligned} \quad (31)$$

[\[Source\]](#)Endpoint test of Prop. B.3 of [S] (lines 9281–9285,9353): at $X_0 = 4/\Lambda$ there is a fixed $c_{\text{ex}} > 0$ (the proof takes $c_{\text{ex}} = .2$) such that

$$p_1 + \frac{p_2^2}{p_1} > 2 + c_{\text{ex}}. \quad (32)$$

The large- C mechanism requires $\chi \leq .99$ and $|Z^*| > \delta_*$, hence $|n_s| \geq \delta_*/2$,

$$|p_2| = C\sqrt{\frac{2}{\Lambda}} \frac{|n_s|}{\varphi^*\Phi} \quad (\text{lines 9344–9349}). \quad (33)$$

On the complementary region $|Z^*| > \delta_*$ one has $|n_s| \geq \delta_*/2$, so this term carries C ; that region does not contain $Z^*(0) = 0$ (lines 9344–9352).

4.2 Parity: $p_2(\cdot, 0) \equiv 0$ exactly

Proposition 4.1 (On the axis at $j_0 = 0$, $p_2 \equiv 0$). [\[Derived\]](#)At $j_0 = 0$, U^* is odd and Π_0 is even(lines 8998–9004) $\Rightarrow Z^*$ is odd $\Rightarrow Z^*(0) = 0$; χ is even, φ^* is even ((30)). The comparison profile of Prop. B.2 of [S] $\Phi_0 = f_0(Y\chi)$ is even, and $u_0 = -YZ^*/(2L)$ is odd(lines 9236). The remainders R_1 is even and R_2 is odd (substitute each term of (lines 9197–9199)). The fixed-point map (lines 9237–9243) preserves this parity. The real fixed point is unique(lines 9261) $\Rightarrow$ the exact solution satisfies $u(Y, 0) = 0$, hence $u_Y(Y, 0) = 0$. [\[Source\]](#) $n_s = -2u_Y$ (lines 9324), so $n_s(Y, 0) = 0$, hence $p_2(4/\Lambda, 0) = 0$. Thus $p_2^2/p_1 = 0$.

Argument sketch. Check the parity of each term of (lines 9197–9199) under $\eta \mapsto -\eta$; the map (lines 9237–9243) preserves the class; uniqueness (lines 9261) forces the exact solution into that class. [Source] “Real data and uniqueness of the fixed point give a real solution” is the statement of (lines 9261); this note quotes it as is and does not isolate it as a separate lemma; the full term-by-term parity verification is not written out. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.* $\square$

Remark 4.2 (A forbidden generic claim). [Derived] One must not write $p_2^2/p_1 \asymp C^2\Lambda^{-1}$ at $\eta = 0$: that asymptotics applies only on the complementary region $|n_s| \geq \delta_*/2$ (lines 9344–9352), which does not contain $Z^*(0) = 0$. The missing step is the jump from (33) to $|n_s| = \Theta(1)$. Even without parity, citing only (B.13) (lines 9172–9182) via (B.18) (lines 9278), at $\eta = 0$ one has $n_s = O(\Lambda^{-1})$ and $p_1 = O(\Lambda^{-1})$ ((lines 9325–9326): the upper bound for p_1 and the approximation for n_s follow from these), so $p_2 \asymp C\Lambda^{-3/2}$, $p_2^2/p_1 \asymp C^2\Lambda^{-2}$, still not Λ^{-1} .

4.3 $p_1 = 2(3 - h)/\Lambda$ and the degeneration of (B.19)

Proposition 4.3 (On-axis $R_1^{(0)}$ and $\partial_Y \Phi_1$). [Derived] At $j_0 = 0$: $\chi(0) = H^*(0) = \zeta^*(0) = Z^*(0) = 0$, $u_0(Y, 0) = 0$, $\Phi_0(Y, 0) = f_0(0) = 1$ (lines 9150–9160). $L(0) = 1$, $W^*(0) = -3$ (lines 9005, 9008), so

$$R_1^{(0)}(Y, 0) = W^*(0) + h = -3 + h, \quad (34)$$

independent of σ_*, δ_*, Y . At $\eta = 0$, $T = J_2\chi/2 = 0$, so $2(Y(\Phi_1)_{YY} + 2(\Phi_1)_Y) = h - 3$. Smoothness at $Y = 0$ excludes the Y^{-2} homogeneous solution, giving

$$\partial_Y \Phi_1(4, 0) = \frac{h - 3}{4} < 0 \quad (0 < h \leq 10^{-2}, \text{ (lines 9002, 1344–1347)}). \quad (35)$$

The remainder $O(\Lambda^{-2})$ does not change the sign for Λ large.

Argument sketch. The fixed-point expansion $\Phi = \Phi_0 + \Lambda^{-1}\Phi_1 + O(\Lambda^{-2})$ comes from the same contraction map (lines 9237–9247), not from an external fit. Cross-check: on the axis $S_q = -W - h = 3 - h$ (lines 1464), consistent with $R_1 = h - 3$. ODE: $Yv' + 2v = (h - 3)/2$ with $v(0)$ finite $\Rightarrow v \equiv (h - 3)/4$. The consistency estimate of the expansion remainder is not written out. *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.* $\square$

Proposition 4.4 ($p_1(4/\Lambda, 0)$). [Source] $p_1 = a = -2Y\Phi_Y/\Phi$ (lines 9324). [Derived] $\Phi_Y(Y, 0) = \Lambda^{-1}(h - 3)/4 + O(\Lambda^{-2})$, $\Phi(Y, 0) = 1 + O(\Lambda^{-1})$, so

$$p_1\left(\frac{4}{\Lambda}, 0\right) = \frac{2(3 - h)}{\Lambda} + O(\Lambda^{-2}) > 0 \quad (\Lambda \text{ sufficiently large}). \quad (36)$$

Cross-check: on the axis $Q_s(0) = S_q(0)/2$ (lines 1487), $S_q(0) \approx 3 - h$, $p_1 = XQ_s/L \approx (4/\Lambda)(3 - h)/2 = 2(3 - h)/\Lambda$.

Proposition 4.5 ((B.19) degenerates at $\eta = 0$). [Derived] By Prop. 4.1, $p_2(\cdot, 0) \equiv 0$, so C does not enter (32). (32) degenerates to

$$p_1 > 2 + c_{\text{ex}} = 2.2. \quad (37)$$

By (36), $p_1 = 2(3 - h)/\Lambda \not> 2.2$ (Λ large). The large C multiplies p_2 (lines 9346–9349) and is ineffective against $p_2 = 0$.

Claim 4.6 (The two cones are separate). [Derived] First cone inequality $a - b_s w > 0$: $b_s = 2D_X U/E$ (lines 1494–1496) vanishes at $\eta = 0$ (U odd), $a = p_1 > 0$, so this item passes. What fails is the second cone test $v_s = p_1 + p_2^2/p_1 > 2$. The two must not be conflated into one “cone”. The literal conditions $p_1 > 0 / \partial_Y \Phi_1 < 0$ pass, and hold for all σ_*, δ_* ; they are not a necessary and sufficient condition for large C to rescue (32) (one still needs $n_s \neq 0$).

Remark 4.7 (The quantifiers of (B.2)). [\[Source\]](#) The quantifiers of (B.2): first choose $\delta_* > 0$ so that $\{|Z^*| \leq \delta_*\}$ avoids a neighborhood of η_0 , then choose σ_* so that $\chi > .99$ (lines 9014–9019). [\[Derived\]](#) Substituting $j_0 = 0$ (an external limit point; the source allows only $0 < j_0 \leq .05$ (lines 9002)): $\eta_0 = 0$ and $Z^*(0) = 0$ ((lines 9004–9008) at $\eta = 0$), so for every $\delta_* > 0$ the point $\eta = 0$ lies both in $\{|Z^*| \leq \delta_*\}$ and in any neighborhood of η_0 . (29) has no solution under the quantifiers of this dichotomy. This is not a choice problem repairable by making σ_*, δ_* smaller/larger. This statement stays at level L0 (this dichotomy fails at the external limit point $j_0 = 0$, see Claim 4.8) and is not extended to an existence statement under other dichotomies.

4.4 The L0 conclusion and the in-class rigidity

Claim 4.8 (L0 (the device extended to the external limit point $j_0 = 0$)). [\[Derived\]](#) $j_0 = 0$ lies outside the source’s domain (the source allows only $0 < j_0 \leq .05$ (lines 9002); this note extends the device to that external limit point): the unique real fixed point lies on the $\eta \mapsto -\eta$ symmetry, so $p_2(\cdot, 0) \equiv 0$ and $b_s(\cdot, 0) = 0$; (32) degenerates at $\eta = 0$ to $p_1 > 2.2$, while $p_1 = 2(3-h)/\Lambda \rightarrow 0$. Hence the endpoint test (32) of Prop. B.3 of [S] fails at that extended point. This is L0: “the device fails when extended to $j_0 = 0$ ”; the nonexistence of a $j_0 = 0$ variant within one class (Prop. 4.9) is L1, and this note does not claim that the source itself fails at some step with $j_0 > 0$.

The necessity at X_0 . [\[Source\]](#) Lemma B.4 of [S] requires $v_r > 2 + c$ on the whole interval $[X_0, X_i]$ (lines 9383–9389); at $\eta = 0$, $v_r = p_{1,r}$. The analytic axis region extends only to $Y \leq 4.1$, i.e. $X \leq 4.1/\Lambda \rightarrow 0$ (lines 9166–9167, 9366). The first collar, where $v_s > 2$, is produced by continuity from $v_r(0) > 2 + c$ (lines 9503–9506). [\[Derived\]](#) The following proposition records ruling (i): L1 (no $j_0 = 0$ variant within the class), not L2.

Proposition 4.9 (In-class rigidity (L1)). *Assume the axis device satisfies the following class definition (H1)–(H3):*

- (H1) *The axis-data class of Appendix B of [S] ((B.1)–(B.3), (lines 9003–9036), $U^* = 4\eta + j_0$), the analytic axis profile defined on $0 \leq Y = \Lambda X \leq 4.1$ (lines 9166–9167), with vanishing leading residual stress on the axis region ($T_0 = 0$, (lines 1904–1905, 2111)).*
- (H2) *Stress activated at $X_0 = 4/\Lambda$: the statement of Prop. B.5 of [S] ((lines 9462–9464), continued to X_i with $X \leq X_0$ unchanged), (B.26) ((lines 9476–9479), the prescription on $0 < y < t_1$), and $B_0(0, \eta) \neq 0$ ((lines 9523–9524)).*
- (H3) *The same cone test: (32) of Prop. B.3 of [S] ((lines 9281–9285), $p_1 + p_2^2/p_1 > 2 + c_{\text{ex}}$) and the $v_r > 2 + c$ of Lemma B.4 of [S] on $[X_0, X_i] \times [-1, 1]$ ((lines 9384–9389)), with the first collar given by the continuity of (lines 9505–9506), uniformly in η .*

[\[Derived\]](#) Then $j_0 = 0$ fails the cone test within this class. Concretely, at $(X, \eta) = (4/\Lambda, 0)$:

1. At $j_0 = 0$, U^* is odd and Π_0 is even (lines 8998–9004) $\Rightarrow Z^*$ odd, χ, φ^* even, the comparison profile $\Phi_0 = f_0(Y\chi)$ even, $u_0 = -YZ^*/(2L)$ odd (lines 9236); the remainders R_1 even and R_2 odd (lines 9197–9199); the fixed-point map (lines 9237–9243) preserves this parity; the real fixed point is unique (lines 9261) $\Rightarrow$ the exact solution has $u(Y, 0) \equiv 0 \Rightarrow n_s = -2u_Y = 0$ (lines 9324) $\Rightarrow p_2(4/\Lambda, 0) = 0$ ((lines 9277, 9346–9349)).
2. $R_1^{\text{lead}}(Y, 0) = W^*(0) + h = -3 + h$ ((lines 9005, 9008) at $j_0 = 0, \eta = 0$; the LR_1 of (B.14) evaluated at $\eta = 0$ (lines 9197); $L(0) = d(0) = 1$), independent of σ_*, δ_*, Y ; the axis-regular solution of the slice ODE $Yv' + 2v = (h-3)/2$ gives $\partial_Y \Phi_1(4, 0) = (h-3)/4 < 0$; hence by Prop. 4.4, $p_1(4/\Lambda, 0) = 2(3-h)/\Lambda + O(\Lambda^{-2}) \rightarrow 0$ (cross-check: the $Q_s(0) = S_q(0)/2$ of (lines 1487) and the $S_q = -W - h = 3 - h$ of (lines 1464)).

3. Hence (32) degenerates at $\eta = 0$ to $p_1 > 2.2$ and fails (Prop. 4.5). The first cone inequality $a - b_s w > 0$ passes ($b_s = 2D_X U/E$ vanishes at $\eta = 0$, (lines 1494–1496)); what dies is the second cone test $v_s > 2$ (Claim 4.6).

Evidence for the negative statement (“must start at X_0 ”). [\[Source\]](#)

1. Prop. B.5 of [S] (lines 9499–9500): for t_1 sufficiently small, $P_c > 2 + c$; by (B.29) ((lines 9494–9498)), $P_c = v_r + O(ye_a)$, so at $y = 0$, $v_r(X_0, \eta) > 2 + c$. Shrinking t_1 only shrinks the error term and cannot raise the endpoint where $v_r(0) < 2$.
2. Prop. B.5 of [S] (lines 9505–9506): “Continuity from $v_r(0) > 2 + c$ gives one positive-width first collar on which $v_s > 2 + c/2$, uniformly in η ” — this is the only sentence in the whole source that names $v_r > 2 + c$ at $y = 0$. (B.26) ((lines 9476–9479)) *does not* use $v_r > 2 + c$.

Where the chain breaks, and downstream uses. [\[Derived\]](#) If the endpoint lower bound is deleted, then at $\eta = 0$: the η -uniform $v_s > 2$ of the first collar cannot start (the quantifier of (lines 9505–9506)) $\rightarrow$ the near-endpoint $P_c > 2$ fails $\rightarrow$ both the admissible and the relaxed branches fail simultaneously (both branches of the dichotomy of (lines 9503–9505) break) $\rightarrow$ while T_0 is already nonzero at $X_0 +$ ((lines 9523–9524)) $\rightarrow$ Cor. B.6 of [S] ((lines 9563–9564)). This is the load-bearing chain. Downstream uses (links in the full existence theorem, *not* independent steps of this proposition): Prop. 4.10(ii) of [S] ((lines 2112–2129)), Appendix C of [S] ((lines 9809–9811)), Theorem 4.6(iii) of [S] ((lines 1908–1909)).

Exclusion of candidate gaps. [\[Source\]](#)

- The $v_s \leq 2$ branch of (lines 9503–9505) still requires $P_c > 2$;
- the $v_s < 1$ of (lines 9538–9544): its onset is after the first collar, still on the scale $Y = O(1)$, and the branch follows the reference extension to X_b (and on to X_i , (lines 9366, 9540));
- the vacuum region $T_0 = 0$ ends exactly at X_0 ((lines 1904–1905, 2111));
- the $4e^{t_c} < 4.1$ of Cor. B.6 of [S] ((lines 9558)) locks the first collar within the analytic axis scale, so it cannot be moved out to $X \asymp 1$.

Scope: this proposition is L1 (no $j_0 = 0$ variant within one function class; $j_0 = 0$ is an external limit point outside the source’s domain, the source allowing only $0 < j_0 \leq .05$ (lines 9002)), valid only within the class (H1)–(H3); it is not L2 (it does not concern “nonexistence of reflection-symmetric forced blowup”). Outward statements use L0 only.

Remark 4.10 (No assertion outside the class). [\[Source\]](#) This proposition covers only the class (H1)–(H3), and makes no assertion outside it. Class-external modifications may change the conclusion, e.g.: dropping “the axis profile is unchanged for $X \leq X_0$ ” (lines 9460–9461) or “ $T_0 = 0$ on the axis region” (lines 1904–1905, 2111); moving the activation point from X_0 into the interior of $(X_0, X_i]$ — Prop. B.5 of [S] pins only the strict relaxed cone on the rest of $(X_0, X_i]$ (lines 9461–9464), and this proposition does not rule out other activation geometries (the $B_0(0, \eta) \neq 0$ of (H2), (lines 9523–9524), is precisely the anchor that pins the activation point); or changing the cone test. For all such class-external cases this proposition is silent; this is exactly why it is not written as L2.

Remark 4.11 (Matching window: a single-point check, not a conclusion of this note). [\[Derived\]](#) At the single point $(4/\Lambda, 0)$: $H^*(0) = j_0$, $Z^*(0) = -(\frac{9}{2} + h)j_0$ exactly (lines 9003–9005); $p_2 = -C\sqrt{2/\Lambda}(\frac{9}{2} + h)j_0(1 + \dots)$ (lines 9346–9349); $p_1 = 2(3 - h)/\Lambda + 2\chi(0) + \dots$ ((lines 9324) together with the S_q and χ terms (lines 1487, 9008, 9299)); hence $p_2^2/p_1 = C^2 k^2 j_0^2 / [(3 - h) + \Lambda\chi(0)]$, and when χ is subdominant the Λ cancels. [\[Assumption\]](#) Regime $j_0 = o(\sigma_*)$: σ_* keeps a positive

lower bound independent of j_0 — the source lines 9014–9015 and 9020–9023 state only “ σ_* may be taken sufficiently small” and “ Ω exists”, and do not state that the holomorphic-neighborhood width does not collapse as $j_0 \rightarrow 0$; this positive lower bound is an additional hypothesis with no source basis. **[Assumption]** The main case additionally requires $C^2 \gg \Lambda/\sigma_*^2$ (the χ term does not dominate p_1) — likewise an additional hypothesis with no source basis: the source lines 9014–9015, 9020–9023 state only that σ_* may be taken sufficiently small and that Ω exists. **[Derived]** The lower bound for $p_2^2/p_1 > 2.3$ is $\Theta(1/C)$ (the coarse bound $p_1 \leq C_1$ loses the Λ^{-1} gain at this point — that upper bound is true in itself; using it to estimate the ratio only yields the overly strong $\sqrt{\Lambda}/C$). The matching bound (lines 2210–2213): $\|G_i - 4\eta\|_{C^k} \leq C_k j_0 + C_k^{\text{nat}}/\Lambda + \dots$; requiring this bound to be smaller than ε_m (lines 2190) gives an upper bound on j_0 of order $\Theta(1/C_k)$. **[Source]** C has no upper bound: (B.16) ((lines 9215–9218)) gives only a lower bound; C may continue to increase ((lines 9350–9352, 9583–9585)). **[Derived]** Hence the $\Theta(1/C)$ lower bound “barely bites”: increasing C can push the lower bound arbitrarily small; the only point where the test truly collapses is the exact value $j_0 = 0$ (the parity mechanism of Prop. 4.9). Ruling (a conclusion of this note): merely “the matching upper bound and the *single-point* cone test do not conflict under the parameter order of (31) (freeze j_0 first, then choose $\delta_*, \sigma_*, \Lambda, C$)”. We do not write: a quantitative window theorem; nor “any $j_0 > 0$ is rescued by large C ” (the η -neighborhood, the worst point of the (B.2) complementary region, and the whole interval $[X_0, X_i]$ of Lemma B.4 of [S] are not covered). *This note does not prove this step; it is stated only as an observation and does not constitute a complete proof.*

5 Acknowledgments and statement

Acknowledgments. The author (Kaiyan Ren) is an M.Sc. student in the Department of Computer Science and Engineering, School of Engineering, The Hong Kong University of Science and Technology, and was formerly an undergraduate in the School of Mathematics, Nanjing University. Gratitude is due to several independent review sessions for their line-by-line scrutiny of the line anchors, the color-code and the scope boundaries: every **[Source]** statement of this note was, on that account, re-checked verbatim against the extracted text, and several wordings and anchors were corrected. The review records themselves do not constitute evidence for this note; the only evidence is the source line numbers.

Statement on the use of artificial-intelligence tools. This note was written with the assistance of generative-AI tools: DeepSeek (line-number extraction, formula rewriting, parity computations and the orchestration of the review rounds), Claude (independent review and the packaging of the release draft), ChatGPT (earlier finite-dimensional notes and material preparation). These tools are not authors of this note. All assertions are anchored to line numbers of the reference text extraction `navier-stokes.txt` identified in §1; the reader should check them line by line, and tool output is not evidence. The author takes full responsibility for the entire content of this note, including line-number and scope errors.

Scope and disclaimer. This note records three structural observations; it is not a completed mathematical paper and has not been peer-reviewed. Prop. 2.2–2.4, 3.1–3.3, 4.1–4.5 give only argument sketches, each explicitly marked as not constituting a complete proof; steps tagged **[Assumption]** are unproven. The in-class rigidity Prop. 4.9 and the single-point check Rem. 4.11 are valid only within the class definition (H1)–(H3) and the stated single-point check, respectively. This note passes no judgment on the correctness of the source construction: the whole text reads the source under the hypothesis “the construction is correct as claimed”, and the L0 statement concerns only the extension of the device to $j_0 = 0$ (an external limit point outside the source’s domain). **No new Navier–Stokes solution is claimed**, no new regularity criterion is claimed, the existence or nonexistence of reflection-symmetric forced blowup is

not claimed, and the finite-dimensional five-moment observation or the shell representation of the terminal force F_0 is not promoted to a uniqueness theorem for the full construction.

References

- [1] OpenAI, *Finite Time Blowup for Navier–Stokes* (these words are the title line of the extraction, (lines 2–3); abstract at (lines 4–6), Theorem 1.1 of the source at (lines 28–37)), made public September 8, 2026. Announcement and full-text entry: <https://openai.com/index/navier-stokes-solution/>. (As of the finalization of this note the paper is not on arXiv, so it is cited via its official publication page, without an arXiv number.) All line numbers of this note refer to the reference text extraction `navier-stokes.txt` of that PDF (10264 lines; fingerprint and availability in the reproducibility paragraph of §1).
- [2] OpenAI, *Lean certificates accompanying Navier–Stokes and Euler results*, GitHub repository <https://github.com/openai/NavierStokesAndEuler>, fixed commit 8937a8f4cbc7abaab5e9e97d1cc7f5d2319d9538 (this note does not cite the Lean proofs).