

NST IB Mathematical Methods II

Yu Lu

Lent, 2023

1 Sturm-Liouville Theory

1.1 The operator

A Sturm-Liouville operator

$$\mathcal{L} = -\frac{d}{dx} \left(\rho(x) \frac{d}{dx} \right) + \sigma(x), \quad \rho(x) > 0, \sigma(x) \in \mathbb{R}$$

can be shown to be self-adjoint under appropriate boundary conditions

$$\rho(vu^{*'} - u^*v') \Big|_{\alpha}^{\beta} = 0,$$

meaning $\langle u|\mathcal{L}v \rangle = \langle \mathcal{L}u|v \rangle$ (equivalently, $\mathcal{L}^{\dagger} = \mathcal{L}$).

Such self-adjointness provides many convenient features. Just like Hermitian matrices, for example, in the eigenvalue equation for a self-adjoint operator $\mathcal{L}y_n = \lambda_n y_n$, the eigenvalues λ_n are real and **non-degenerate** eigenfunctions y_n are mutually orthogonal. Moreover, the eigenfunctions commonly form a complete basis, from which we could solve inhomogeneous linear differential equations.

1.2 A generalisation to any 2nd order linear ODE

Despite being seemingly restrictive, the Sturm-Liouville operator could express any second order linear differential operator ($a = -p, b = p' - q$)

$$\tilde{\mathcal{L}} = p(x) \frac{d^2}{dx^2} + q(x) \frac{d}{dx} + r(x) = -\frac{d}{dx} \left(a(x) \frac{d}{dx} \right) - b(x) \frac{d}{dx} + r(x)$$

could be converted into a Sturm-Liouville form by choosing $w(x)$ such that

$$w\tilde{\mathcal{L}} = -\frac{d}{dx} \left(aw \frac{d}{dx} \right) + (aw' - bw) \frac{d}{dx} + rw$$

has a vanishing coefficient for the first derivative term:

$$w(x) = C \exp \left[\int^x \frac{b(x')}{a(x')} dx' \right] > 0.$$

This way, a Sturm-Liouville operator

$$\mathcal{L} = w(x)\tilde{\mathcal{L}} = -\frac{d}{dx} \left(\rho(x) \frac{d}{dx} \right) + \sigma(x), \quad \rho = aw, \sigma = rw$$

could be constructed. From this, an eigenvalue problem $\tilde{\mathcal{L}}y_n = \lambda y_n$ involving $\tilde{\mathcal{L}}$ could be converted to a generalised eigenvalue problem $\mathcal{L}y_n = \lambda w(x)y_n$ of the Sturm-Liouville operator \mathcal{L} . Alternatively, we could stick with the original operator $\tilde{\mathcal{L}}$ and introduce the weight in the definition of inner product such that

$$\langle u|v \rangle_w = \int_{\alpha}^{\beta} w^* uv \, dx \implies \|v\|_m^2 = \int_{\alpha}^{\beta} w^* |v|^2 \, dx,$$

and the self-adjoint condition becomes $\langle u|\mathcal{L}v \rangle_w = \langle \mathcal{L}u|v \rangle_w$. Under this convention, $\tilde{\mathcal{L}}$ satisfying the original eigenvalue equation is self-adjoint with weight $w(x)$ while \mathcal{L} satisfying the generalised eigenvalue equation is self-adjoint with weight 1.

2 Laplace's and Poisson's equation

Many physical systems, including incompressible irrotational fluids, heat conduction, and electrostatics, can be described by Poisson's equation

$$\nabla^2 \Phi(\mathbf{r}) = \rho(\mathbf{r}), \quad (1)$$

where $\Phi(\mathbf{r})$ is defined in some region V with **Dirichlet** boundary conditions $\Phi(\mathbf{r}) = \Psi(\mathbf{r})$ or **Neumann** boundary conditions $\nabla \Phi \cdot \hat{n} = \Psi(\mathbf{r})$ on ∂V . Sometimes Φ is itself an observable, like temperature, while sometimes it is only $\nabla \Phi$ that is physical, like in electrostatics. The distinction sometimes brings about a difference in boundary conditions. We are sometimes (like in heat conduction) also interested in the flux $\mathbf{F} = \kappa \nabla \Phi$, where κ is (piecewise) constant.

2.1 Laplace's equation

When $\rho(\mathbf{r}) = 0$ in Eq.(1), it simplifies into Laplace's equation

$$\nabla^2 \Phi(\mathbf{r}) = 0,$$

where $\Phi(\mathbf{r})$ satisfies the same types of boundary conditions as described above. We could express $\Phi(\mathbf{r})$ as a superposition of separable solutions in cartesian coordinates, planar polar, or spherical polar coordinates.

2.1.1 Cartesian coordinates

Separation into cartesian coordinates gives $\Phi(x, y, z) = X(x)Y(y)Z(z)$ and the Laplace's equation reads

$$\begin{aligned} \frac{X''(x)}{X} + \frac{Y''(y)}{Y} + \frac{Z''(z)}{Z} &= 0, \\ \Rightarrow \frac{d^2 X}{dx^2} &= -k_x^2 X, \quad \frac{d^2 Y}{dy^2} = -k_y^2 Y, \quad \frac{d^2 Z}{dz^2} = (k_x^2 + k_y^2) Z. \end{aligned}$$

Therefore, one part of the solution must be qualitatively different from the other two. The most general solution is

$$X = A \cos(k_x x + B) \quad Y = C \cos(k_y y + D), \quad Z = E \exp\left(\sqrt{k_x^2 + k_y^2} z\right) + F \exp\left(-\sqrt{k_x^2 + k_y^2} z\right),$$

but the boundary conditions often immensely simplifies the expression. For example, k_x and k_y may be quantised to integer multiplies of some quantities and $B = D = 0$ if oscillatory boundary conditions are given in X and Y . Furthermore, the exponential growth solution in Z are often killed due to regularity constraint at infinity.

2.1.2 Planar polar coordinates

In planar polar coordinates, the solution can be separated as $\Phi(r, \phi) = P(\phi)R(r)$ and the Laplace's equation reads

$$\nabla^2 \Phi = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \Phi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \Phi}{\partial \phi^2} = 0,$$

giving the separated ODE as

$$\frac{d^2 P}{d\phi^2} + n^2 P = 0, \quad r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} - n^2 R = 0,$$

where $n = 0, 1, 2, \dots$ from the physical requirement on periodicity of P or P' : if Φ is physically meaningful, then we require $P(\phi + 2\pi) = P(\phi)$ and if only $\nabla \Phi$ is physical, only P' needs to be periodic. The ODE on P gives oscillatory solution while that on R is of Euler's type and gives a power function solution $R = r^{\pm n}$ or $R = \ln r$. Thus, the general solution is

$$\Phi(r, \phi) = A_0 + B_0 \phi + C_0 \ln r + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} r^n (A_n \cos(n\phi) + B_n \sin(n\phi)).$$

The $B_0\phi$ term is only present when the physical quantity is $\nabla\Phi$. Regularity condition at $r = 0$ kills $\ln r$ and all the r^{-n} while regularity at infinity kills all the r^{-n} dependence (and $\ln r$ if it is posted on Φ itself). Specifying one more boundary condition then gives A_n and B_n in terms of Fourier coefficients.

The only solution to the Laplace's equation in all space is a constant function, which is consistent with Liouville's theorem in complex analysis (though this is true in 3D as well). Therefore, we usually study the behaviour parts of the space (e.g. a circular disc). Alternatively, we could allow $\hat{n} \cdot \nabla\Phi$ to have a jump discontinuity across some boundaries and split the solution into two piecewise parts. This may correspond to a point source or a discontinuity in the conductivity κ (conserving flux \mathbf{F}).

2.1.3 Spherical polar coordinates

Laplace's equation in axisymmetric spherical polar coordinates (no ϕ dependence) is

$$\nabla^2\Phi = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial\Phi}{\partial r} \right) + \frac{1}{r^2 \sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial\Phi}{\partial\theta} \right) = 0,$$

where $\Phi(r, \theta) = \Theta(u)R(r)$ and $u = \cos\theta$. The equation can then be separated as

$$\frac{d}{du} \left[(1-u^2) \frac{d\Theta}{du} \right] + l(l+1)\Theta = 0, \quad r^2 \frac{d^2R}{dr^2} + 2r \frac{dR}{dr} - l(l+1)R = 0.$$

The ODE on Θ is of the Legendre type and is only regular at $u = \pm 1$ when $l = 0, 1, \dots$. The second ODE is of Euler's type and admits power function solutions $R = r^l, r^{-l-1}$. Thus, the general solution is

$$\Phi(r, \theta) = \sum_{l=0}^{\infty} (A_l r^l + B_l r^{-l-1}) P_l(\cos\theta),$$

where $P_l(x)$ is the Legendre polynomial with normalisation $P_l(1) = 1$.

(*) When the system is not axisymmetric, the Laplace's equation becomes

$$\nabla^2\Phi = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial\Phi}{\partial r} \right) + \frac{1}{r^2 \sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial\Phi}{\partial\theta} \right) + \frac{1}{r^2 \sin\theta} \frac{\partial^2\Phi}{\partial\phi^2} = 0.$$

It needs to be separated into $\Phi(r, \theta, \phi) = R(r)\Theta(\theta)P(\phi)$ with two separation constants, as shown below:

$$\begin{cases} \frac{\sin^2\theta}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) + \frac{\sin\theta}{\Theta} \frac{d}{d\theta} \left(\sin\theta \frac{d\Theta}{d\theta} \right) = -\frac{1}{P} \frac{d^2P}{d\phi^2} = \lambda_1, \\ \frac{1}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) = -\frac{1}{\Theta \sin\theta} \frac{d}{d\theta} \left(\sin\theta \frac{d\Theta}{d\theta} \right) + \frac{\lambda_1}{\sin^2\theta} = \lambda_2. \end{cases}$$

2.2 Uniqueness of solutions to Poisson's equation

The uniqueness theorem states that if $\nabla^2\Phi = \rho(\mathbf{r})$ in V and Dirichlet or Neumann boundary conditions are satisfied on ∂V , then the solution must be unique, or unique up to a constant for Neumann conditions. This can be proved by showing that $\Psi = \Phi_2 - \Phi_1$ satisfies

$$\int_V |\nabla\Psi|^2 dV = \oint_{\partial V} \Psi \nabla\Psi \cdot d\mathbf{S} = 0$$

using the vector identity $\nabla \cdot (\Psi \nabla\Psi) = |\nabla\Psi|^2 + \Psi \nabla^2\Psi$.

This allows us to use the method of image to convert Poisson's equation in a confined space to a Poisson's equation over all space.

2.3 Green's function to a point source

Green's function to the Poisson's equation in V bounded by S is

$$\nabla_r^2 G(\mathbf{r}, \mathbf{r}') = \delta^{(3)}(\mathbf{r} - \mathbf{r}')$$

satisfying homogeneous $\boxed{G = 0 \text{ on } S}$ for a Dirichlet problem or $\boxed{\nabla \cdot G = 1/A \text{ on } S}$ for a Neumann problem, where A is surface area of S and \mathbf{r}' can be any point inside V . It is not hard to verify that $G(\mathbf{r}, \mathbf{r}')$ is symmetric under $\mathbf{r} \leftrightarrow \mathbf{r}'$. The condition on Neumann problem comes from conserving the flux

$$\oint_S \nabla G \cdot d\mathbf{S} = \int_V \nabla^2 G dV = \int_V \delta^{(3)}(\mathbf{r} - \mathbf{r}') dV = 1.$$

2.3.1 3D fundamental solution

A fundamental solution $G(\mathbf{r}, \mathbf{r}')$ is the Green's function satisfying the boundary conditions in the whole space. Just as what we did in finding the Green's function of an ODE, G can be obtained by solving the Laplace's equation but allowing for discontinuity with determined strength at the origin.

Since the proposed PDE is radially symmetric, we would expect G to only have radial dependence, giving

$$\nabla^2 G = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial G}{\partial r} \right) = 0 \implies G = \frac{C}{r} + A \quad (r \neq 0),$$

where A, C are arbitrary constants. The constant offset A must be 0 for a Dirichlet problem but can be left undetermined for a Neumann problem. The strength C can be determined by the flux out from the origin:

$$\int_{r < \epsilon} \nabla^2 G dV = \oint_{r=\epsilon} \frac{\partial G}{\partial r} dS = 1,$$

giving (shifting the origin)

$$\boxed{G = -\frac{1}{4\pi |\mathbf{r} - \mathbf{r}'|}}.$$

2.3.2 2D fundamental solution

Similarly, the fundamental solution to the 2D Poisson's equation

$$\nabla_r^2 G(\mathbf{r}, \mathbf{r}') = \delta^{(2)}(\mathbf{r} - \mathbf{r}')$$

is

$$\boxed{G(\mathbf{r}, \mathbf{r}') = \frac{1}{2\pi} \ln |\mathbf{r} - \mathbf{r}'| + \text{const.}}$$

Note that in two dimensions we can no longer require $G \rightarrow 0$ at infinity, so the Dirichlet boundary conditions are usually specified in other ways.

2.3.3 The method of image for complicated geometry

We can seek Green's function over a infinite domain using the fundamental solution. However, we often want to find Green's functions in some more complicated domains, like a half-plane or a circle. This calls for the method of image which relies on uniqueness of solution to Poisson's equation. If we can cleverly introduce point sources **outside** the domain V such that the fundamental solution over the whole space also *happens* to satisfy the homogeneous boundary condition on the finite domain V , we can claim to find the **unique** solution.

For example, for a infinite plane/line boundary, we could simply reflect the source, as shown in Fig.1 to give the Green's function as a superposition of fundamental solutions:

$$G = -\frac{1}{4\pi} \left(\frac{1}{|\mathbf{r} - \mathbf{r}_0|} \pm \frac{1}{|\mathbf{r} - \mathbf{r}_1|} \right),$$

where \mathbf{r}_1 is the position of the image charge. With Dirichlet boundary conditions, an image source of opposite sign is introduced and we essentially have a dipole field with all flux through the wall (none to infinity). With Neumann boundary conditions, on the other hand, all the flux spreads out to infinity.

For a spherical or circular boundary with Dirichlet boundary condition, we could put an image charge at the inverse point $\mathbf{x}_1 = (a^2/|\mathbf{r}_0|)\mathbf{r}_0$ (so that a is the geometric mean of r_0 and r_1). For a 2D circle, the strength of the image charge is just $q = -Q$ while for a 3D sphere the image charge must have a weaker strength $q = -a/r_0$.

If the boundary is in more complicated geometry, some numerical methods are probably needed to find the Green's function. However, an inhomogeneous forcing or an inhomogeneous boundary condition with a simple geometry could be resolved by the following integral solution.




Figure 1: Image charges (red) for an infinite plane boundary conditions.




Figure 2: Inverse point for a spherical or circular Dirichlet boundary.

2.4 Integral solution to Poisson's equation

With the Green's function in hand, we could solve the inhomogeneous Poisson's equation $\nabla^2 \Phi = \rho(\mathbf{r})$ with inhomogeneous boundary condition $\Phi = f(\mathbf{r})$ on S or $\nabla \Phi \cdot \hat{\mathbf{n}} = f(\mathbf{r})$ on S . From the same vector calculus identity we used in proving uniqueness of Poisson's equation, we could prove Green's second identity ($\Psi = G$)

$$\int_V (\Phi \nabla^2 \Psi - \Psi \nabla^2 \Phi) dV = \oint_S \left(\Phi \frac{\partial \Psi}{\partial n} - \Psi \frac{\partial \Phi}{\partial n} \right) dS.$$

For a Dirichlet boundary condition, this simplifies to

$$\Phi(\mathbf{r}') = \int_V \rho(\mathbf{r}) G(\mathbf{r}, \mathbf{r}') dV + \oint_S f(\mathbf{r}) \nabla G \cdot d\mathbf{S},$$

where the first term comes from superposition of different sources and the second term comes from inhomogeneous boundary conditions. Similarly, the solution from Neumann boundary condition is

$$\Phi(\mathbf{r}') = \int_V \rho(\mathbf{r}) G(\mathbf{r}, \mathbf{r}') dV - \oint_S f(\mathbf{r}) G(\mathbf{r}, \mathbf{r}') dS + C.$$