

Wednesday, November 02, 2016
11:34 AM

Operations and Service Delivery sub-committee

1.	Maureen Kimani- Chair	NASCOP	Maureen.nyambura@gmail.com
2.	Patricia Oluoch- Co-chair	CDC	Hpk7@cdc.gov
3.	Tom Mawro		
4.	Brian Chirombo	WHO	pazvakavambwab@who.int
5.	Olivia Njathi	CHAI	onjathi@clintonhealthaccess.org
6.	Anangwe Samson	HJFMRI	Samson.anangwe@usamru-k.org
7.	Jason Reed	JHPIEGO	Jason.reed@jhpiego.org
8.	Elizabeth Irungu	Partners Scale Up	eirungu@pipsthika.org
9.	Mohamud Mohamed	NASCOP	Mudmoha09@yahoo.com
10.	Anthony Kamau	PSK	Kamau.anthony@ku.ac.ke
11.	Stephen Wanjala	MSF FRANCE	Msff-nairobi-medco-deputy@paris.msf.org
12.	Esther Karimi	PSK	ceo@psk.or.ke
13.	Dorcas Abuya	NHRL	abuyadorcas@gmail.com
14.	Patricia Muthea	NASCOP	patuerest@gmail.com
15.	Linda Misiko	NASCOP	tindilinda@yahoo.com
16.	Elizabeth Kubo	UON	En_kubo@yahoo.com
17.	Timothy Nzomo	NHRL	Jaytim2010@gmail.com
18.	Patricia Macharia	NASCOP	
19.	Lenet Bundi	NASCOP	bundilenet@gmail.com
20.	Barbara Mambo	NASCOP	bmambo@nascop.or.ke

TORs

Objective: Develop an operation tool kit (operational guidelines, training tools) by January 2018

Timeline: by 16th November

We need to pool together existing materials (Consultants also need to review what other material is available on the web- WHO/CDC)

Who will do the initial risk assessment?

Risk assessment: the revised guidelines already have a risk assessment tool that can be adapted to include what other implementing partners have developed. It Needs to be written in the form of a short tool that can be used at service delivery level to determine client eligibility. Past experience has shown there is need to be careful about the questions posed in the screening tool- short, precise and sensitive questions

The risk assessment tools will be availed to all HTS providers and lay counsellors, all clinicians, other entry points, pharmacies with HTS including where HIV self testing is provided.

Where will the risk assessment be done?

FP, STI, ANCs, DICES, Community prevention centres (previous VCT centres) private providers, pharmacies, community and facility testing points, youth friendly centres

Who will initiate PrEP?

PrEP will need to be initiated in places where there are backbone support structures e.g. laboratory

Pharmacies- pharmacists, pharmacy technicians can provide for re-fills but will need to be able to do HIV testing/ provide HIV self test kits

How long will PrEP be provided for?

This assumes HIV negative status is confirmed (for both initiation and re-fills).

PrEP should be given until the risk assessment determines there is 'zero risk' and no adverse reaction ascertained by relevant laboratory tests as provided in the guidelines; or when the client wishes to stop

PrEP should be given on the same day as enrollment, then 1 month and then 2 months after that (3 months post initiation day). Each visit will need to have HIV negative status confirmed

How will mobile KP be tracked? potential use of encrypted unique IDs/biometric; use of green cross pharmacy system countrywide