

Lawvere-Tierney Sheafification in Homotopy Type Theory

By

Kevin Quirin

Major Subject: Computer Science

Approved by the
Examining Committee:

Pierre Cointe, Thesis Advisor

Nicolas Tabareau, Thesis Advisor

Carl F. Gauß, Göttingen

Euclid, Athens

Mines de Nantes

2016

Abstract

The main goal of this thesis is to define an extension of Gödel not-not translation to all truncated types, in the setting of homotopy type theory. This goal will be achieved using the Lawvere-Tierney sheaves theory, by extending and formalizing the Lawvere-Tierney sheafification functor.

The definition of this functor needs concepts that were not defined in homotopy type theory, such as colimits over graphs and truncated modalities.

Almost all results presented in this thesis are formalized with the proof assistant Coq.

Write a better abstract

Acknowledgments

Write the acknowledgments

Contents

Acknowledgments	iii
Contents	iv
1 Introduction	1
2 Homotopy type theory	3
2.1 Dependent type theory	3
2.2 Identity types and Univalence axiom	8
2.3 Higher Inductives Types	11
2.4 Introduction to homotopy type theory	14
3 Higher modalities	21
3.1 Modalities	21
3.2 Examples of modalities	25
3.3 New type theories	27
3.4 Truncated modalities	27
3.5 Formalization	28
3.6 Translation	28
4 Colimits	31
4.1 Colimits over graphs	31
4.2 Van Doorn’s and Boulier’s constructions	37
4.3 Formalization	41
4.4 Towards groupoid objects	41
5 Sheaves in homotopy type theory	43
5.1 Sheaves in topoi	43
5.2 Sheaves in homotopy type theory	45
5.3 Formalization	59
5.4 Forcing in type theory	61
6 Conclusion and future works	63
Nomenclature	67
Bibliography	69

1 Introduction

Homotopy type theory is a very new branch of mathematics and computer science, exhibiting a strong, but surprising link between the theory of ω -categories and type theory. This topic hence live at the borderline between pure mathematics and computer science. One of the goals of researches on this topic is to use homotopy type theory as a new foundation for mathematics, replacing for example Zermelo-Fr enkel set theory. Its strong links with type theory would allow mathematicians to formalize their work with a proof assistant such as Coq [The12], Agda[Nor07] or Lean [Mou+15]. Indeed, errors in mathematical research papers seem to be inevitable [Voe14], and a computer-checked proof might be more trustable than a human-checked proof. One of the main difference between ZFC and HoTT foundations is that ZFC is defined in two steps (we need to define the logic, then the actual theory) while HoTT defines both at the same time.

The main ingredient in homotopy type theory, linking mathematics and computer science, is the Curry-Howard isomorphism: one can speak equivalently about proofs or about programs, they describe the same objects *via* a correspondance. For example, in type theory, the sequence of symbols $A \rightarrow B$ can be seen as the type of programs taking an argument of type A and producing an output of type B as well as the types of proofs that A implies B . Something implied by this correspondance is that there might exist different proofs of “ A implies B ”, since there are probably several ways to construct an output of type B from an input of type A . This property is called *proof-relevance*, while ZFC is considered as *proof-irrelevant*: if a lemma has been proved, the way it was proved can be forgotten, as it does not matter at all.

Go on!

Aims of the thesis The main goal of this thesis is to give a definition of a Lawvere-Tierney sheafification functor in the setting of homotopy type theory. In order to do this, we need in a first time to develop a theory of colimits in homotopy type theory, leading to thoughts on the definition of equivalence relation in type theory. Then, as our definition of sheafification is done inductively on the truncation levels, we need to define a truncated version of the just defined theory of colimits, as well as a truncated version of left-exact modalities. All these developments have been computer-checked by the proof assistant Coq, most of these available on my Github account <https://github.com/KevinQuirin>.


 Check plugin

Our deep study of modalities also lead us to define the translation of type theories associated to a (left-exact) modality, and write a Coq plugin to handle automatically that translation.

Plan of the thesis Let us describe the contents of this thesis. Chapter 2 recall the basic definitions in homotopy type theory. It is mainly based on [HoTT]¹, and serves more as a way for us to introduce notations to be consistent in the whole thesis. If the reader is not supposed to know anything about homotopy type theory before reading this thesis, this short introduction might be not enough to understand this setting, and he's strongly encouraged to take a look at [HoTT] before.

Chapter 3 introduces in its first part the theory of modalities as explained in [HoTT]. Then, we describe what we will call a *truncated modality*, which is a restricted version of modalities. Finally, we exhibit the translation of type theories induced by a left-exact modality.

In chapter 4, we describe a basic theory of colimits over graphs, and discuss an extension defining colimits over “graph with compositions”. This chapter, in a large part, has been formalized by Simon Boulrier in a library available at <https://github.com/SimonBoulrier/hott-colimits>. At the end of this chapter, we share our thoughts about groupoid objects, or equivalence relations in homotopy type theory. This section can easily be skipped by the reader, as it does not contain anything useful for the rest of the thesis.

The central (and last) chapter of this thesis is chapter 5. It describes our construction of the Lawvere-Tierney sheafification functor, which is a way to extend the not-not Gödel translation, valid only on h-propositions, to all truncated types. This result is the main contribution of the thesis. This chapter uses almost all the theory defined in previous chapters, and thus can hardly be read on its own. This section as been (almost) fully formalized, in a library available at <https://github.com/KevinQuirin/sheafification>.

¹Note that, to ease the reading, some references are shortcutted by their usual names: [HoTT] is the Homotopy Type Theory book, [HoTT/Coq] is the Coq/HoTT library, [HTT] is Lurie's monograph Higher topos theory, *etc.*

Homotopy type theory

Mathematics is the art of giving
the same name to different
objects.

Henri Poincaré

This chapter is an overview of the setting we will use in the rest of the thesis, homotopy type theory. The first part describes the formal system *dependent type theory*, which is a formal basis for homotopy type theory. This description is not intended to form a complete introduction to type theory accessible by anyone, but rather to clarify the formal system we use. A neophyte willing to read this thesis should read first a complete introduction to type theory, [Hof97] for an extensive one, but [HoTT] might be more accessible, and sufficient for the thesis. As we will see, we can view homotopy type theory as in

$$\text{HoTT} = \text{MLTT} + \text{UA} + \text{HIT}.$$

UA stands for *univalence axiom*, and is introduced in the second part of this chapter. HIT is the usual abbreviation for *higher inductive types*, which is a way, extensively discussed in [HoTT], to build new type, by giving constructors for the type as well as for its identity types. General principles will not be discussed here ; we rather present usual examples of HIT in section 2.3.

Finally, the last section of this chapter presents our point of view on identity paths, leading to the terminology *homotopy* type theory. This section also introduces the notion of *truncation level*, which plays a central role in the thesis.

2.1 Dependent type theory

In Zermelo-Frankel set theory, the most basic assertion is

$$x \in E$$

where x and E are sets. In dependent type theory, a similar judgement can be

$$a : A,$$

to be read as “ a is of type A ”. The main difference with membership relation is that an element a has one and only one type, while we can say $x \in E$ and $x \in F$ for the same element x in set theory (it is the definition of $x \in E \cap F$).

Dependent (or Martin-L f, due to its inventor [Mar98]) type theory is based on the Curry-Howard correspondance [How80], or propositions-as-types principle. Indeed, we do not need to make a difference between types and propositions. Hence $a : A$ will be read “ a is of type A ” when A is seen as a type, and “ a is a proof of A ” when A is seen as a proposition. In the rest of this section 2.1, we present the different types used to build dependent type theory. It is not intended to be complete ; for example we sometimes only give the non-dependent elimination rules to ease the reading. The reader can refer to [NPS01] or [HoTT] for a more detailed introduction.

2.1.1 Universes

In dependent type theory, types are also terms of a universe Type . Of course, we want the universe Type to be itself a type, and the Russel’s paradox¹ is close here. We solve the problem by using a cumulative hierarchy of universes

$$\text{Type}^0 : \text{Type}^1 : \text{Type}^2 : \dots$$

where every universe Type^i is of type Type^{i+1} . Cumulativity means that if $A : \text{Type}^k$ and $k < m$, then $A : \text{Type}^m$. The handling of universes level can be done automatically² by proof assistants such as Coq, and we will thus use only a “type of type” Type , to be understood in a polymorphic way.

2.1.2 Empty and Unit types

The first two types we will see are the Empty type (denoted $\mathbf{0}$) and the Unit type (denoted $\mathbf{1}$). These are respectively the types with zero and one elements (named \star). Those two types are dual to each other:

- having a term of type $\mathbf{0}$ in the context allows to prove anything, while having a term of type $\mathbf{1}$ in the context is useless
- dually, giving a term of type $\mathbf{1}$ is trivial, while giving a term of type $\mathbf{0}$ is impossible (if the theory is consistent).

Here are the (non-trivial) introduction and elimination rules for these types:

$$\frac{\Gamma \vdash x : \mathbf{0} \quad X : \text{Type}}{\Gamma \vdash X} \mathbf{0}\text{-ELIM} \qquad \frac{}{\Gamma \vdash \star : \mathbf{1}} \mathbf{1}\text{-INTRO}$$

Under the propositions-as-types principle, $\mathbf{0}$ is the type always false, and $\mathbf{1}$ the type always true. With a categorical point of view, $\mathbf{0}$ is an initial object and $\mathbf{1}$ is a terminal object.

¹In type theory, the argument is harder to prove, and is called Girard’s paradox.

²We will see in chapters 3 and 5 that we sometimes need to “help” Coq

2.1.3 Coproduct

The coproduct of A and B , noted $A + B$, is seen as the disjoint sum of A and B . It is described by the inductive type generated by

$$\left| \begin{array}{ll} \text{inl} & : A \rightarrow A + B \\ \text{inr} & : B \rightarrow A + B \end{array} \right.$$

The introduction and elimination rules of coproduct are:

$$\begin{array}{c} \frac{\Gamma \vdash a : A}{\Gamma \vdash \text{inl } a : A + B} \text{+INTRO}_L \quad \frac{\Gamma \vdash b : B}{\Gamma \vdash \text{inr } b : A + B} \text{+INTRO}_R \\[10pt] \frac{\Gamma \vdash p : A + B \quad \Gamma, x : A \vdash c_A : C \quad \Gamma, x : B \vdash c_B : C}{\Gamma \vdash \text{sum_rect}(p, c_A, c_B) : C} \text{+ELIM} \end{array}$$

Under the propositions-as-types principle, $A + B$ is seen as the disjunction of A and B .

2.1.4 Dependent product

One of the things both mathematicians and computer scientists love to do is to define functions. If A and B are types, one can consider the type of functions from A to B , taking an inhabitant of type A (called the source type) and giving an inhabitant of type B (the target type). What is new in dependent type theory is that the target type is allowed to depend on the argument of the function.

EXAMPLE – For example, one can consider a function taking a natural number n and giving a natural number greater than n . The target source depends indeed of n .

The type of dependent functions with source type A and target type Bx is called “dependent product over B ” (or “pi-type over B ”), and will be denoted $\prod_{x:A} Bx$ or $(x : A) \rightarrow (Bx)$. When B does not depend on A , we just say “arrow type” and note it $A \rightarrow B$.

The introduction and elimination rules of dependent products are:

$$\frac{\Gamma, x : A \vdash b : B}{\Gamma \vdash \lambda (x : A), b : \prod_{x:A} B} \prod\text{-INTRO} \quad \frac{\Gamma \vdash f : \prod_{x:A} B \quad \Gamma \vdash a : A}{\Gamma \vdash f(a) : B[a/x]} \prod\text{-ELIM}$$

Under the propositions-as-types, type of non-dependent functions $A \rightarrow B$ is seen as implication $A \Rightarrow B$, and type of dependent functions $\prod_{x:A} Bx$ is seen as universally quantified formulas $\forall x, Bx$.

2.1.5 Dependent sum

If A and B are two types, we would like to define the type of pairs (a, b) , where $a : A$ and $b : B$. The resulting type is called the product of A and B , noted $A \times B$.

As for functions, dependent type theory allows the second type to depend on the first type. Thus, the type of pairs where the first element x is in type A

and the second element y is in type Bx is called “dependent sum over B ” (or sigma-type over B), noted $\sum_{a:A} Ba$.

The introduction and elimination rules are:

$$\frac{\Gamma, x : A \vdash B : \text{Type} \quad \Gamma \vdash x : A \quad \Gamma \vdash b : B[a/x]}{\Gamma \vdash (a, b) : \sum_{x:A} B} \Sigma\text{-INTRO}$$

$$\frac{\Gamma \vdash p : \sum_{x:A} B}{\Gamma \vdash \pi_1 p : A} \Sigma\text{-INTRO}_1 \quad \frac{\Gamma \vdash p : \sum_{x:A} B}{\Gamma \vdash \pi_2 p : B[\pi_1 p/x]} \Sigma\text{-INTRO}_2$$

The projections π_1 and π_2 will sometimes be noted, when applied to a term u , u_1 and u_2 .

NOTE – We note that the terminology might be confusing: the dependent generalization of products are dependent sums, while dependent products are generalization of functions.

Under the propositions-as-types principle, $A \times B$ is seen as the conjunction $A \wedge B$, and $\sum_{x:A} Ba$ is seen as the existentially quantified formula $\exists x Bx$.

If A and B are types, $f : A \rightarrow B$ and $b : B$, the particular dependent sum $\sum_{a:A} fa = b$ is called the fiber of f over b , noted $\text{fib}_f(b)$. It describes the inverse image of b by f .

2.1.6 Inductive types

Dependent type theory actually allows us to define any inductive type. An inductive type is a type defined only by its introduction rules, in a free way. Its elimination rules are automatically determined. We have already seen examples of inductive types: Empty, Unit, Coproduct.

The most basic example of inductive types might be the type \mathbb{N} of naturals. Its introduction rules are

$$\frac{}{0 : \mathbb{N}} \mathbb{N}\text{-INTRO}_0 \quad \frac{\Gamma \vdash n : \mathbb{N}}{\Gamma \vdash S n : \mathbb{N}} \mathbb{N}\text{-INTRO}_S$$

We also say that its *constructors* are $0 : \mathbb{N}$ and $S : \mathbb{N} \rightarrow \mathbb{N}$: \mathbb{N} is thus the free monoid generated by 0 and S . The elimination rule for \mathbb{N} is the famous induction principle

$$\frac{\Gamma, x : \mathbb{N} \vdash P : \text{Type} \quad \Gamma \vdash c_0 : C[0/x] \quad \Gamma \vdash n : \mathbb{N} \quad \Gamma, n : \mathbb{N}, y : C \vdash c_S : C[S n/x]}{\Gamma \vdash \text{nat_ind}(C, c_0, c_S(n, y), n) : C[n/x]} \mathbb{N}\text{-ELIM}$$

This elimination rule allows us to define basic operators on natural numbers: addition, multiplication, order, *etc.*

For a reader interested by the theory of general inductive types, we refer him to [AGS12].

2.1.7 Paths type

One of the most powerful tool in dependent type theory might be the identity types. They allow us to talk about propositional equality between inhabitants of a type. The identity type over A will be noted $a =_A b$ or $a = b$ if A can be inferred from context. What is great about identity type over A is that its definition does not depend on the type A . If $a, b : A$, $a =_A b$ is defined as the inductive type whose only constructor is

$$\text{idpath} : \prod_{a:A} a =_A a$$

(idpath_x will sometimes be just noted 1_x , or even 1).

$$\frac{\Gamma \vdash a : A}{\Gamma \vdash \text{idpath}_a : a =_A a} =\text{-INTRO}$$

$$\frac{\Gamma \vdash a, b : A \quad \Gamma, x : A, y : A, p : x =_A y \vdash P : \text{Type} \quad \Gamma \vdash q : a =_A b \quad \Gamma, z : A \vdash w : P[z, z, 1/x, y, p]}{\Gamma \vdash \text{path_ind}_P^q(w) : P[a, b, q/x, y, p]} =\text{-ELIM}$$

The elimination principle should be looked closer. Lets simplify to understand how it works: suppose that the type family P depends only on two objects $x, y : A$. Then if C is reflexive, i.e if $C(x, x)$ is always inhabited, then whenever $x = y$, $C(x, y)$ is inhabited too.

Path-induction, also known as “J principle”, allow us to state the following:

Lemma 1 : [HoTT, Lemma 2.3.1]

Let $A : \text{Type}$ and $P : A \rightarrow \text{Type}$ a type family over A . Then if $p : x =_A y$, there is a function $\text{transport}_P^p : P x \rightarrow P y$.

Proof. Let’s do this proof to use path-induction for the first time. Let Q be the type family indexed by $x, y : A$

$$Q(x, y) := P x \rightarrow P y.$$

Then by path-induction, it suffices to define $Q(x, x)$. The latter is clearly inhabited by $\text{idmap}_{P x}$. \square

Path-induction might be the most powerful tool we can use. For example, most of the lemmas we will see in section 2.4 can be proved using path-induction.

2.1.8 Summary

We can summarize the situation in the following array:

Name	Notation	Proposition-as-types
Empty	$\mathbf{0}$	\perp
Unit	$\mathbf{1}$	\top
Coproduct, sum	$A + B$	$A \vee B$
Function	$A \times B$	$A \Rightarrow B$
Dependent function, pi-type	$\prod_{x:A} Bx$	$\forall x, Bx$
Product	$A \times B$	$A \wedge B$
Dependent sum, sigma-type	$\sum_{x:A} Bx$	$\exists x, Bx$

2.2 Identity types and Univalence axiom

Identity types are very useful to assert propositional equalities between object. Note that it does not characterize *judgemental* equality, which we consider here as belonging to the meta-theory (some theories, as Voevodsky’s Homotopy Type System [HTS], implemented in proof assistant Andromeda [m31]). One can prove that identity types give each type a structure of a ω -groupoid, *i.e* the structure of a ω -category where all arrows are invertible. For example, identity types over a type A satisfies:

- Reflexivity: for all $x : A$,

$$1_x : x = x$$

- Transitivity: for all $x, y, z : A$, $p : x = y$ and $q : y = z$,

$$p \cdot q : x = z$$

- Symmetry: for all $x, y : A$ and $p : x = y$, $p^{-1} : y = x$

- Reflexivity and symmetry behave well together: for all $x, y : A$ and $p : x = y$,

$$p \cdot p^{-1} = 1_x \text{ and } p^{-1} \cdot p = 1_y$$

Note that these paths are respectively in types $x = x$ and $y = y$.

- Reflexivity and associativity behave well together: for all $x, y : A$ and $p : x = y$,

$$p \cdot 1 = p \text{ and } 1 \cdot p = p$$

- Associativity: for all $w, x, y, z : A$ and $p : w = x, q : x = y, r : y = z$,

$$p \cdot (q \cdot r) = (p \cdot q) \cdot r$$

- MacLane pentagon: for all $v, w, x, y, z : A, p : v = w, q : w = x, r : x = y$ and $s : y = z$, the following diagram involving associativities commutes

$$\begin{array}{ccc}
 & p \cdot (q \cdot (r \cdot s)) & \\
 \swarrow & & \searrow \\
 (p \cdot q) \cdot (r \cdot s) & & p \cdot ((q \cdot r) \cdot s) \\
 \searrow & & \swarrow \\
 ((p \cdot q) \cdot r) \cdot s & \longrightarrow & (p \cdot (q \cdot r)) \cdot s
 \end{array}$$

- The coherences goes on and on.

A thing one can notice here is that we work in a proof-relevant setting. The types $x = y$ are just types, and thus can be inhabited by different objects. It might sound unfamiliar for mathematicians, but we will give in section 2.4 an interpretation of identity types justifying proof-relevance.

Identity types also have a good behavior with respect to function. Let $A, B : \text{Type}$ and $f : A \rightarrow B$; then

- For all $x, y : A$, there is a map $\text{ap}_f : x = y \rightarrow f(x) = f(y)$
- It is coherent with inverse: $\text{ap}_f p^{-1} = (\text{ap}_f p)^{-1}$
- It actually is coherent with the whole ω -groupoid structure. We refer to [HoTT] for more detailed results.

The fact that the definition of identity types does not depend on the type might seem strange, but it actually allows to catch the equality one would want ; for example, we can characterize the equalities between dependent pairs:

Lemma 2 : Paths in dependent sums

Let $A : \text{Type}$ and $B : A \rightarrow \text{Type}$. Then, for any $u, v : \sum_{x:A} Bx$, we have a term

$$\text{path}_\Sigma : \sum_{p:u_1=A} \text{transport}_p^B u_2 = v_2 \rightarrow u = v.$$

This characterization goes even further, as we can prove that path_Σ is an equivalence. Equivalences are very important objects in homotopy type theory, and are defined as

Definition 3 : Equivalence

Let $A, B : \text{Type}$ and $f : A \rightarrow B$. We say that f is an equivalence, noted $\text{IsEquiv}(f)$, if there are:

- a map $g : B \rightarrow A$, called the inverse of f
- a term $\text{retr}_f : \prod_{x:A} g(f(x)) = x$, called the retraction of the equivalence
- a term $\text{sect}_f : \prod_{x:A} f(g(x)) = x$, called the section of the equivalence
- a term $\text{adj}_f : \prod_{x:A} \text{ap}_f \text{retr}_f x = \text{sect}_f(f x)$, called the adjunction of the equivalence

We say that A and B are equivalent, noted $A \simeq B$, if there is a function $f : A \rightarrow B$ which is an equivalence.

Hence, path_Σ allows us to prove equalities in a dependent sum $\sum_{x:A} B a$, but its inverse allows us to prove, from an equality in the dependent sum, equalities in A and in $B a$.

Unfortunately, identity types does not allow to characterize equalities for all types. One example is paths in dependent products. In usual mathematics, one would like pointwise equality to be a sufficient (actually, a necessary and sufficient) property to state equality between function. In type theory, that would be phrased

$$\prod_{x:A} f x = g x \rightarrow f = g. \quad (2.1)$$

But one can prove (see [Str93] for a semantical proof) that this property, called *functional extensionality* cannot be proved from rules of dependent type theory. All we can say is that the backward function of 2.1 can be defined, called *happly*:

$$\text{happly} : f = g \rightarrow \prod_{x:A} f x = g x.$$

Functional extensionality will thus be stated as the axiom:

Axiom 4 : Functional extensionality

For any $A : \text{Type}$ and $B : A \rightarrow \text{Type}$, the arrow *happly* is an equivalence.

In a way, that could solve our problem. But dependent products are not the only types for which we cannot characterize identity types; the same problem arises with identity types of the universe. An answer was proposed first by Martin Hofmann and Thomas Streicher in 1996 in [HS96], and later (around 2005) by Vladimir Voevodsky: the univalence axiom. As for functional extensionality, it asserts that a certain arrow is an equivalence.

Axiom 5 : Univalence axiom

For any types $A, B : \text{Type}$, the function

$$\text{idtoequiv} : A = B \rightarrow A \simeq B$$

is an equivalence.

What is great about univalence axiom is that it seems to imply all other characterizations of identity types. For example:

Lemma 6 : [BL11][Lic]

Univalence axiom implies functional extensionality.

Another example the reader can think of is the type of streams (and more generally coinductive) extensionality: we want two streams (infinite list) to be equal when they are pointwise equal. It seems that univalence axiom implies it [Lic].

The main issue with univalence is that it clearly obliges us to work in a proof-relevant setting ; indeed, there are for examples two distinct proofs of $1 + 1 = 1 + 1$: the identity, and the function swapping left and right.

2.3 Higher Inductives Types

As seen in section 2.1.6, one can define a type by giving only its constructors ; but we also saw in section 2.2 that our setting is proof-relevant. It means that a type is characterized not only by its objects, but also by its identity types between these objects, and identity types of these identity types, *etc.* Hence, we would like to be able, like for inductive types, to define a type by giving constructors for the objects, and constructors for the identity types, *etc.* Then, the constructed type is the type freely generated by the constructors, and whose identity types are freely generated by the constructors, *etc.* Unlike inductive types, it is not known if elimination rules of higher inductive type can easily (and automatically) be inferred from the constructors. At the moment, we prefer to express them explicitly. As a general theory of higher inductive types would be very hard, we will only give some fundamental examples.

2.3.1 The circle

The most basic example of higher inductive type is the circle S^1 . The circle consists of just a point, with a non-trivial path above this point. It is defined as the higher inductive type generated by

$$\begin{array}{lcl} \text{base} & : & S^1 \\ \text{loop} & : & \text{base} = \text{base} \end{array}$$

It can be pictured as



As said, we give the elimination principle of the circle. We start by the non-dependent eliminator.

Lemma 7 : Non-dependent elimination of S^1

Let P be a type. If $b : P$ and $p : b = b$, then there is a map

$$S^1_{\text{rec}} : S^1 \rightarrow P$$

such that $S^1_{\text{rec}}(\text{base}) \equiv b$ (judgementally) and $\text{ap}_{S^1_{\text{rec}}}(\text{loop}) = p$ (propositionally).

It says that if in P you can find a “copy” of S^1 , then there is a “good” function $S^1 \rightarrow P$. The dependent eliminator is a bit more complicated, but still understandable.

Lemma 8 : Dependent elimination of S^1

Let $P : S^1 \rightarrow \text{Type}$ be a type family over S^1 . If $b : P(\text{base})$ and $p : \text{transport}_p^{\text{loop}}(b) = b$, then there is a term

$$S^1_{\text{ind}} : \prod_{x:S^1} P x$$

such that $S^1_{\text{ind}}(\text{base}) \equiv b$ and $\text{ap}_{S^1_{\text{ind}}}(\text{loop}) = p$.

The most famous result about the circle is the computation of its homotopy group $\pi_1(S^1)$.

Proposition 9 : Fundamental group of S^1 [LS13]

There is an equivalence

$$(\text{base} = \text{base}) \simeq \mathbb{Z}$$

where $\mathbb{Z} \stackrel{\text{def}}{=} \mathbb{N} + \mathbb{N} + 1$.

2.3.2 Coequalizers

Another example of higher inductive type is the computation of coequalizers. In category theory, a coequalizer of two objects a and b with arrow $f, g \in \text{Hom}(a, b)$ is a $c \in \text{Obj}$ together with an arrow $q \in \text{Hom}(b, c)$ such that

$a \begin{smallmatrix} \xrightarrow{f} \\ \xrightarrow{g} \end{smallmatrix} b \xrightarrow{q} c$ commutes, and which is universal with respect to this property, in the sense that for any other object $c' \in \text{Obj}$ with $q' \in \text{Hom}(b, c')$, there is an unique arrow $d : \text{Hom}(c, c')$. It might be seen as the diagram

$$\begin{array}{ccccc} a & \begin{smallmatrix} \xrightarrow{f} \\ \xrightarrow{g} \end{smallmatrix} & b & \xrightarrow{q} & c \\ & & & \searrow & \vdots \\ & & & q' & c' \end{array}$$

In homotopy type theory, the coequalizer of two functions $f, g : A \rightarrow B$ is defined as the higher inductive type $\text{Coeq}^{f,g}$ generated by

$$\left| \begin{array}{l} q : B \rightarrow \text{Coeq}^{f,g} \\ \alpha : \prod_{x:A} q \circ f(x) = q \circ g(x) \end{array} \right.$$

Its elimination principles are:

Lemma 10 : Elimination of coequalizer

Let A, B, f, g be as above.

- Let $P : \text{Type}$. If there are terms $q' : B \rightarrow P$ and $\alpha' : \prod_{x:A} q' \circ f(x) = q' \circ g(x)$, then there is a map

$$\text{Coeq}_{\text{rec}}^{f,g} : \text{Coeq}^{f,g} \rightarrow P$$

such that for all $b : B$, $\text{Coeq}_{\text{rec}}^{f,g}(qb) \equiv q'b$ and $\text{ap}_{\text{Coeq}_{\text{rec}}^{f,g}}(\alpha b) = \alpha' b$.

- Let $P : \text{Coeq}^{f,g} \rightarrow \text{Type}$. If there are terms $q' : \prod_{b:B} P(qb)$ and $\alpha' : \prod_{b:B} \text{transport}_P^{\alpha b}(q' \circ f(b)) = q' \circ g(b)$, then there is a dependent map

$$\text{Coeq}_{\text{ind}}^{f,g} : \prod_{x:\text{Coeq}^{f,g}} P x$$

such that for all $b : B$, $\text{Coeq}_{\text{ind}}^{f,g}(qb) \equiv q'b$ and $\text{ap}_{\text{Coeq}_{\text{ind}}^{f,g}}(\alpha b) = \alpha' b$.

It satisfies the desired universal property in the following sense:

- If Q is a type and $\varphi : \text{Coeq}^{f,g} \rightarrow Q$, then one can define a map $\psi : B \rightarrow Q$ such that for all $b : B$, $q \circ f(b) = q \circ g(b)$.
- One can prove that the map $\phi \mapsto \psi$ is an equivalence: from any other type such that the diagram commutes, there is an unique arrow $\text{Coeq}^{f,g} \rightarrow Q$.

Some results about coequalizers in homotopy type theory can be found in the form of Coq code in the HoTT/Coq library [HoTT/Coq].

2.3.3 Suspension

There is a way to see any type A as the identity type of another type: the suspension of A , noted ΣA . It is the higher inductive type generated by

$$\left| \begin{array}{ll} N & : \Sigma A \\ S & : \Sigma A \\ \text{merid} & : A \rightarrow (N = S) \end{array} \right.$$

Its elimination principle are:

Lemma 11 : Elimination of suspension

Let A be a type.

- Let $P : \text{Type}$. If there are $n, s : B$ and $m : A \rightarrow (n = s)$, then there is a map

$$\Sigma_{\text{rec}}^A : \Sigma A \rightarrow B$$

such that $\Sigma_{\text{rec}}^A(N) \equiv n$, $\Sigma_{\text{rec}}^A(S) \equiv s$ and for all $a : A$, $\text{ap}_{\Sigma_{\text{rec}}^A}(\text{merid}(a)) = m(a)$

- Let $P : \Sigma A \rightarrow \text{Type}$. If there are $n : P(N)$, $s : P(S)$ and $m : \prod_{a:A} \text{transport}_P^{\text{merid}(a)} n = s$, then there is a dependent map

$$\Sigma_{\text{ind}}^A : \prod_{a:A} P a$$

such that $\Sigma_{\text{ind}}^A(S) \equiv s$, $\Sigma_{\text{ind}}^A(N) \equiv n$ and $\text{ap}_{\Sigma_{\text{ind}}^A}(\text{merid}(a)) = m(a)$.

The first thing we can notice is that $\Sigma(\mathbf{1} + \mathbf{1}) \simeq \mathbb{S}^1$ [HoTT, Lemma 6.5.1]. Actually, we can define the sequence of n -spheres inductively by

$$\begin{aligned} \mathbb{S}^0 & \stackrel{\text{def}}{=} \mathbf{1} + \mathbf{1} \\ \mathbb{S}^{n+1} & \stackrel{\text{def}}{=} \Sigma \mathbb{S}^n \end{aligned}$$

2.4 Introduction to homotopy type theory

It is now time to really introduce homotopy type theory: it is the sum of sections 2.1, 2.2 and 2.3. In other words, homotopy type theory is just dependent type theory, augmented with univalence axiom and higher inductive types. We promised in section 2.2 an interpretation of identity types. In homotopy type theory, we view types as topological spaces and inhabitants of types as points of these spaces. Then, if $x, y : A$, the identity type $x = y$ is viewed as the topological space of paths (or continuous maps $f : [0, 1] \rightarrow A$ such that $f(0) = x$ and $f(1) = y$) between points x and y in A . As said, $x = y$ is itself a topological space, thus we can iterate the analogy. A path $r : p = q$ where $p, q : x = y$ are themselves paths is called a *homotopy*; a path between paths




Figure 2.1: Paths: a and b are points, p_1 and p_2 paths, H_1 and H_2 homotopies.

between paths is called a *2-homotopy*, etc. The situation can be depicted as in figure 2.1. Indeed, that interpretation allows to explain all properties of identity types.

- The equality 1_x is just the constant path

$$1_x: \begin{array}{ccc} [0,1] & \longrightarrow & A \\ t & \longmapsto & x \end{array}$$

- The inverse \bullet^{-1} is the function changing a path f into

$$g: \begin{array}{ccc} [0,1] & \longrightarrow & A \\ t & \longmapsto & f(1-t) \end{array}$$

- The concatenation $\bullet \cdot \bullet$ is the function changing paths f and g into

$$h: \begin{array}{ccc} [0,1] & \longrightarrow & A \\ t & \longmapsto & \begin{cases} f(2t) & \text{if } t \in [0, 1/2] \\ g(2t-1) & \text{if } t \in [1/2, 1] \end{cases} \end{array}$$

- A path between paths f and g between points x and y is a continuous function

$$H: \begin{array}{ccc} [0,1] & \longrightarrow & A \\ (u,v) & \longmapsto & H(u,v) \end{array}$$

such that $H(t,0) = f(t)$, $H(t,1) = g(t)$, $H(0,s) = x$ and $H(1,s) = y$. A homotopy is then a continuous deformation of f into g , fixing the ending points x and y .

The table 2.1 summarize the three points of view (type theoretic, groupoidal, homotopy theoretic). In the rest of the thesis, we will use any of the following name (e.g we will talk about “points of a type”, “path between inhabitants”, etc.).

What homotopy theorists love to do is to compute fundamental groups (i.e the group of paths between two points) of different spaces ([WX10][Hut11]). We can then categorized spaces with the level at which their homotopy groups become trivial. Voevodsky has realized that this notion admits a compact inductive definition internal to type theory, given by

These references were added without even opening them. Maybe a check would be good

Type theory	Homotopy theory	ω -groupoid
Type	Topological space	ω -groupoid
Inhabitant	Point	Object
Equality	Path	Morphism
idpath_x	Constant path	Identity morphism
\bullet^{-1}	Inverse path	Inverse morphism
$\bullet \cdot \bullet$	Concatenation of paths	Composition of morphisms
Equality between equalities	Homotopy	2-morphism

Table 2.1: Three points of view about HoTT

Definition 12 : Truncated types

$\text{Is-}n\text{-type}$ is defined by induction on $n \geq -2$:

- $\text{Is-}(-2)\text{-type}(X)$ if X is a contractible type, i.e X is inhabited by $c : X$, and every other point in X is connected to c .
- $\text{Is-}(n+1)\text{-type}(X) \stackrel{\text{def}}{=} \prod_{x,y:X} \text{Is-}n\text{-type}(x = y)$.

Then, $\text{Type}_n \stackrel{\text{def}}{=} \sum_{X:\text{Type}} \text{Is-}n\text{-type}(X)$.

For the first values of n , there are different names: (-2) -truncated types are called *contractible types*, (-1) -truncated types are called *h-propositions*, 0 -truncated types are called *h-sets*. Following this, $\text{Is-}(-2)\text{-type}$ is just Contr , $\text{Is-}(-1)\text{-type}$ is just IsHProp and Type_{-1} is HProp , and $\text{Is-}0\text{-type}$ is just IsHSet and Type_0 is HSet . An explanation of this terminology might be helpful. Contractible types are types, inhabited by a center c , with paths between any point and c . A kind of magic thing about contractible types is the lemma

Lemma 13

If $A : \text{Type}$ is contractible, then for any $x, y : A$, the type $x = y$ is contractible.

Inductively, it means that paths types of any level over a contractible type is contractible. It can be seen as the fact that a contractible type contains only one point c , for which there is only one path $c = c$, etc. The canonical example of contractible type is $\mathbf{1}$, and actually, any contractible type is equivalent to $\mathbf{1}$. Then, *h-propositions* are types where any two points are connected by a path. The only difference with contractible types is that we allow the type not to be inhabited. *H-propositions* are then proof-irrelevant types, in the sense that under the *propositions-as-types* principle, any points x and y in an HProp are equal, with a unique equality, which is thus irrelevant.

Lemma 14 : Example of h-propositions

The following types are h-propositions:

- $0, 1$
- $\text{IsEquiv}(f)$ for any $A, B : \text{Type}$ and $f : A \rightarrow B$.
- $\text{Is-}n\text{-type}(A)$ for any type $A : \text{Type}$ and truncation index n .

Now, by definition, h-sets are types for which identity types are in HProp . The world of h-sets can thus be seen as the “usual mathematical” world, as it satisfies proof-irrelevance.

The n -truncated types are stable under a lot of operations, as:

Lemma 15

Let $n \geq -2$ be a truncation index. Then

- If $A : \text{Type}_n$ and $x, y : A$, then $\text{Is-}n\text{-type}(x = y)$.
- If $A : \text{Type}_n$ and $B : A \rightarrow \text{Type}_n$, then $\text{Is-}n\text{-type}(\sum_{x:A} Bx)$.
- If $A : \text{Type}$ and $B : A \rightarrow \text{Type}_n$, then $\text{Is-}n\text{-type}(\prod_{x:A} Bx)$. Note that the source type needs not to be truncated.
- If $X : \text{Type}_n$, $Y : \text{Type}$ and $X \simeq Y$, then $\text{Is-}n\text{-type}(Y)$.
- If $X : \text{Type}_n$, then $\text{Is-}(n+1)\text{-type}(X)$.
- We have $\text{Is-}(n+1)\text{-type}\text{Type}_n$.

NOTA – Some types are not n -truncated for any n [HoTT, Example 8.8.6] ; it is highly suspected for example that the sphere S^2 is one of these ∞ -truncated types (it is at least true in homotopy theory).

This stratification of types induces a stratification of functions. A function $f : A \rightarrow B$ is said to be n -truncated if all its homotopy fibers $\text{fib}_f(b)$ are n -truncated. If $B : \text{Type}$, a type A together with a n -truncated map $f : A \rightarrow B$ will be called a n -subobject of B .

As in [RS13], we can define a sequence of subobject classifiers, namely, Type_n classifies n -subobjects of a type B , in the sens that there is an equivalence

$$\Xi : \sum_{A:\text{Type}} \sum_{f:A \rightarrow B} \prod_{b:B} \text{Is-}n\text{-type} \text{fib}_f(b) \xrightarrow{\sim} (B \rightarrow \text{Type}_n)$$

such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{t_f} & \text{Type}_n^\bullet \\ f \downarrow & & \downarrow \pi_1 \\ B & \xrightarrow{\chi_f} & \text{Type}_n \end{array}$$

is a pullback for any f with n -truncated homotopy fibers where $\text{Type}_n^\bullet \stackrel{\text{def}}{=} \sum_{A:\text{Type}} A$ is the universe of pointed n -truncated types and

$$t_f = \lambda a, (\text{fib}_f(f(a)), (a, \text{idpath})).$$

We will use the equivalence Ξ to define n -subobjects of a type B either as a n -truncated map $A \rightarrow B$, either as a characteristic map $\chi : B \rightarrow \text{Type}_n$.

2.4.1 Truncations

We present here a way to change any type into a n -truncated type, using truncations. The interested reader can read Nicolai Kraus' PhD thesis [Kra15] consecrated to truncation levels in HoTT.

Let $n \geq -1$ be a truncation index. The n -truncation of a type A is the higher inductive type $\|A\|_n$ generated by

$$\left| \begin{array}{ll} \text{tr}_n & : A \rightarrow \|A\|_n \\ \alpha_{\text{tr}}^n & : \text{Is-}n\text{-type}(\|A\|_n) \end{array} \right|$$

If $a : A$, $\text{tr}_n(a)$ will be noted $|a|_n$. The elimination principles are:

Lemma 16 : Elimination principle of truncations

Let $A : \text{Type}$ and $n \geq -1$ be a truncation index.

- If $P : \text{Type}$ such that $\text{Is-}n\text{-type}(P)$ and $f : A \rightarrow P$, then there is a map

$$|f|_n : \|A\|_n \rightarrow P$$

such that for all $a : A$, $|f|_n(|a|_n) \equiv f(a)$.

- If $P : \|A\|_n \rightarrow \text{Type}$ such that $\prod_{x:\|A\|_n} \text{Is-}n\text{-type}(Px)$ and $f : \prod_{a:A} P(|a|_n)$, then there is a dependent map

$$|f|_n : \prod_{x:\|A\|_n} Px$$

such that for all $a : A$, $|f|_n(|a|_n) \equiv f(a)$.

Basically, this induction principle says that $\|A\|_n$ contains as much data about A than A itself, but it can only be used to define a n -truncated type. It can be expressed as the following universal property:

Lemma 17 : Universal property of truncations

Let $A : \text{Type}$ and $P : \text{Type}_n$. Then the map

$$\text{precompose}_{\text{tr}_n} : \begin{array}{ccc} A \rightarrow P & \longrightarrow & \|A\|_n \rightarrow P \\ f & \longmapsto & f \circ \text{tr}_n \end{array}$$

is an equivalence.

We will see in chapter 3 that all $(\text{Is-}n\text{-type}, \|\bullet\|_n)$ define *modalities*. In particular, every truncation index n yields a factorization system:

Proposition 18

Let $n \geq -1$ a truncation index. Then any map $f : A \rightarrow B$ can be factored as a n -connected map followed by a n -truncated map, where a map is n -connected if for all $b : B$, $\|\text{fib}_f(b)\|_n$ is contractible.

In the rest of this thesis, we will only use this result with $n = -1$. In that case, (-1) -truncated functions are called *embeddings*, and (-1) -connected functions are called *surjections*. The factorization of a map $f : A \rightarrow B$ is given by

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow \tilde{f} & \nearrow \pi_1 \\ & \text{Im}(f) & \end{array}$$

$$\begin{aligned} \text{Im}(f) &\stackrel{\text{def}}{=} \sum_{b:B} \left\| \sum_{a:A} f a = b \right\|_{-1} \\ \tilde{f} &\stackrel{\text{def}}{=} \lambda a : A, (f a, |(a, 1_b)|_{-1}) \end{aligned}$$

Higher modalities

As said in the introduction, the main purpose of our work is to build, from a model \mathfrak{M} of homotopy type theory, another model \mathfrak{M}' satisfying new principles. Of course, \mathfrak{M}' should be describable *inside* \mathfrak{M} . In set theory, it corresponds to building *inner models* ([Kun]). In type theory, it can be rephrased in terms of left-exact modalities: it consists of an operator \circ on types such that for any type A , $\circ A$ satisfies a desired property. If the operator has a “good” behaviour, then it is a modality, and the universe of all types satisfying the chosen property forms a new model of homotopy type theory.

3.1 Modalities

Definition 19

A left exact modality is the data of

- (i) A predicate $P : \text{Type} \rightarrow \text{HProp}$
- (ii) For every type A , a type $\circ A$ such that $P(\circ A)$
- (iii) For every type A , a map $\eta_A : A \rightarrow \circ A$

such that

- (iv) For every types A and B , if $P(B)$ then

$$\left\{ \begin{array}{ll} (\circ A \rightarrow B) & \rightarrow (A \rightarrow B) \\ f & \mapsto f \circ \eta_A \end{array} \right.$$

is an equivalence.

- (v) for any $A : \text{Type}$ and $B : A \rightarrow \text{Type}$ such that $P(A)$ and $\prod_{x:A} P(Bx)$, then $P(\sum_{x:A} B(x))$
- (vi) for any $A : \text{Type}$ and $x, y : A$, if $\circ A$ is contractible, then $\circ(x = y)$ is contractible.

Conditions (i) to (iv) define a reflective subuniverse, (i) to (v) a modality.

NOTATION – The inverse of $- \circ \eta_A$ from point (iv) will be denoted $\circ_{\text{rec}} : (A \rightarrow B) \rightarrow (\circ A \rightarrow B)$, and its computation rule $\circ_{\text{rec}}^\beta : \prod_{f:A \rightarrow B} \prod_{x:A} \circ_{\text{rec}}(f)(\eta_A x) = f x$.

If \circ is a modality, the type of modal types will be denoted Type° . Let us fix a left-exact modality \circ for the rest of this section. A modality acts functorially on Type , in the sense that

Lemma 20 : Functoriality of modalities

Let $A, B : \text{Type}$ and $f : A \rightarrow B$. Then there is a map $\circ f : \circ A \rightarrow \circ B$ such that

- $\circ f \circ \eta_A = \eta_B \circ f$
- if $g : B \rightarrow C$, $\circ(g \circ f) = \circ g \circ \circ f$
- if $\text{IsEquiv } f$, then $\text{IsEquiv } \circ f$.

Proposition 21

Any left-exact modality \circ satisfies the following properties¹.

- (R) A is modal if and only if η_A is an equivalence.
- (R) $\mathbf{1}$ is modal.
- (R) Type° is closed under dependent products, i.e $\prod_{x:A} Bx$ is modal as soon as all Bx are modal.
- (R) For any types A and B , the map

$$\circ(A \times B) \rightarrow \circ A \times \circ B$$

is an equivalence.

- (R) If A is modal, then for all $x, y : A$, $(x = y)$ is modal.
- (M) For every type A and $B : \circ(A) \rightarrow \text{Type}^\circ$, then

$$-\circ \eta_A : \begin{array}{ccc} \prod_{z:\circ A} Bx & \longrightarrow & \prod_{a:A} B(\eta_A a) \\ f & \longmapsto & f \circ \eta_A \end{array}$$

is an equivalence.

- (M) If $A, B : \text{Type}$ are modal, then so are $\text{Is-}n\text{-type } A$, $A \simeq B$ and $\text{IsEquiv } f$ for all $f : A \rightarrow B$.
- (L) If $A : \text{Type}_n$, then $\circ A : \text{Type}_n$.
- (L) If $X, Y : \text{Type}$ and $f : X \rightarrow Y$, then the map

$$\circ(\text{fib}_f(y)) \rightarrow \text{fib}_{\circ f}(\eta_B y)$$

¹Properties needing only a reflective subuniverse are annotated by (R), a modality by (M), a left-exact modality by (L)

is an equivalence, and the following diagram commutes

$$\begin{array}{ccc} \text{fib}_f(y) & \xrightarrow{\eta} & \circ(\text{fib}_f(y)) \\ \gamma \downarrow & \swarrow & \\ \text{fib}_{\circ f}(\eta_B y) & & \end{array}$$

NOTATION – Again, the inverse of $- \circ \eta_A$ will be denoted $\circ_{\text{ind}} : \prod_{a:A} B(\eta_A a) \rightarrow \prod_{x:\circ A} Bx$, and its computation rule $\circ_{\text{ind}}^\beta : \prod_{f:\prod_{a:A} B(\eta_A a)} \prod_{x:A} \circ_{\text{ind}}(f)(\eta_A x) = fx$

Proof. We only prove the last point. It is straightforward to define a map

$$\phi : \sum_{x:X} fx = y \rightarrow \sum_{x:\circ X} \circ fx = \eta_Y y,$$

using η functions. We will use the following lemma to prove that the function induced by ϕ defines an equivalence:

Lemma 22

Let $X : \text{Type}$, $Y : \text{Type}^\circ$ and $f : X \rightarrow Y$. If for all $y : Y$, $\circ(\text{fib}_f(y))$ is contractible, then $\circ X \simeq Y$.

Hence we just need to check that every \circ -fiber $\circ(\text{fib}_\phi(x;p))$ is contractible. Technical transformations allow one to prove

$$\text{fib}_\phi(x;p) \simeq \text{fib}_s(y;p^{-1})$$

for

$$s : \begin{array}{ccc} \text{fib}_{\eta_X}(x) & \longrightarrow & \text{fib}_{\eta_Y}(\circ f x) \\ (a,q) & \longmapsto & (f a, -) \end{array}$$

But left-exactness allows to characterize the contractibility of fibers:

Lemma 23

Let $A, B : \text{Type}$. Let $f : A \rightarrow B$. If $\circ A$ and $\circ B$ are contractible, then so is $\text{fib}_f(b)$ for any $b : B$.

Thus, we just need to prove that $\circ(\text{fib}_{\eta_X}(a))$ and $\circ(\text{fib}_{\eta_Y}(b))$ are contractible. But one can check that η maps always satisfy this property. Finally, $\circ(\text{fib}_s(y;p^{-1}))$ is contractible, so $\circ(\text{fib}_\phi(x;p))$ also, and the result is proved. \square

Let us finish these properties by the following proposition, giving an equivalent characterization of left-exactness.

Proposition 24

Let \circ be a modality. Then \circ is left-exact if and only if \circ preserves path spaces, i.e

$$\prod_{A:\text{Type}} \prod_{x,y:A} \text{IsEquiv}(\circ(\text{ap}_{\eta_A}))$$

where $\circ(\text{ap}_{\eta_A}) : \circ(x = y) \rightarrow \eta_A x = \eta_A y$.

Proof. We will rather prove something slightly more general, using an encode-decode proof [HoTT, Section 8.9] ; we will characterize, for a type A and a fixed inhabitant $x : A$ the type

$$\eta_A x = y$$

for any $y : \circ A$.

Let $\text{Cover} : \circ A \rightarrow \text{Type}^\circ$ be defined by induction by

$$\text{Cover}(y) \stackrel{\text{def}}{=} \circ_{\text{rec}}(\lambda y, \circ(x = y)).$$

Note that for any $y : \circ A$, $\text{Cover}(y)$ is always modal. We will show that $\eta_A x = y \simeq \text{Cover}(y)$. Now, let $\text{Encode} : \prod_{y:\circ A} \eta_A x = y \rightarrow \text{Cover}(y)$ be defined by

$$\text{Encode}(y, p) \stackrel{\text{def}}{=} \text{transport}_{\text{Cover}}^p \left(\text{transport}_{\text{idmap}}^{\circ_{\text{rec}}^\beta((\lambda z, \circ(x=z)), x)} (\eta_{x=x} 1) \right)$$

and $\text{Decode} : \prod_{y:\circ A} \text{Cover}(y) \rightarrow \eta_A x = y$ by

$$\text{Decode} \stackrel{\text{def}}{=} \circ_{\text{ind}} \left(\lambda y p, \circ(\text{ap}_{\eta_A}) \left(\text{transport}_{\text{idmap}}^{\circ_{\text{rec}}^\beta((\lambda z, \circ(x=y)), y)} p \right) \right)$$

Then one can show, using \circ -induction and path-induction, that for any $y : \circ A$, $\text{Encode}(y, -)$ and $\text{Decode}(y, -)$ are each other inverses. Then, taking $y' = \eta_A y$, we have just shown that $\eta_A x = \eta_A y \simeq \text{Cover}(\eta_A y)$, which is itself equivalent, by \circ_{rec}^β , to $\circ(x = y)$. It is straightforward to check that the composition $\circ(x = y) \rightarrow \text{Cover}(\eta_A y) \rightarrow \eta_A x = \eta_A y$ is exactly $\circ(\text{ap}_{\eta_A})$.

Now, let us prove the backward implication. Let A be a type such that $\circ A$ is contractible, and $x, y : A$. As $\eta_A x, \eta_A y : \circ A$, we know that $\eta_A x = \eta_A y$ is contractible. But as $\eta_A x = \eta_A y \simeq \circ(x = y)$ by assumption, $\circ(x = y)$ is also contractible. \square

As this whole thesis deals with truncation levels, it should be interesting to see how they are changed under a modality. We already know that if a type T is (-2) -truncated, i.e contractible, then it is unchanged by the reflector:

$$\circ T \simeq \circ \mathbf{1} \simeq \mathbf{1} \simeq T.$$

Thus, Type_{-2} is closed by any reflective subuniverse. Now, let $T : \text{HProp}$. To check that $\circ T$ is an h-proposition, it suffices to check that

$$\prod_{x,y:\circ T} x = y$$

For any $x : \circ T$, the type $\prod_{y:\circ T} x = y$ is modal, as all $x = y$ are ; by the same argument, $\prod_{x:\circ T} x = y$ is modal too for any $y : \circ T$. Using twice the dependent eliminator of \circ , it now suffices to check that

$$\prod_{x,y:T} \eta_T x = \eta_T y.$$

As T is supposed to be an h-proposition, this is true. It suffices to state

Lemma 25

For any modality, Type_{-1} is closed under the reflector \circ , i.e

$$\prod_{P:\text{HProp}} \text{IsHProp}(\circ P).$$

A simple induction on the truncation level, together with the left-exactness property allows to state

Lemma 26

For any left-exact modality, all Type_p are closed under the reflector \circ , i.e

$$\prod_{P:\text{Type}_p} \text{Is-}p\text{-type}(\circ P).$$

We note that this is not a equivalent characterization of left-exactness, as it is satisfied by truncations, and we will see they are not left-exact.

3.2 Examples of modalities

3.2.1 The identity modality

Let us begin with the most simple modality one can imagine: the one doing nothing. We can define it by letting $\circ A \stackrel{\text{def}}{=} A$ for any type A , and $\eta_A \stackrel{\text{def}}{=} \text{idmap}$. Obviously, the desired computation rules are satisfied, so that the identity modality is indeed a left-exact modality.

It might sound useless to consider such a modality, but it can be precious when looking for properties of modalities: if it does not hold for the identity modality, it cannot hold for an abstract one.

3.2.2 Truncations

The first class of non-trivial examples might be the *truncations* modalities, seen in 2.4.1.

3.2.3 Double negation modality

Proposition 27

The double negation modality $\circ A \stackrel{\text{def}}{=} \neg\neg A$ is a modality.

Proof. We define the modality with

- (i) We will define the predicate P later.
- (ii) \circ is defined by $\circ A = \neg\neg A$
- (iii) We want a term η_A of type $A \rightarrow \neg\neg A$. Taking

$$\eta_A \stackrel{\text{def}}{=} \lambda x : A, \lambda y : \neg A, y a$$

do the job.

Now, we can define P to be exactly $\prod_{A:\text{Type}} \text{IsEquiv } \eta_A$.

- (iv) Let $A, B : \text{Type}$, and $\varphi : A \rightarrow \neg\neg B$. We want to extend it into $\psi : \neg\neg A \rightarrow \neg\neg B$. Let $a : \neg\neg A$ and $b : \neg B$. Then $a(\lambda x : A, \varphi x b) : \mathbf{0}$, as wanted. One can check that it forms an equivalence.
- (v) Let $A : \text{Type}$ and $B : A \rightarrow \text{Type}$ such that $P(A)$ and $\prod_{a:A} P(Ba)$. There is a map

$$\sum_{x:A} Bx \rightarrow A$$

thus by the previous point, we can extend it into

$$\kappa : \neg\neg \sum_{x:A} Bx \rightarrow A.$$

It remains to check that for any $x : \neg\neg \sum_{x:A} Bx$, $B(\kappa x)$.

But the previous map can be easily extended to the dependent case, and thus it suffices to show that for all $x : \sum_{x:A} Bx$, $B(\kappa(\eta x))$. As $\kappa \circ \eta = \text{idmap}$, the goal is solved by $\pi_2 x$.

□

Unfortunately, it appears that the only type which can be modal are h-propositions, as they are equivalent to their double negation which is always an h-proposition. Thus, the type of modal types is consisted only of h-proposition, which is not satisfactory. The main purpose of this thesis, in particular chapter 5 is to extend this modality into a better one.

3.3 New type theories

New type theory: Enhance me

We suppose here that \circ is a left-exact modality such that Type° is modal. This is for example the case when the modality is *accessible* (see [HoTT/Coq] for definition and proof).

Proposition 28

A left exact modality \circ induces a consistent type theory if and only if $\circ\mathbf{0}$ can not be inhabited in the initial type theory. In that case, the modality is said to be consistent.

Proof. By condition (iv) of Definition 19, $\circ\mathbf{0}$ is an initial object of Type° , and thus corresponds to false for modal mere proposition. As $\circ\mathbf{1} = \mathbf{1}$, Type° is consistent when $\circ\mathbf{0} \neq \mathbf{1}$, that is when there is no proof of $\circ\mathbf{0}$. \square

3.4 Truncated modalities

As for colimits, we define a truncated version of modalities, in order to use it in chapter 5. Basically, a truncated modality is the same as a modality, but restricted to Type_n .

Definition 29 : Truncated modality

Let $n \geq -1$ be a truncation index. A left exact modality at level n is the data of

- (i) A predicate $P : \text{Type}_n \rightarrow \text{HProp}$
- (ii) For every n -truncated type A , a n -truncated type $\circ A$ such that $P(\circ A)$
- (iii) For every n -truncated type A , a map $\eta_A : A \rightarrow \circ A$

such that

- (iv) For every n -truncated types A and B , if $P(B)$ then

$$\begin{cases} (\circ A \rightarrow B) & \rightarrow & (A \rightarrow B) \\ f & \mapsto & f \circ \eta_A \end{cases}$$

is an equivalence.

- (v) for any $A : \text{Type}_n$ and $B : A \rightarrow \text{Type}_n$ such that $P(A)$ and $\prod_{x:A} P(Bx)$, then $P(\sum_{x:A} B(x))$
- (vi) for any $A : \text{Type}_n$ and $x, y : A$, if $\circ A$ is contractible, then $\circ(x = y)$ is contractible.

Properties of truncated left-exact modalities described in 21 are still true when restricted to n -truncated types, except the one that does not make sense: Type_n° cannot be modal, as it is not even a n -truncated type.

3.5 Formalization

Let us discuss here about the formalization of the theory of modalities. General modalities are formalized in the Coq/HoTT library [HoTT/Coq], thanks to a huge work of Mike Shulman [Shu15]. The formalization might seem to be straightforward, but the universe levels (at least, their automatic handling by Coq) are here a great issue. Hence, we have to explicitly give the universe levels and their constraints in a large part of the library. For example, the reflector \circ of a modality is defined, in [HoTT/Coq] as

$$\circ : \text{Type}^i \rightarrow \text{Type}^i;$$

it maps any universe to itself.

In chapter 5, we will need a slightly more general definition of modality. The actual definitions stay the same, but the universes constraints we consider change. The reflector \circ will now have type

$$\circ : \text{Type}^i \rightarrow \text{Type}^j, \quad i \leq j;$$

it maps any universe to a possibly higher one. Other components of the definition of a modality are changed in the same way.

Fortunately, this change seems small enough for all properties of modalities to be kept. Of course, the examples of modalities mapping any universe to itself are still an example of generalized modality, it just does not use the possibility to inhabit a higher universe. This has been computer-checked, it can be found at https://github.com/KevinQuirin/HoTT/tree/extended_modalities.

We would like to have the same generalization for truncated modalities. But there are a lot of new universe levels popping out, mostly because in $\text{Type}_n = \sum_{T:\text{Type}} \text{Is-}n\text{-type } T$, $\text{Is-}n\text{-type}$ come with its own universes. Hence, handling “by hand” so many universes together with their constraints quickly go out of control. One idea to fix this issue could be to use *resizing rules* [Voe11], allowing h -propositions to live in the smallest universe. We could then get rid of the universes generated by $\text{Is-}n\text{-type}$, and treat the truncated modality exactly as generalized modalities.

3.6 Translation

As said in section 3.3, left-exact modalities allows to perform model transformation. But it can be enhanced a bit by exhibiting a *translation* of type theories, as it has been done for forcing [JTS12]. Let us explain here how this translation works.

Let \circ be an accessible left-exact modality. We describe, for each type constructor, how to build its translation. We denote $\pi_{\circ}(A)$ the proof that $\circ(A)$ is always modal.

- For types

$$[\text{Type}] \stackrel{\text{def}}{=} (\text{Type}^{\circ}, \pi_{\text{Type}^{\circ}})$$

where $\pi_{\text{Type}^{\circ}}$ is a proof that Type° is itself modal. To ease the reading in what follows, we introduce the notation

$$\llbracket A \rrbracket \stackrel{\text{def}}{=} \pi_1 [A]$$

- For dependent sums

$$\begin{aligned} [\sum_{x:A} B] &\stackrel{\text{def}}{=} (\sum_{x:\llbracket A \rrbracket} \llbracket B \rrbracket, \pi_{\Sigma}^{[A],[B]}) \\ [(x, y)] &\stackrel{\text{def}}{=} ([x], [y]) \\ [\pi_i t] &\stackrel{\text{def}}{=} \pi_i [t] \end{aligned}$$

where $\pi_{\Sigma}^{A,B}$ is a proof that $\sum_{x:A} B$ is modal when A and B are.

- For dependent products

$$\begin{aligned} [\prod_{x:A} B] &\stackrel{\text{def}}{=} (\prod_{x:\llbracket A \rrbracket} \llbracket B \rrbracket, \pi_{\Pi}^{[A],[B]}) \\ [\lambda x : A, M] &\stackrel{\text{def}}{=} \lambda x : \llbracket A \rrbracket, [M] \\ [t t'] &\stackrel{\text{def}}{=} [t][t'] \end{aligned}$$

where $\pi_{\Pi}^{A,B}$ is a proof that $\prod_{x:A} B$ is modal when B is.

- For paths

$$\begin{aligned} [A = B] &\stackrel{\text{def}}{=} ([A] = [B], \pi_{=}^{[A],[B]}) \\ [1] &\stackrel{\text{def}}{=} 1 \\ [J] &\stackrel{\text{def}}{=} J \end{aligned}$$

where $\pi_{=}^{A,B}$ is a proof that $A = B$ is modal when A and B are, if $A, B : \text{Type}$, or a proof that $A = B$ is modal as soon as their type is modal if $A, B : X$.

- For positive types (we only treat the case of the sum as an example)

$$\begin{aligned} [A + B] &\stackrel{\text{def}}{=} (\circ(\llbracket A \rrbracket + \llbracket B \rrbracket); \pi_{\circ}(\llbracket A \rrbracket + \llbracket B \rrbracket)) \\ [\text{in}_{\ell} t] &\stackrel{\text{def}}{=} \eta(\text{in}_{\ell} [t]) \\ [\text{in}_r t] &\stackrel{\text{def}}{=} \eta(\text{in}_r [t]) \\ [\langle f, g \rangle] &\stackrel{\text{def}}{=} \circ_{\text{rec}}^{\llbracket A \rrbracket + \llbracket B \rrbracket} \langle [f], [g] \rangle \end{aligned}$$

- For truncations ($i \leq n$)

$$\begin{aligned} \llbracket A \rrbracket_i &\stackrel{def}{=} (\circ \llbracket A \rrbracket_i; \pi_\circ(\llbracket A \rrbracket_i)) \\ \llbracket t \rrbracket_i &\stackrel{def}{=} \eta \llbracket t \rrbracket_i \\ \llbracket f \rrbracket_i &\stackrel{def}{=} \circ_{\text{rec}}^{\llbracket A \rrbracket_i} \llbracket f \rrbracket_i \end{aligned}$$

As in the forcing translation [JTS12], the main issue is that convertibility might not be preserved. For example, let $f : A \rightarrow X$ and $g : B \rightarrow X$. Then $\langle f, g \rangle(\text{in}_\ell t)$ is convertible to $f t$, but $\llbracket \langle f, g \rangle(\text{in}_\ell t) \rrbracket$ reduces to $\circ_{\text{rec}}^{\llbracket A \rrbracket + \llbracket B \rrbracket} \langle \llbracket f \rrbracket, \llbracket g \rrbracket \rangle(\eta(\text{in}_\ell t))$, which is only equal to $\llbracket f t \rrbracket$.

Two solutions to this problem can come to our mind:

- We can use the eliminator J of equality in the conversion rule, but this would require to use a type theory with explicit conversion in the syntax, like in [JTS12].
- We can ask the modality \circ to be a strict modality, in the sense that retraction in the equivalence of $(- \circ \eta)$ is conversion instead of equality, like for the closed modality. That way, we keep the conversion rule, *i.e.* if $\Gamma \vdash u \equiv v$, then $\llbracket \Gamma \rrbracket \vdash \llbracket u \rrbracket \equiv \llbracket v \rrbracket$. That is the solution we chose.

The usual result we want about any translation is its soundness:

Proposition 30 : Soundness of the translation

Let Γ is a valid context, A a type and t a term. If $\Gamma \vdash t : A$, then

$$\llbracket \Gamma \rrbracket \vdash \llbracket t \rrbracket : \llbracket A \rrbracket.$$

NOTA – We note that if the modality is not left-exact (or not accessible), like truncations modalities, then Type° is not itself modal. It is although still possible to write a translation, but we can only define it on a type theory with only one universe. Indeed, the judgement $\Gamma \vdash \text{Type}^i : \text{Type}^j$ cannot be expressed, and thus cannot be translated to $\llbracket \Gamma \rrbracket \vdash \llbracket \text{Type}^i \rrbracket : \llbracket \text{Type}^j \rrbracket$.

An implementation of this translation has been made, in the form of a Coq plugin, available at <https://github.com/KevinQuirin/translation-mod/>.

Code the plugin, then plug it here

4 Colimits

A comathematician is a device
turning cotheorems into ffee.

Co-Alfréd Rényi

As seen in chapter 2, adding sigma-types to type theory results in adding limits over graphs in the underlying category, and adding higher inductive types results in adding colimits over graphs. If limits has been extensively studied in [AKL15], theory of colimits was not completely treated.

The following is conjoint work with Simon Boulrier and Nicolas Tabareau, helped by precious discussions with Egbert Rijke. The sections 4.1 and 4.2 are extended version of the blog post [Bou16].

4.1 Colimits over graphs

As colimits are just dual to limits, it seems that it would be very easy to translate the work on limits to colimits. Although, even if it might be because we are more habituated to manipulate sigma-types than higher inductive types, it seems way harder.

4.1.1 Definitions

Let's recall the definitions of graphs and diagrams over graphs, introduced in [AKL15].

Definition 31 : Graph

A graph G is the data of

- a type G_0 of vertices ;
- for any $i, j : G_0$, a type $G_1(i, j)$ of edges.

Definition 32 : Diagram

A diagram D over a graph G is the data of

- for any $i : G_0$, a type $D_0(i)$;
- for any $i, j : G_0$ and all $\phi : G_1(i, j)$, a map $D_1(\phi) : D_0(i) \rightarrow D_0(j)$

When the context is clear, G_0 will be simply denoted G , $G_1(i, j)$ will be noted $G(i, j)$, $D_0(i)$ will be noted $D(i)$ or D_i , and $D_1(\phi)$ will be noted $D_{i,j}(\phi)$ or simply $D(\phi)$ (i and j can be inferred from ϕ).

EXAMPLES :

- One can consider the following graph, namely the graph of (co)equalizers

$$\bullet \rightrightarrows \bullet$$

Here, $G_0 = 2$, $G_1(\top, \perp) = 2$ and other $G_1(i, j)$ are empty.

A diagram over this graph consists of two types A and B , and two maps $f, g : A \rightarrow B$, producing the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \rightrightarrows \\ \xrightarrow{g} \end{array} B$$

- The graph of the mapping telescope is

$$\bullet \longrightarrow \bullet \longrightarrow \dots$$

In other words, $G_0 = \mathbb{N}$ and $G_1(i, i+1) = 1$.

A diagram over the mapping telescope is a sequence of types $P : \mathbb{N} \rightarrow \text{Type}$ together with arrows $f_n : P_n \rightarrow P_{n+1}$:

$$P_0 \xrightarrow{f_0} P_1 \xrightarrow{f_1} \dots$$

What we would like now would be to define the colimits of these diagrams over graphs, that would satisfy type theoretic versions of usual properties: it should make the diagram commute, and be universal with respect to this property. From now on, let G be a graph and D a diagram over this graph.

The commutation of the diagram is easy: the colimit should be the tip of a cocone.

Definition 33 : Cocone

Let Q be a type. A cocone over D into Q is the data of arrows $q_i : D_i \rightarrow Q$, and for any $i, j : G$ and $g : G(i, j)$, an homotopy $q_j \circ D(g) \sim q_i$.

If Q and Q' are type with an arrow $f : Q \rightarrow Q'$, and if C is a cocone over D into Q , one can easily build a cocone on D into Q' by postcomposing all maps of the cocone by f , giving a map

$$\text{postcompose}_{\text{cocone}} : \text{cocone}_D(Q) \rightarrow (Q' : \text{Type}) \rightarrow (Q \rightarrow Q') \rightarrow \text{cocone}_D(Q')$$

The other way around (from a cocone into Q' , give a map $Q \rightarrow Q'$) is exactly the second condition we seek:

Definition 34 : Universality of a cocone

Let Q be a type, and C be a cocone over D into Q . C is said universal if for any type Q' , $\text{postcompose}_{\text{cocone}}(C, Q')$ is an equivalence.

We can finally define what it means for Q to be a colimit of D .

Definition 35 : Colimit

A type Q is said to be a colimit of D if there is a cocone C over D into Q , which is universal.

EXAMPLE – Let A, B be types and $f, g : A \rightarrow B$. Let Q be the HIT generated by

$$\left| \begin{array}{l} q : B \rightarrow Q \\ \alpha : q \circ f \sim q \circ g \end{array} \right.$$

. Then Q is a colimit of the coequalizer diagram associated to A, B, f, g . We say that Q is a coequalizer of f and g .

Note that for any diagram D , one can build a free colimit of D , namely the higher inductive type $\text{colim}(D)$ generated by

$$\left| \begin{array}{l} \text{colim} : \prod_{i:G} D_i \rightarrow \text{colim}(D) \\ \alpha_{\text{colim}} : \prod_{ij:G} \prod_{g:G(i,j)} \prod_{x:D_i} \text{colim}_j \circ D(g) \sim \text{colim}_i \end{array} \right.$$

4.1.2 Properties of colimits

Diagrams can be thought as functors from type of graphs to Type, and hence one can define morphisms between diagrams as natural transformation.

Definition 36 : Morphism of diagram

Let D and D' be two diagrams over the same graph G . A morphism of diagram m between D and D' is the data of

- for all $i : G$, a map $m_i : D_i \rightarrow D'_i$
- for all $g : G(i, j)$, a path $D'(g) \circ m_i = m_j \circ D(g)$

This definition leads to obvious definition of identity morphism of diagrams, and composition of morphisms of diagrams.

Then, two diagrams D and D' are said to be equivalent if there is a morphism of diagrams m between D and D' such that all D_i are equivalences. It can easily be checked that the morphism m^{-1} given by inverting all m_i 's satisfies $m \circ m^{-1} = \text{id}$ and $m^{-1} \circ m = \text{id}$.

In the previous section, we defined the map $\text{postcompose}_{\text{cocone}}$ changing a cocone into Q and a map $Q \rightarrow Q'$ into a cocone into Q' . We now define a map $\text{precompose}_{\text{cocone}}$ taking a morphism of diagrams m between D and D' , and a cocone over D' into X , and giving a cocone over D into X :

Definition 37

Let m be a morphism of diagrams between D and D' , and X a type. Any cocone over D' into X can be changed into a cocone over D into X , by precomposing all maps in the cocone by the m_i 's:

$$\text{precompose}_{\text{cocone}} : \text{Hom}(D, D') \rightarrow \text{cocone}_{D'}(X) \rightarrow \text{cocone}_D(X).$$

Precomposition by a morphism of diagrams and postcomposition by a morphism are compatible with composition and identities, in the sense that

$$\begin{aligned} \text{precompose}_{\text{cocone}}(m \circ m') &= (\text{precompose}_{\text{cocone}} m) \circ (\text{precompose}_{\text{cocone}} m') \\ \text{precompose}_{\text{cocone}}(\text{idmap}) &= \text{idmap} \\ \text{postcompose}_{\text{cocone}}(\varphi \circ \varphi') &= (\text{postcompose}_{\text{cocone}} \varphi) \circ (\text{postcompose}_{\text{cocone}} \varphi') \\ \text{postcompose}_{\text{cocone}}(\text{idmap}) &= \text{idmap} \end{aligned}$$

These properties allows us to express functoriality properties of colimits. If m is a morphism between diagrams D and D' , then $\text{postcompose}_{\text{cocone}}^{-1} \circ \text{precompose}_{\text{cocone}}$ is a map $Q \rightarrow Q'$, where Q (resp. Q') is a colimit of D (resp. D'). One can even check that if m is an equivalence of diagrams, then the produced map $Q \rightarrow Q'$ is an equivalence of types. This is the lemma:

Lemma 38

Let D and D' be two equivalent diagrams, with respective colimits Q and Q' . Then $Q \simeq Q'$.

In particular, identity of diagrams being an equivalence, it asserts that the colimit of a diagram is unique (up to equivalence). From now on, as we supposed since chapter 2 the univalence axiom, we will say *the* colimit of a diagram.

One more interesting property of colimits is that it is stable by dependent sums. More precisely, let X be a type, and D_x a diagram over a graph G for all $x : X$. We want to link the colimits Q_x of diagrams D_x with the colimit of $\sum_{x:X} D_x$.

Definition 39 : Dependent diagram

Let G be a graph, X a type and D_x a diagram over G for all $x : X$. The diagram $\sum_{x:X} D_x$ is the diagram defined by

- $(\sum_{x:X} D_x)(i) = \sum_{x:X} D_x(i)$
- $(\sum_{x:X} D_x)(\phi) = (\text{idmap}, D_x(\phi))$

Note that a family C_x of cocones over D_x can be extended to a cocone $\sum_{x:X} C_x$ on $\sum_{x:X} D_x$. Then, the expected result is true

Proposition 40

If, for all $x : X$, Q_x is the colimit of D_x , then $\sum_{x:X} Q_x$ is the colimit of $\sum_{x:X} D_x$.

4.1.3 Truncated colimits

As said in the introduction, we now give a truncated version of colimits. Colimits actually behave well with respect to truncations. Indeed, if D is a diagram and P a colimit of D , then $\|P\|_n$ is the n -colimit of the n -truncated diagram $\|D\|_n$. Let's make it more precise.

Definition 41 : Truncation of a diagram

Let D be a diagram over a graph G , and n a truncation index. Then the diagram $\|D\|_n$ is the diagram over G defined by

- $(\|D\|_n)_0(i) \stackrel{\text{def}}{=} \|D_0(i)\|_n : \text{Type}_n$
- $(\|D\|_n)_1(\phi) \stackrel{\text{def}}{=} |D_1(\phi)|_n : \|D_0(i)\|_n \rightarrow \|D_0(j)\|_n$

Definition 42

Let D be a diagram over a graph G , P be a type, and C a cocone over D into P . C is said n -universal if for any $Q : \text{Type}_n$, $\text{postcompose}_{\text{cocone}}(C, Q)$ is an equivalence.

Then, P is said to be a n -colimit of D if there is a cocone C over D into P which is n -universal.

We can now give the fundamental proposition linking colimit and n -colimit.

Proposition 43

Let D be a diagram, and $P : \text{Type}$. Then, if P is a colimit of D , $\|P\|_n$ is a n -colimit of $\|D\|_n$.

The proof of this is really straightforward: a cocone over D into P can be changed equivalently into a cocone over $\|D\|_n$ into $\|P\|_n$, using the elimination principle 16 of truncations, and then we can show that the following diagram commutes for any $X : \text{Type}_n$

$$\begin{array}{ccc}
 \|P\|_n \rightarrow X & \longrightarrow & \text{cocone}(\|D\|_n, X) \\
 \downarrow \wr & & \uparrow \wr \\
 P \rightarrow X & \xrightarrow{\sim} & \text{cocone}(D, X)
 \end{array}$$

NOTA – This result does not hold for limits. If it were true, then applying it to the following equalizer diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[\lambda_{\perp}, y]{} \end{array} B$$

with $A, B : \text{Type}$, $f : A \rightarrow B$ and $y : B$ would lead to an equivalence

$$\left\| \sum_{a:A} f a = y \right\|_n \simeq \sum_{a:\|A\|_n} |f|_n a = |y|_n,$$

proving left-exactness of n -truncation.

4.1.4 Towards highly coherent colimits

In category theory, a diagram over a graph G in a category \mathcal{C} is a functor $G \rightarrow \mathcal{C}$. Our definition of diagrams doesn't reflect this definition, as we don't take care of composition of morphisms in the category (actually, we only define a map $G \rightarrow \mathcal{C}$). Diagrams over a category are defined in [HoTT, Exercise 7.16]. We will here define these diagrams, and the associated colimits.

NOTA – This new definition of diagrams is not sufficient to reflect what we would want to express. If we view homotopy type theory as ω -topos theory, what we want is rather a ω -functor from a graph into the ω -topos. That is an important open problem in homotopy type theory (even defining simplicial types is an open problem). It requires to handle an infinity of coherence levels, with finite data.

However, we could define, for any fixed n , diagrams and colimits coherent up to coherence level n (but that would be a real nightmare).

The main goal of this section is to define the colimit of the coequalizer with degeneracy δ

$$\begin{array}{c} \delta \\ \curvearrowright \\ A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{} \end{array} B \end{array}$$

and compositions $c_f : f \circ \delta = \text{idmap}$ and $c_g : g \circ \delta = \text{idmap}$. We will use this coequalizer to define the Boulier's construction in next section 4.2.

If we only take the previous definition of colimits, what we miss is the commutation of the following diagram

$$\begin{array}{ccc} q \circ f \circ \delta & \xrightarrow{q \circ c_f} & q \\ \alpha \circ \delta \downarrow & \nearrow q \circ c_g & \\ q \circ g \circ \delta & & \end{array}$$

Our idea to define the colimit of this diagram with degeneracy is to add it directly into the definition of the higher inductive type:

$$\left| \begin{array}{ll} \text{colim} & : B \rightarrow Q \\ \text{colim}_\alpha & : \prod_{x:A} f x = g x \\ \text{colim}_\alpha^\circ & : \prod_{y:B} \text{colim}_\alpha(\delta y) = c_f(y) \cdot c_g(y)^{-1} \end{array} \right.$$

NOTA – We indeed see here why this idea cannot be extended to a fully coherent notion of colimits: we cannot define a higher inductive type with an infinity of constructors. Hopefully, adding this first level of coherence will be sufficient for our applications.

The new constructor $\text{colim}_\alpha^\circ$ indeed asserts the commutation of the desired diagram. We can view this higher inductive type as the same type as the colimit over the coequalizer graph, but which does not add already existing paths.

One can easily (but very technically) check that properties in section 4.1.2 are still satisfied by this new notion of colimit.

4.2 Van Doorn's and Boulrier's constructions

In topos theory, there is a result that we would want to use in chapter 5:

Lemma 44 : [MM92, p. IV.7.8]

In a topos \mathcal{E} , if $f \in \text{Hom}_{\mathcal{E}}(A, B)$ is an epimorphism, then the colimit of

$$A \times_B A \rightrightarrows A$$

is B . The pullback $A \times_B A$ is called the kernel pair of f .

Unfortunately, this result fails in higher topos ; the kernel pair should be replaced by the Čech nerve of f .

$$\dots \rightrightarrows A \times_B A \times_B A \rightrightarrows A \times_B A \rightrightarrows A \xrightarrow[\text{colim}]{f} B$$

The issue we face in homotopy type theory is that the definition of the Čech nerve, and in general of simplicial objects is a hard open problem. It involves an infinite tower of coherences, and we do not know how to handle this. However, there is a way to define a diagram depending on a map f , which colimit is $\text{Im}(f)$.

The starting point of the construction is Floris Van Doorn's construction of proposition truncation [Van16].

Proposition 45 : Van Doorn's construction

Let $A : \text{Type}$. We define the higher inductive type TA as the coequalizer of

$$A \times A \xrightarrow[\pi_2]{\pi_1} A.$$

The colimit of the diagram

$$A \xrightarrow{q} TA \xrightarrow{q} TTA \xrightarrow{q} TTTA \xrightarrow{q} \dots$$

is $\|A\|_{-1}$.

Let's compare the direct definitions of $\|A\|_{-1}$ and TA

$$\|A\|_{-1} \left| \begin{array}{l} \text{tr} : A \rightarrow \|A\|_{-1} \\ \alpha_{\text{tr}} : \prod_{x,y:\|A\|_{-1}} x = y \end{array} \right. \quad TA \left| \begin{array}{l} q : A \rightarrow TA \\ \alpha : \prod_{x,y:A} qx = qy \end{array} \right.$$

The definitions are almost the same, except that the path constructor of TA quantifies over A , while the one of $\|A\|_{-1}$ quantifies over $\|A\|_{-1}$ itself: such a higher inductive type is a *recursive* inductive type. Thus, proposition 45 allows us to build the truncation in a non-recursive way. The counterpart is that we have to iterate the construction. In [Bou16], we found a way to generalize a bit this result. The main idea is that iterating the kernel pair construction will result in a diagram of colimit $\text{Im}(f)$.

Now, let $A, B : \text{Type}$ and $f : A \rightarrow B$. We define the kernel pair of f as the coequalizer of

$$A \times_B A \xrightarrow[\pi_2]{\pi_1} A.$$

In other words, $\text{KP}(f)$ is the higher inductive type generated by

$$\left| \begin{array}{l} \text{kp} : A \rightarrow \text{KP}(f) \\ \alpha : \prod_{x,y:A} fx = fy \rightarrow kpx = kp y \end{array} \right.$$

Using the eliminator of coequalizers, one can build a map $\widehat{f} : \text{KP}(f) \rightarrow B$, such that the following commutes

$$\begin{array}{ccc} A & \xrightarrow{\text{kp}} & \text{KP}(f) \\ & \searrow f & \downarrow \widehat{f} \\ & & B \end{array}$$

Then we can build $\text{KP}(\widehat{f})$ and build a map $\widehat{\widehat{f}} : \text{KP}(\widehat{f}) \rightarrow B$, etc. We have the following result

Proposition 46 : Boulier's construction

For any $f : A \rightarrow B$, $\text{Im}(f)$ is the colimit of the iterated kernel pair diagram of f

$$A \longrightarrow \text{KP}(f) \longrightarrow \text{KP}(\widehat{f}) \longrightarrow \text{KP}(\widehat{\widehat{f}}) \longrightarrow \dots$$

In particular, if f is a surjection, the colimit of this diagram is B .

Proof. The main idea of the proof is the equivalence between the diagrams

$$A \longrightarrow \text{KP}(f) \longrightarrow \text{KP}(\widehat{f}) \longrightarrow \dots$$

$$\sum_{y:B} \text{fib}_f(y) \longrightarrow \sum_{y:B} T(\text{fib}_f(y)) \longrightarrow \sum_{y:B} TT(\text{fib}_f(y)) \longrightarrow \dots$$

Let's begin by showing the first non-trivial equivalence:

$$s : \text{KP}(f) \simeq \sum_{y:B} T(\text{fib}_f(y)). \quad (4.1)$$

$\text{KP}(f)$ is the colimit of $A \times_B A \xrightarrow[\pi_2]{\pi_1} A$, and $\sum_{y:B} T(\text{fib}_f(y))$ is the colimit of $\sum_{y:B} \text{fib}_f(y) \times \text{fib}_f(y) \xRightarrow{\quad} \sum_{y:B} \text{fib}_f(y)$. As the two diagrams are equivalent, their colimits are equivalent. We will need the following fact, easily checked

$$\pi_1 \circ s = \widehat{f}.$$

Now, let's prove the other equivalences. We need the following lemma

Lemma 47

Let $X, Y : \text{Type}$ and $\varphi : X \rightarrow Y$. Then $\text{fib}_{\widehat{\varphi}}(y) \simeq T(\text{fib}_{\varphi}(y))$.

Proof. We have the following sequence of equivalences:

$$\begin{aligned} \text{fib}_{\widehat{\varphi}}(y) &\stackrel{\text{def}}{=} \sum_{x:\text{KP}(\varphi)} \widehat{\varphi} x = y \\ &\simeq \sum_{x:\sum_{y:B} T(\text{fib}_{\varphi}(y))} \widehat{\varphi} \circ s^{-1}(x) = y \quad \text{by 4.1} \\ &\simeq \sum_{x:\sum_{y:B} T(\text{fib}_{\varphi}(y))} \pi_1(x) = y \\ &\simeq T(\text{fib}_{\varphi}(y)) \end{aligned}$$

◇

Then, using the sum-of-fibers property, we can change the iterated kernel pair of f into

$$\sum_{y:B} \text{fib}_f(y) \longrightarrow \sum_{y:B} \text{fib}_{\widehat{f}}(y) \longrightarrow \sum_{y:B} \text{fib}_{\widehat{\widehat{f}}}(y) \longrightarrow \dots$$

With the just proved lemma, and a bit of induction, we can prove the desired equivalence of diagrams.

As colimits are stable under dependent sum, we know that the colimit of the diagram is thus $\sum_{y:B} Qy$, where Q is the colimit of

$$\text{fib}_f(y) \longrightarrow T(\text{fib}_f(y)) \longrightarrow TT(\text{fib}_f(y)) \longrightarrow \dots$$

But proposition 45 asserts that $Q \simeq \|\text{fib}_f(y)\|_{-1}$, and the result is proved. \square

The main issue with Van Doorn's construction is that it does not preserve truncations levels at all. For example, when computing $\|\mathbf{1}\|_{-1}$, the first step is $T\mathbf{1} \simeq S^1$, which is a 1-type. Asking for preservation of all truncation levels along the diagram might be too much, but the least we could ask is that when starting with a $P : \mathbf{HProp}$, the diagram should be the constant diagram $P \rightarrow P \rightarrow P \rightarrow \dots$. The Boulier counterpart of this is, when starting with an embedding f , all \widehat{f} are embeddings.

This can be achieved by changing operators T and KP , by asking them to preserve identities:

$$TA \left| \begin{array}{l} q : A \rightarrow TA \\ \alpha : \prod_{x,y:A} qx = qy \\ \alpha_1 : \prod_{x:A} \alpha(x,x) = 1 \end{array} \right. ; \quad KP(f) \left| \begin{array}{l} kp : A \rightarrow KP(f) \\ \alpha : \prod_{x,y:A} fx = fy \rightarrow qx = qy \\ \alpha_1 : \prod_{x:A} \alpha(x,x,1) = 1 \end{array} \right.$$

This new KP can be thought of as the coequalizer preserving the “degeneracy” δ , as in section 4.1.4:

$$\begin{array}{c} \delta \\ \curvearrowright \\ A \times_B A \xrightarrow[\pi_2]{\pi_1} A \end{array}$$

Then, the following is still true

Proposition 48 : Boulier's construction

For any $f : A \rightarrow B$, $\text{Im}(f)$ is the colimit of the iterated kernel pair diagram of f

$$A \longrightarrow KP(f) \longrightarrow KP(\widehat{f}) \longrightarrow KP(\widehat{\widehat{f}}) \longrightarrow \dots$$

Moreover, if f is an embedding, \widehat{f} also is.

The proof is almost the same as for proposition 46, except that the new constructors in the higher inductive types introduce another level of coherence, which is very technical to handle.

4.3 Formalization

Except the next section 4.4, all this chapter has been formalized in Coq, mostly by Simon Boulrier, and the library is available at <https://github.com/SimonBoulrier/hott-colimits>.

Formalization of colimits: what could we say ?...

4.4 Towards groupoid objects

Defining groupoid objects in homotopy type theory is an open problem, just as defining simplicial or even semisimplicial types. We will just give here some of our thoughts about this problem.

The first thing to say is that defining a groupoid object in type theory is exactly the same problem as defining a highly coherent equivalence relation. At the moment, the only fully coherent equivalence relation we know in homotopy type theory is the equality, which we can slightly modify to create another ones. For example, if $f : A \rightarrow B$, the relation $f x = f y$ defines an equivalence relation on A . Actually, when seeing homotopy type theory as the language of ω -topoi, the Giraud-Rezk-Lurie axioms in [HTT] asserts that these are the only equivalence relation we can define.

We will call a *uncoherent equivalence relation*, or UER, on a type A any symmetric, reflexive and transitive relation R on A , i.e R is a UER on A if

1. For all $x, y : A$, $R(x, y) : \text{Type}$
2. For all $x, y : A$, $\text{Sym}_R(x, y) : R(x, y) \rightarrow R(y, x)$
3. For all $x : A$, $\text{Refl}_R(x) : R(x, x)$
4. For all $x, y, z : A$, $\text{Trans}_R(x, y, z) : R(x, y) \rightarrow R(y, z) \rightarrow R(x, z)$

Then, if we want to mimic the behaviour of equalities, we can think of an equivalence relation on A as the data of an UER R_0 on A , and for all $x, y : A$, an UER $R_1^{x,y}$ on $R(x, y)$, and for all $u, v : R(x, y)$, an UER $R_2^{u,v}$ on $R_1(u, v)$, etc. This can be formalized by a coinductive type $R : \text{ER}(A)$ with constructors

- R_0 is an UER on A
- $R_1 : \prod_{x,y:A} \text{ER}(R_0(x, y))$

The issue with such a definition is that all these uncoherent equivalence relation have to be coherent with each other, for example we want

$$R_1^{x,x}(\text{Trans}_{R_0}(x, x, x) \text{Refl}_{R_0}(x) \text{Refl}_{R_0}(x), \text{Refl}_{R_0}(x))$$

Adding all these desired coherences by hand is not an option, as there are an infinity of them.

The next idea is to use the Boulier's construction. Let R_0 be an UER on a type A . We define then the higher inductive type Ξ_0 generated by

$$\left| \begin{array}{ll} \xi_0 & : A \rightarrow \Xi_0 \\ \alpha_0 & : \prod_{x,y:A} R_0(x,y) \rightarrow \xi_0 x = \xi_0 y \\ \alpha_0^{\text{Refl}} & : \prod_{x:A} \alpha_0(x,x, \text{Refl}_{R_0}(x)) = 1 \\ \alpha_0^{\text{Sym}} & : \prod_{x,y:A} \prod_{u:R_0(x,y)} \alpha_0(y,x, \text{Sym}_{R_0}(u)) = (\alpha_0(x,y,u))^{-1} \\ \alpha_0^{\text{Trans}} & : \prod_{x,y,z:A} \prod_{u:R_0(x,y)} \prod_{v:R_0(y,z)} \alpha_0(x,z, \text{Trans}_{R_0}(u,v)) = \alpha_0(x,y,u) \cdot \alpha_0(y,z,v) \end{array} \right.$$

Ξ_0 can be seen as the quotient type of A by R_0 . In other words, Ξ_0 is the same type as A , but where R_0 has been changed by equalities. Then, an equivalence relation on A could be defined coinductively as an UER R_0 on A together with an equivalence relation on Ξ_0 .

Our hope is that the Boulier's construction is still valid in this context, *i.e* that if we consider the colimit Ξ_ω of the diagram

$$\begin{array}{ccccc} A & \xrightarrow{\quad} & \Xi_0 & \xrightarrow{\quad} & \Xi_1 & \xrightarrow{\quad} & \cdots \\ & \searrow \zeta & \downarrow \zeta_0 & & \downarrow \zeta_1 & \nearrow & \\ & & \Xi_\omega & & & & \end{array}$$

then the equivalence relation defined by (R_i) is equivalent to the equivalence relation on A defined by $x \sim y \stackrel{\text{def}}{=} \zeta x = \zeta y$.

Define groupoid objects in HoTT, then write it here

5 Sheaves in homotopy type theory

Reductio ad absurdum, which
Euclid loved so much, is one of a
mathematician's finest weapons.
It is a far finer gambit than any
chess gambit: a chess player may
offer the sacrifice of a pawn or
even a piece, but a mathematician
offers the game

Godfrey Harold Hardy

Section 5.1 briefly recalls the topos-theoretic version of Lawvere-Tierney sheaves theory, and section 5.2 presents the main result of this thesis: the construction of the sheafification modality in homotopy type theory. Section 5.4 tries to link our construction with forcing in type theory.

5.1 Sheaves in topoi

In this section, we will rather work in an arbitrary topos rather in type theory. The next section will present a generalisation of the results presented here.

Let us fix for the whole section a topos \mathcal{E} , with subobject classifier Ω . A *Lawvere-Tierney topology* on \mathcal{E} is a way to modify slightly truth values of \mathcal{E} . It allows to speak about *locally true* things instead of *true* things.

Definition 49 : Lawvere-Tierney topology [MM92]

A *Lawvere-Tierney topology* is an endomorphism $j : \Omega \rightarrow \Omega$ preserving \top ($j \top = \top$), idempotent ($j \circ j = j$) and commuting with products ($j \circ \wedge = \wedge \circ (j, j)$).

A classical example of Lawvere-Tierney topology is given by double negation. Other examples are given by Grothendieck topologies, in the sense

Theorem 50 : [MM92, p. V.1.2]

Every Grothendieck topology J on a small category \mathbf{C} determines a Lawvere-Tierney topology j on the presheaf topos $\mathbf{Sets}^{\mathbf{C}^{\text{op}}}$.

Any Lawvere-Tierney topology j on \mathcal{E} induces a closure operator $A \mapsto \overline{A}$ on subobjects. If we see a subobject A of E as a characteristic function χ_A , the closure \overline{A} corresponds to the subobject of E whose characteristic function is

$$\chi_{\overline{A}} = j \circ \chi_A.$$

A subobject A of E is said to be dense when $\overline{A} = E$.

Then, we are interested in objects of \mathcal{E} for which it is impossible to make a distinction between objects and their dense subobjects, *i.e* for which “true” and “locally true” coincide. Such objects are called *sheaves*, and are defined as

Definition 51 : Sheaves[MM92, p. V.2]

On object F of \mathcal{E} is a sheaf (or j -sheaf) if for every dense monomorphism $m : A \hookrightarrow E$ in \mathcal{E} , the canonical map $\text{Hom}_{\mathcal{E}}(E, F) \rightarrow \text{Hom}_{\mathcal{E}}(A, F)$ is an isomorphism.

One can show that $\text{Sh}_{\mathcal{E}}$, the full sub-category of \mathcal{E} given by sheaves, is again a topos, with classifying object

$$\Omega_j = \{P \in \Omega \mid jP = P\}.$$

Lawvere-Tierney sheafification is a way to build a left adjoint \mathbf{a}_j to the inclusion $\mathcal{E} \hookrightarrow \text{Sh}(\mathcal{E})$, exhibiting $\text{Sh}(\mathcal{E})$ as a reflective subcategory of \mathcal{E} . In particular, that implies that logical principles valid in \mathcal{E} are still valid in $\text{Sh}(\mathcal{E})$.

For any object E of \mathcal{E} , $\mathbf{a}_j(E)$ is defined as in the following diagram

$$\begin{array}{ccc} E & \xrightarrow{\{\cdot\}_E} & \Omega^E \\ \theta_E \downarrow & & \downarrow j^E \\ E' & \xrightarrow{\quad} & \Omega_j^E \\ \text{closure} \searrow & & \nearrow \\ & \mathbf{a}_j(E) & \end{array}$$

The proof that \mathbf{a}_j defines a left adjoint to the inclusion can be found in [MM92].

One classical example of use of sheafification is the construction, from any topos, of a boolean topos negating the continuum hypothesis. More precisely:

Theorem 52 : Negation of CH [MM92, p. VI.2.1]

There exists a Boolean topos satisfying the axiom of choice, in which the continuum hypothesis fails.

The proof actually follows almost exactly the famous proof of the construction by Paul Cohen of a model of ZFC negating the continuum hypothesis [CD66]. Together with the model of constructible sets \mathfrak{L} by Kurt Gödel [Göd40], it proves that CH is independent of ZFC, solving first Hilbert’s problem.

5.2 Sheaves in homotopy type theory

The idea of this section is to consider sheafification in topoi as only the first step towards sheafification in type theory. We note that axioms for a Lawvere-Tierney topology on the subobject classifier Ω of a topos are very close to those of a modality on Ω . We will extensively use this idea, applying it to every subobject classifier Type_n we described in 2.4. The subobject classifier Ω of a topos is seen as the *truth values* of the topos, which corresponds to the type HProp in our setting ; the topos is considered proof irrelevant, corresponding to our HSet . Sheafification in topoi is thus a way, when translated to the setting of homotopy type theory, to build, from a left-exact modality on HProp , a left-exact modality on HSet . Our hope in this section is to iterate this construction by applying it to the subobject classifier HSet equipped with a left-exact modality, to build a new left-exact modality on Type_1 , and so on.

The first thing we can note is that such a construction will not allow to reach every type: it is known that there exists types with no finite truncation level [HoTT, Example 8.8.6]. Even worse, some types are not even the limit of its successive truncations, even in a hypercomplete setting [MV98]. It suggests that defining a sheafification functor for all truncated types won't give (at least easily) a sheafification functor on whole Type . Another issue that can be pointed is the complexity of proofs. If, in a topos-theoretic setting, everything is proof-irrelevant, it won't be the case for higher settings, forcing us to prove results that were previously true on the nose. This will oblige us to write long and technical proofs of coherence, and more deeply, to modify completely some lemmas, such as Proposition [MM92, p. IV.7.8], stating that epimorphisms are coequalizers of their kernel pair.

The main idea is thus to follow as closely as possible the topos-theoretic construction, and change as few times as possible to make it work in our higher setting.

Note that the principles we want to add are added directly from the HProp level, the extension to all truncated types is automatic. The choice of the left-exact modality on HProp is thus crucial. For the rest of the section, we fix one, note \circ_{-1} . The reader can think of the double negation \circ_{-n} defined in 3.2.3. We will define, by induction on the truncation level, left-exact modalities on all Type_n , as in the following theorem.

Theorem 53

The sequence defined by induction by

$$\begin{aligned} \circ &: \forall (n : \text{nat}), \text{Type}_n \rightarrow \text{Type}_n \\ \circ_{-1} (T) &\stackrel{\text{def}}{=} j T \\ \circ_{n+1} (T) &\stackrel{\text{def}}{=} \sum_{u: T \rightarrow \text{Type}_n^{\circ}} \circ_{-1} \left\| \sum_{a: T} u = (\lambda t, \circ_n(a = t)) \right\| \end{aligned}$$

defines a sequence of left-exact modalities, coherent with each others in the

sense that the following diagram commutes for any $P : \text{Type}_n$, where \hat{P} is P seen as an inhabitant of Type_{n+1} .

$$\begin{array}{ccc} P & \xrightarrow{\sim} & \hat{P} \\ \eta_n \downarrow & & \downarrow \eta_{n+1} \\ \circ_n P & \xrightarrow{\sim} & \circ_{n+1} \hat{P} \end{array}$$

5.2.1 Sheaf theory

Let n be a truncation index greater than -1 , and \circ_n be the left-exact modality given by our induction hypothesis. As in the topos-theoretic setting, we will define what it means for a type to be a n -sheaf (or just “sheaf” if the context is clear), and consider the reflective subuniverses of these sheaves; the reflector will exactly be the sheafification functor. The main issue to give the “good” definition is the choice of the subobject classifier in which dense subobjects will be chosen: two choices appears, HProp and Type_n ; we will actually use both. What guided our choice is the crucial property that the type of all n -sheaves has to be a $(n+1)$ -sheaf.

From the modality \circ_n , one can build a *closure operator*.

Definition 54

Let E be a type.

- The closure of a subobject of E with n -truncated homotopy fibers (or n -subobject of E , for short), classified by $\chi : E \rightarrow \text{Type}_n$, is the subobject of E classified by $\circ_n \circ \chi$.
- An n -subobject of E classified by χ is said to be closed in E if it is equal to its closure, i.e if $\chi = \circ_n \circ \chi$.
- An n -subobject of E classified by χ is said to be dense in E if its closure is E , i.e if $\circ_n \circ \chi = \lambda e, 1$

Topos-theoretic sheaves are characterized by a property of existence and uniqueness, which will be translated, as usual, into a proof that a certain function is an equivalence.

Definition 55 : Restriction

Let $E, F : \text{Type}$ and $\chi : E \rightarrow \text{Type}$. We define the restriction map Φ_E^χ as follows

$$\Phi_E^\chi : \begin{array}{ccc} E \rightarrow F & \longrightarrow & \sum_{e:E} \chi e \rightarrow F \\ f & \longmapsto & f \circ \pi_1 \end{array} .$$

Here, we need to distinguish between dense (-1) -subobjects, that will be used in the definition of sheaves, and dense n -subobjects, that will be used in the definition of separated types.

Definition 56 : Separated Type

A type F in Type_{n+1} is separated if for any type E , and all dense n -subobject of E classified by χ , Φ_E^χ is an embedding.

With topos theory point of view, it means that given a map $\sum_{e:E} \chi e \rightarrow F$, if there is an extension $\tilde{f} : E \rightarrow F$, then it is unique, as in

$$\begin{array}{ccc} \sum_{e:E} \chi e & \xrightarrow{f} & F \\ \pi_1 \downarrow & \nearrow \tilde{f} & \\ E & & \end{array}$$

Definition 57 : Sheaf

A type F of Type_{n+1} is a $(n+1)$ -sheaf if it is separated, and for any type E and all dense (-1) -subobject of E classified by χ , Φ_E^χ is an equivalence.

In topos-theoretic words, it means that given a map $f : \sum_{e:E} \chi e \rightarrow F$, one can extend it uniquely to $\tilde{f} : E \rightarrow F$, as in

$$\begin{array}{ccc} \sum_{e:E} \chi e & \xrightarrow{f} & F \\ \pi_1 \downarrow & \nearrow \tilde{f} & \\ E & & \end{array}$$

Note that these definitions are almost the same as the ones in [MM92]. The main difference is that separated is defined for n -subobjects, while sheaf only for (-1) -subobjects. It might seem bizarre to make such a distinction, but the following proposition gives a better understanding of the situation.

Proposition 58

A type F is Type_{n+1} is separated if, and only if all its path types are n -modal, ie

$$\prod_{x,y:F} (\odot_n(x=y)) = (x=y).$$

A $(n+1)$ -sheaf is hence just a type satisfying the usual property of sheaves (i.e existence of uniqueness of arrow extension from dense (-1) -subobjects), with the condition that all its path types are n -sheaves. It is a way to force the compatibility of the modalities we are defining.

One can check that the property `IsSeparated` (resp. `IsSheaf`) is `HProp`: given a $X : \text{Type}_{n+1}$, there is at most one way for it to be separated (resp. a sheaf). In particular, when needed to prove equality between two sheaves, it suffices to show the equality between the underlying types.

As said earlier, these definitions allow us to prove the fundamental property that the type of all n -sheaves is itself a $(n+1)$ -sheaf (this can be viewed as an equivalent definition of left-exactness).

Proposition 59

Type_n° is a $(n+1)$ -sheaf.

Proof. We have two things to prove here : separation, and sheafness.

- Let $E : \text{Type}$ and $\chi : E \rightarrow \text{Type}$, dense in E . Let $\phi_1, \phi_2 : E \rightarrow \text{Type}_n^\circ$, such that $\phi_1 \circ \pi_1 = \phi_2 \circ \pi_1$ and let $x : E$. We show $\phi_1(x) = \phi_2(x)$ using univalence.

As χ is dense, we have a term $m_x : \circ_n(\chi x)$. But as $\phi_2(x)$ is modal, we can obtain a term $h_x : \chi x$. As ϕ_1 and ϕ_2 are equal on $\sum_{e:E} \chi e$, we have an arrow $\phi_1(x) \rightarrow \phi_2(x)$. The same method leads to an arrow $\phi_2(x) \rightarrow \phi_1(x)$, and one can prove that they are each other inverse.

- Now, we prove that Type_n° is a sheaf. Let $E : \text{Type}$ and $\chi : E \rightarrow \text{HProp}$, dense in E . Let $f : \sum_{e:E} \chi e \rightarrow \text{Type}_n^\circ$. We want to extend f into a map $E \rightarrow \text{Type}_n^\circ$.

We define g as $g(e) = \circ_n(\text{fib}_\phi(e))$, where

$$\phi : \sum_{b : \sum_{e:E} \chi e} (f b) \rightarrow E$$

defined by $\phi(x) = (x_1)_1$. Using the following lemma, one can prove that the map $f \mapsto g$ defines an inverse of Φ_E^χ .

Lemma 60

Let $A, B, C : \text{Type}_n$, $f : A \rightarrow B$ and $g : B \rightarrow C$. Then if all fibers of f and g are n -truncated, then

$$\left(\prod_{c:C} \circ_n(\text{fib}_{g \circ f}(c)) \right) \simeq \circ_n \left(\sum_{w:\text{fib}_g(c)} \circ_n(\text{fib}_f(w_1)) \right).$$

Proof. This is just a modal counterpart of the property characterizing fibers of composition of function. \diamond

□

Another fundamental property on sheaves we will need is that the type of (dependent) functions is a sheaf as soon as its codomain is a sheaf.

Proposition 61

If $A : \text{Type}_{n+1}$ and $B : A \rightarrow \text{Type}_{n+1}$ such that for any $a : A$, $(B\ a)$ is a sheaf, then $\prod_{a:A} B\ a$ is a sheaf.

Proof. Again, when proving equivalences, we will only define the maps. The proofs of section and retraction are technical, not really interesting, and present in the formalisation.

- *Separation:* Let $E : \text{Type}$ and $\chi : E \rightarrow \text{Type}_n$ dense in E . Let $\phi_1, \phi_2 : E \rightarrow \prod_{a:A} B\ a$ equal on $\sum_{e:E} \chi\ e$ i.e. such that $\phi_1 \circ \pi_1 = \phi_2 \circ \pi_1$. Then for any $a : A$, $(\lambda x : E, \phi_1(x, a))$ and $(\lambda x : E, \phi_2(x, a))$ coincide on $\sum_{e:E} (\chi\ e)$, and as $B\ a$ is separated, they coincide also on all E .
- *Sheaf:* Let $E : \text{Type}$, $\chi : E \rightarrow \text{HProp}$ dense in E and $f : \sum_{e:E} \chi\ e \rightarrow \prod_{a:A} B\ a$. Let $a : A$; the map $(\lambda x : E, f(x, a))$ is valued in the sheaf $B\ a$, so it can be extended to all E , allowing f to be extended to all E .

□

5.2.2 Sheafification

The sheafification process will be defined in two steps. The first one will build, from any $T : \text{Type}_{n+1}$, a separated object $\square_{n+1} T : \text{Type}_{n+1}$; one can show that \square_{n+1} defines a modality on Type_{n+1} . The second step will build, from any separated type $T : \text{Type}_{n+1}$, a sheaf $\circ_{n+1}(T)$; one can show that \circ_{n+1} is indeed the left-exact modality we are searching.

Let n be a fixed truncation index, and \circ_n a left-exact modality on Type_n , compatible with \circ_{-1} as in

Condition 62

For any mere proposition P (where \widehat{P} is P seen as a Type_n), $\circ_n \widehat{P} = \circ_{-1} P$ and the following coherence diagram commutes

$$\begin{array}{ccc} P & \xrightarrow{\sim} & \widehat{P} \\ \eta_{-1} \downarrow & & \downarrow \eta_n \\ \circ_{-1} P & \xrightarrow{\sim} & \circ_n \widehat{P} \end{array}$$

From types to separated types

Let $T : \text{Type}_{n+1}$. We define $\square_{n+1} T$ as the image of $\circ_n^T \circ \{\cdot\}_T$, as in

$$\begin{array}{ccc} T & \xrightarrow{\{\cdot\}_T} & (\text{Type}_n)^T, \\ \mu_T \downarrow & & \downarrow \circ_n^T \\ \square_{n+1} T & \longrightarrow & (\text{Type}_n^\circ)^T \end{array}$$

where $\{\cdot\}_T$ is the singleton map $\lambda(t : T), \lambda(t' : T), t = t'$. $\square_{n+1} T$ can be given explicitly by

$$\begin{aligned} \square_{n+1} T &\stackrel{\text{def}}{=} \text{Im}(\lambda t : T, \lambda t', \circ_n(t = t')) \\ &\stackrel{\text{def}}{=} \sum_{u : T \rightarrow \text{Type}_n^\circ} \left\| \sum_{a : A} (\lambda t, \circ_n(a = t)) = u \right\|. \end{aligned}$$

This corresponds to the free separated object used in the topos-theoretic construction, but using Type_n° instead of the j -subobject classifier Ω_j .

Proposition 63

For any $T : \text{Type}_{n+1}$, $\square_{n+1} T$ is separated.

Proof. We use the following lemma:

Lemma 64

A $(n+1)$ -truncated type T with an embedding $f : T \rightarrow U$ into a separated $(n+1)$ -truncated type U is itself separated.

Proof. Let $E : \text{Type}$ and $\chi : E \rightarrow \text{Type}_n$ dense in E . Let $\phi_1, \phi_2 : \sum_{e:E} \chi e \rightarrow T$ such that $\phi_1 \circ \pi_1 \sim \phi_2 \circ \pi_1$. Postcomposing by f yields an homotopy $f \circ \phi_1 \circ \pi_1 \sim f \circ \phi_2 \circ \pi_1$. As $f \circ \phi_1, f \circ \phi_2 : \sum_{e:E} \chi e \rightarrow U$, and U is separated, we can deduce $f \circ \phi_1 \sim f \circ \phi_2$. As f is an embedding, $\phi_1 \sim \phi_2$. \diamond

As $\square_{n+1} T$ embeds in $(\text{Type}_n^\circ)^T$, we only have to show that the latter is separated. But it is the case because Type_n° is a sheaf (by Proposition 59) and a function type is a sheaf as soon as its codomain is a sheaf (by Proposition 61). \square

We will now show that \square_{n+1} defines a modality, with unit map μ . The left-exactness of \square_{n+1} will come from the second part of the process. The first thing to show that $\square_{n+1} T$ is universal among separated type below T . In the topos-theoretic sheafification, it comes easily from the fact that epimorphisms are coequalizers of their kernel pairs. As it is not true anymore in our setting,

we will use its generalization, proposition 48. Here is a sketch of the proof: as μ_T is a surjection (it is defined by the surjection-embedding factorization), $\square_{n+1} T$ is the colimit of its iterated kernel pair. Hence, for any type Q defining a cocone on $\text{KP}(\mu_T)$, there is a unique arrow $\square_{n+1} T \rightarrow Q$. What remains to show is any separated type Q defines a cocone on $\text{KP}(\mu_T)$; we will actually show that any separated type Q defines a cocone on $\|\text{KP}(\mu_T)\|_{n+1}$, which is enough. We do it by defining another diagram \hat{T} , equivalent to $\|\text{KP}(\mu_T)\|_{n+1}$, for which it is easy to define a cocone into any separated type Q .

This comes from the following construction which connects $\square_{n+1} T$ to the colimit of the iterated kernel pair of μ_T .

Definition 65

Let $X : \text{Type}$. Let \hat{T}_X be the higher inductive type generated by

- $\mathfrak{i} : \|X\|_{n+1} \rightarrow \hat{T}_X$
- $\mathfrak{a} : \forall a, b : \|X\|_{n+1}, \circ(a = b) \rightarrow \mathfrak{i}(a) = \mathfrak{i}(b)$
- $\mathfrak{a}_1 : \forall a : \|X\|_{n+1}, \mathfrak{a}(a, a, \eta_{a=a} 1) = 1$

We view \hat{T} as the coequalizer of

$$\sum_{a, b : \|X\|_{n+1}} \circ(a = b) \xrightarrow[\pi_2]{\pi_1} \|X\|_{n+1}$$

preserving $\eta_{a=a} 1$.

We consider the diagram \hat{T} :

$$\|X\|_{n+1} \longrightarrow \|\hat{T}_X\|_{n+1} \longrightarrow \|\hat{T}_{\hat{T}_X}\|_{n+1} \longrightarrow \dots$$

The main result we want about \hat{T} is the following:

Lemma 66

Let $T : \text{Type}_{n+1}$. Then $\square_{n+1} T$ is the $(n+1)$ -colimit of the diagram \hat{T} .

The key point of the proof is that diagrams \hat{T} and $\|\text{KP}(\mu_T)\|_{n+1}$ are equivalent. We will need the following lemma:

Lemma 67

Let $A, S : \text{Type}_{n+1}$, S separated, and $f : A \rightarrow S$. Then if

$$\forall a, b : A, f(a) = f(b) \simeq \circ(a = b), \quad (5.1)$$

then

$$\forall a, b : \|\text{KP}_f\|_{n+1}, |\tilde{f}|_{n+1}(a) = |\tilde{f}|_{n+1}(b) \simeq \circ(a = b).$$

Sketch of proof. By induction on truncation, we need to show that

$$\forall a, b : \mathbf{KP}_f, \tilde{f}(|a|_{n+1}) = \tilde{f}(|b|_{n+1}) \simeq \circ(|a|_{n+1} = |b|_{n+1}).$$

We use the encode-decode [HoTT, Section 8.9] method to characterize $\tilde{f}(|a|_{n+1}) = x$, and the result follows. We refer to the formalization for details. \square

This lemma allows to prove that, in the iterated kernel pair diagram of f

$$\begin{array}{ccccccc} X & \longrightarrow & \mathbf{KP}(f) & \longrightarrow & \mathbf{KP}(f_1) & \longrightarrow & \mathbf{KP}(f_2) \longrightarrow \dots \\ & & \searrow f & & \downarrow f_2 & & \swarrow f_3 \\ & & & & S & & \end{array}$$

if f satisfies 5.1, then each $|f_i|_{n+1}$ does.

NOTA – It is clear that if A and B are equivalent types, and $\forall a, b : A, f(a) = f(b) \simeq \circ(a = b)$, then

$$\mathrm{Coeq}_1 \left(\sum_{a,b:A} f a = f b \xrightarrow[\pi_2]{\pi_1} A \right) \simeq \mathrm{Coeq}_1 \left(\sum_{a,b:B} \circ(a = b) \xrightarrow[\pi_2]{\pi_1} B \right)$$

Proof of lemma 66. As said, it suffices to show that $\|\mathbf{C}(\mu_T)\|_{n+1} = \mathring{T}$.

$$\begin{array}{ccccccc} \|\mathbf{KP}^0(\mu_T)\|_{n+1} & \longrightarrow & \|\mathbf{KP}^1(\mu_T)\|_{n+1} & \longrightarrow & \|\mathbf{KP}^2(\mu_T)\|_{n+1} & \longrightarrow & \dots \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr & & \\ \mathring{T}_0 & \longrightarrow & \mathring{T}_1 & \longrightarrow & \mathring{T}_2 & \longrightarrow & \dots \end{array}$$

The first equivalence is trivial. Let's then start with the second. What we need to show is

$$\|\mathbf{KP}(\mu_T)\|_{n+1} \simeq \|\mathring{T}_T\|_{n+1},$$

i.e

$$\mathrm{Coeq}_1 \left(\sum_{a,b:T} \mu_T a = \mu_T b \xrightarrow[\pi_2]{\pi_1} T \right) \simeq \mathrm{Coeq}_1 \left(\sum_{a,b:T} \circ(a = b) \xrightarrow[\pi_2]{\pi_1} T \right).$$

By the previous remark, it suffices to show that μ_T satisfies condition (5.1), *i.e* $\forall a, b : T, \circ_n(a = b) = (\mu_T a = \mu_T b)$. By univalence, we want arrows in both ways, forming an equivalence.

- Suppose $p : (\mu_T a = \mu_T b)$. Then projecting p along first components yields $q : \prod_{t:T} \circ_n(a = t) = \circ_n(b = t)$. Taking for example $t = b$, we deduce $\circ_n(a = b) = \circ_n(b = b)$, and the latter is inhabited by $\eta_{b=b}1$.

- Suppose now $p : \circ_n(a = b)$. Let ι be the first projection from $\square_{n+1} T \rightarrow (T \rightarrow \text{Type}_n^\circ)$. ι is an embedding, thus it suffices to prove $\iota(\mu_T a) = \iota(\mu_T b)$, i.e. $\prod_{t:T} \circ_n(a = t) = \circ_n(b = t)$. The latter remains true by univalence.

The fact that these two form an equivalence is technical, we refer to the formalization for an explicit proof.

Let's show the other equivalences by induction. Suppose that, for a given $i : \mathbb{N}$, $\|\text{KP}^i(\mu_T)\|_{n+1} \simeq \hat{T}_i$. We want to prove $\|\text{KP}^{i+1}(\mu_T)\|_{n+1} \simeq \hat{T}_{i+1}$, i.e.

$$\begin{aligned} & \left\| \text{Coeq}_1 \left(\sum_{a,b:\text{KP}^i(\mu_T)} f_i a = f_i b \xrightarrow[\pi_2]{\pi_1} \text{KP}^i(\mu_T) \right) \right\|_{n+1} \\ & \simeq \left\| \text{Coeq}_1 \left(\sum_{a,b:\|\hat{T}_i\|_{n+1}} \circ(a = b) \xrightarrow[\pi_2]{\pi_1} \|\hat{T}_i\|_{n+1} \right) \right\|_{n+1} \end{aligned}$$

where f_i is the map $\text{KP}^i(\mu_T) \rightarrow \square_{n+1} T$. But lemma 67 just asserted that f_i satisfies 5.1, hence the previous nota yields the result.

One would need to show that, modulo these equivalences, the arrows of the two diagrams are equal. We leave that to the reader, who can refer to the formalization if needed. \square

Now, let Q be any separated Type_{n+1} , and $f : X \rightarrow Q$. Then the following diagram commutes

$$\begin{array}{ccccccc} \|X\|_{n+1} & \longrightarrow & \|\hat{T}_X\|_{n+1} & \longrightarrow & \|\hat{T}_{\hat{T}_X}\|_{n+1} & \longrightarrow & \dots \\ & \searrow & \downarrow & \swarrow & & & \\ & & Q & & & & \end{array}$$

But we know (lemma 66) that $\square_{n+1} T$ is the $(n+1)$ -colimit of the diagram \hat{T} , thus there is an universal arrow $\square_{n+1} T \rightarrow Q$. This is enough to state the following proposition.

Proposition 68

(\square_{n+1}, μ) defines a reflective subuniverse on Type_{n+1} .

To show that \square_{n+1} is a modality, it remains to show that separation is a property stable under sigma-types. Let $A : \text{Type}_{n+1}$ be a separated type and $B : A \rightarrow \text{Type}_{n+1}$ be a family of separated types. We want to show that $\sum_{x:A} Bx$ is separated. Let E be a type, and $\chi : E \rightarrow \text{Type}_n$ a dense subobject of E .

Let f, g be two maps from $\sum_{e:E} \chi e$ to $\sum_{x:A} Bx$, equal when precomposed with π_1 .

$$\begin{array}{ccc}
 \sum_{e:E} \chi e & \xrightarrow[\substack{g \circ \pi_1 \\ f}]{f \circ \pi_1} & \sum_{x:A} Bx \\
 \text{dense} \downarrow & \nearrow \substack{f \\ g} & \\
 E & &
 \end{array}$$

We can restrict the previous diagram to

$$\begin{array}{ccc}
 \sum_{e:E} \chi e & \xrightarrow[\substack{\pi_1 \circ g \circ \pi_1 \\ \pi_1 \circ f}]{\pi_1 \circ f \circ \pi_1} & A \\
 \text{dense} \downarrow & \nearrow \substack{\pi_1 \circ f \\ \pi_1 \circ g} & \\
 E & &
 \end{array}$$

and as A is separated, $\pi_1 \circ f = \pi_1 \circ g$. For the second components, let $x : E$. Notice that $\sum_{y:E} x = y$ has a dense n -subobject, $\sum_{y:\sum_{e:E} \chi e} x = y_1$:

$$\begin{array}{ccc}
 \sum_{y:\sum_{e:E} \chi e} x = y_1 & \xrightarrow[\substack{\pi_2 \circ g \circ \pi_1 \circ \pi_1 \\ \pi_2 \circ f \circ \pi_1}]{\pi_2 \circ f \circ \pi_1 \circ \pi_1} & Bx \\
 \text{dense} \downarrow & \nearrow \substack{\pi_2 \circ f \circ \pi_1 \\ \pi_2 \circ g \circ \pi_1} & \\
 \sum_{y:E} x = y & &
 \end{array}$$

Using the separation property of Bx , one can show that second components, transported correctly along the first components equality, are equal. The complete proof can be found in the formalization. This proves the following proposition

Proposition 69

(\Box_{n+1}, μ) defines a modality on Type_{n+1} .

As this modality is just a step in the construction, we do not need to show that it is left exact (actually, it is not), we will only do it for the sheafification modality.

From Separated Type to Sheaf

For any T in Type_{n+1} , $\circ_{n+1} T$ is defined as the closure of $\Box_{n+1} T$, seen as a subobject of $T \rightarrow \text{Type}_n^\circ$. $\circ_{n+1} T$ can be given explicitly by

$$\circ_{n+1} T \stackrel{\text{def}}{=} \sum_{u:T \rightarrow \text{Type}_n^\circ} \circ_{-1} \left\| \sum_{a:T} (\lambda t, \circ_n(a = t)) = u \right\|.$$

To prove that $\circ_{n+1}T$ is a sheaf for any $T : \text{Type}_{n+1}$, we use the following lemma.

Lemma 70

Any closed (-1) -subobject of a sheaf is a sheaf.

Proof. Let U be a sheaf, and $\kappa : U \rightarrow \text{HProp}$ be a closed (-1) -subobject. Let $E : \text{Type}$ and $\chi : E \rightarrow \text{HProp}$ dense in E . Let $\phi : \sum_{e:E} \chi e \rightarrow \sum_{u:U} \kappa u$. As $\pi_1 \circ \phi$ is a map $\sum_{e:E} \chi e \rightarrow U$ and U is a sheaf, it can be extended into $\psi : E \rightarrow U$. As κ is closed, it suffices now to prove $\prod_{e:E} \circ_n(\kappa(\psi e))$ to obtain a map $E \rightarrow \sum_{u:U} \kappa u$.

Let $e : E$. As χ is dense, we have a term $w : \circ_n(\chi e)$, and by \circ_n -induction, a term $\tilde{w} : \chi e$. Then, by retraction property, $\psi(e) = \phi(e, \tilde{w})$, and by $\pi_2 \circ \phi$, we have hence our term of type $\kappa(\psi e)$. □

As $T \rightarrow \text{Type}_n^\circ$ is a sheaf, and $\circ_{n+1}T$ is closed in $T \rightarrow \text{Type}_n^\circ$, $\circ_{n+1}T$ is a sheaf. We now prove that it forms a reflective subuniverse.

Proposition 71

(\circ_{n+1}, ν) defines a reflective subuniverse.

Proof. Let $T, Q : \text{Type}_{n+1}$ such that Q is a sheaf. Let $f : T \rightarrow Q$. Because Q is a sheaf, it is in particular separated; thus we can extend f to $\square_{n+1}f : \square_{n+1}T \rightarrow Q$.

But as $\circ_{n+1}T$ is the closure of $\square_{n+1}T$, $\square_{n+1}T$ is dense into $\circ_{n+1}T$, so the sheaf property of Q allows to extend $\square_{n+1}f$ to $\circ_{n+1}f : \circ_{n+1}T \rightarrow Q$.

As all these steps are universal, the composition is. □

The next step is the closure under dependent sums, to state:

Proposition 72

(\circ_{n+1}, ν) defines a modality.

Proof. The proof use the same ideas as in subsection 5.2.2. Let $A : \text{Type}_{n+1}$ a sheaf and $B : A \rightarrow \text{Type}_{n+1}$ a sheaf family. By proposition 69, we already know that $\sum_{a:A} Ba$ is separated. Let E be a type, and $\chi : E \rightarrow \text{HProp}$ a dense subobject. Let $f : \sum_{e:E} \chi e \rightarrow \sum_{x:A} Bx$; we want to extend it into a map $E \rightarrow \sum_{x:A} Bx$.

$$\begin{array}{ccc}
\sum_{e:E} \chi e & \xrightarrow{f} & \sum_{x:A} Bx \\
\downarrow & \nearrow \text{dotted} & \\
E & &
\end{array}$$

As A is a sheaf, and $\pi_1 \circ f : \sum_{e:E} \chi e \rightarrow A$, we can recover an map $g_1 : E \rightarrow A$. We then want to show $\prod_{e:E} B(g_1 e)$. Let $e : E$. As χ is dense, we have a term $w : \circ_n(\chi e)$, and as $B(g_1 e)$ is a sheaf, we can recover a term $\tilde{w} : \chi e$. Then $g_1(e) = f(e, \tilde{w})$, and $\pi_2 \circ f$ gives the result. \square

It remains to show that \circ_{n+1} is left exact and is compatible with \circ_{-1} . To do that, we need to extend the notion of compatibility and show that actually every modality \circ_{n+1} is compatible with \circ_n on lower homotopy types.

Proposition 73

If $T : \text{Type}_n$, then $\circ_{n+1} \widehat{T} = \circ_n T$, where \widehat{T} is T seen as a Type_{n+1} .

Proof. We prove it by induction on n :

- For $n = -1$: Let $T : \text{HProp}$. Then

$$\begin{aligned}
\circ_0 \widehat{T} &\stackrel{\text{def}}{=} \sum_{u:T \rightarrow \text{Type}_n^\circ} \circ_{-1} \left\| \sum_{a:T} (\lambda t, \circ_{-1}(a=t)) = u \right\|_{-1} \\
&= \sum_{u:T \rightarrow \text{Type}_n^\circ} \circ_{-1} \left(\sum_{a:T} (\lambda t, \circ_{-1}(a=t)) = u \right)
\end{aligned}$$

because the type inside the truncation is already in HProp . Now, let define $\phi : \circ_{-1} T \rightarrow \circ_0 T$ by

$$\phi t = (\lambda t', \mathbf{1}; \kappa)$$

where κ is defined by \circ_{-1} -induction on t . Indeed, as T is an HProp , $(a=t) \simeq \mathbf{1}$. Let $\psi : \circ_0 T \rightarrow \circ_{-1} T$ by obtaining the witness $a : T$ (which is possible because we are trying to inhabit a modal proposition), and letting $\psi(u; x) = \eta_T a$. These two maps form an equivalence (the section and retraction are trivial because the equivalence is between mere propositions).

- Suppose now that \circ_{n+1} is compatible with all \circ_k on lower homotopy types. Let \circ_{n+2} be as above, and let $T : \text{Type}_{n+1}$. Then, as \circ_{n+1} is compatible with \circ_n , and $(a=t)$ is in Type_n ,

$$\circ_{n+2} \widehat{T} = \sum_{u:T \rightarrow \text{Type}_{n+1}^\circ} \circ_{-1} \left\| \sum_{a:T} (\lambda t, \circ_n(a=t)) = u \right\|_{-1}.$$

It remains to prove that for every (u, x) inhabiting the Σ -type above, u is in $T \rightarrow \text{Type}_n^\circ$, i.e. that for every $t : T$, $\text{Is-}n\text{-type}(u\ t)$. But for any truncation index p , the type $\text{Is-}p\text{-type } X : \mathbf{HProp}$ is a sheaf as soon as X is, so we can get rid of \circ_{-1} and of the truncation, which tells us that for every $t : T$, $u\ t = \circ_n(a = t) : \text{Type}_n$. \square

This proves in particular that \circ_{n+1} is compatible with \circ_{-1} in the sense of condition 62.

The last step is the left-exactness of \circ_{n+1} . Let T be in Type_{n+1} such that $\circ_{n+1}T$ is contractible. Thanks to the just shown compatibility between \circ_{n+1} and \circ_n for Type_n , left exactness means that for any $x, y : T$, $\circ_n(x = y)$ is contractible.

Using a proof by univalence as we have done for proving $\circ_n(a = b) \simeq (\mu_T(a) = \mu_T(b))$ in Proposition 66, we can show that:

Proposition 74

For all $a, b : T$, $\circ_n(a = b) \simeq (v_T a = v_T b)$.

As $\circ_{n+1}T$ is contractible, path spaces of $\circ_{n+1}T$ are contractible, in particular $(v_T a = v_T b)$, which proves left exactness.

5.2.3 Summary

Starting from any left-exact modality \circ_{-1} on \mathbf{HProp} , we have defined for any truncation level n , a new left-exact modality \circ_n on Type_n , which corresponds to \circ_{-1} when restricted to \mathbf{HProp} .

When \circ_{-1} is consistent (in the sense of proposition 28), $\circ_n \mathbf{0} = \circ_{-1} \mathbf{0}$ is also not inhabited, hence the homotopy type theory induced by \circ_n is consistent. In particular, the modality induced by the double negation modality on \mathbf{HProp} is consistent.

In topos theory, the topos of Lawvere-Tierney sheaves for the double negation topology is a boolean topos. In homotopy type theory, this result can be expressed as:

Proposition 75

$(\circ_{\neg\neg})_n$, the modality obtained by sheafification of the double negation modality, induces a type theory where the propositional excluded middle law holds, i.e

$$\prod_{P:\mathbf{HProp}} P + \neg P.$$

Proof. For now, the only thing we know is that $\prod_{P:\mathbf{HProp}} \neg\neg P \rightarrow P$. Let $P : \mathbf{HProp}$, and pose $Q \stackrel{\text{def}}{=} P + \neg P$. Then, as P and $\neg P$ are disjoint h-propositions, $P + \neg P$ is itself a h-proposition [HoTT/Coq, ishprop_sum].

We want to find a proof of $Q : \mathbf{HProp}$, and thus we only need a proof of $\neg\neg Q$, which is inhabited by

$$\lambda (x : \neg Q), x(\text{inr}(\lambda y : P, x(\text{inl } y))).$$

□

5.2.4 Extension to Type

In the previous section, we have defined a (countably) infinite family of modalities $\text{Type}_i \rightarrow \text{Type}_i$. One can extend them to whole Type by composing with truncation:

Lemma 76

Let $\circ_i : \text{Type}_i \rightarrow \text{Type}_i$ be a modality. Then $\circ \stackrel{\text{def}}{=} \circ_i \circ \|\cdot\|_i : \text{Type} \rightarrow \text{Type}$ is a modality in the sense of section 3.1

If \circ_{-1} is the double negation modality on \mathbf{HProp} and $i = -1$, \circ is exactly the double negation modality on Type described in 3.2.3. Choosing $i \geq 0$ is a refinement of this double negation modality on Type: it will collapse every type to a Type_i , instead of an \mathbf{HProp} .

Obviously, as truncation modalities are not left-exact [HoTT, Exercise 7.11], \circ isn't either. But in the following sense, when restricted to i -truncated types, it is:

Lemma 77

Let $A : \text{Type}_i$. Then if $\circ(A)$ is contractible, for any $x, y : A$, $\circ(x = y)$ is contractible.

Proof. For i -truncated types, $\circ = \circ_i$, and \circ_i is left-exact. □

The compatibility between the modalities \circ_n and between the modalities $\|\cdot\|_n$ allow us to choose the truncation index as high as desired. Taking it as a non-fixed parameter allows to work in an universe where the new principle (e.g. mere excluded middle) is true for any explicit truncated type. Indeed, i can be chosen dynamically along a proof, and thus be increased as much as needed, without changing results for lower truncated types.

By proposition 28, these left-exact modalities induces a consistent type theory. Furthermore, the univalence remains true in this new type theory in the following sense:

Proposition 78

Let n be a given truncation index, and \circ the modality associated to n as de-

defined in lemma 76. Then, for any type $A, B : \text{Type}_n^\circ$, if φ is the canonical arrow

$$A = B \rightarrow A \simeq B,$$

then $\text{IsEquiv}(\varphi)$ is modal.

Proof. The first thing to notice is that, if X and Y are modal, and $f : X \rightarrow Y$, then the mere proposition $\text{IsEquiv } f$ is also modal. Therefore, it suffices to show that both $A = B$ and $A \simeq B$ are modal. By proposition 21, $A = B$ is modal. Moreover, $(A \simeq B) \simeq \sum_{f:A \rightarrow B} \text{IsEquiv } f$. Therefore, as A and B are modal, $A \simeq B$ is too.

Hence, $\text{IsEquiv } \varphi$ is modal. \square

We can view sheafification in terms of model of type theory but because of the resulting modality on Type is not left exact, we need to restrict ourselves to a type theory with only one universe. Let \mathfrak{M} be a model of homotopy type theory with one universe. Using the modality $\circlearrowleft_{\neg, \neg}$ (for any level n) associated to the sheafification, there is a model $\circlearrowleft_{\neg, \neg} \mathfrak{M}$ of type theory with one universe (using results in Section 3.3), where excluded middle is true, and which is univalent (as shown in Proposition 78).

5.3 Formalization

A Coq formalization of the sheafification process based on the Coq/HoTT library [HoTT/Coq] is available at <https://github.com/KevinQuirin/sheafification>.

After reviewing the content and some statistics about the formalization in Section 5.3.1, we present the limitations of our formalization in Section 5.3.2, in particular the issues relative to universe polymorphism.

5.3.1 Content of the formalization

We provide a more detailed insight of the structure of our formalization:

- Colimits and iterated kernel pairs are formalized in `Limit.v`, `OT.v`, `OT_Tf.v`, `T_telescope.v`, `Tf_0mono_sep.v`.
- Reflective subuniverses and modalities are formalized in `reflective_subuniverse.v`, `modalities.v`.
- The definition of the dense topology as a left exact modality on HProp is given in `sheaf_base_case.v`.
- Section 5.2.1 is formalized in `sheaf_def_and_thm.v`.
- Section 5.2.2 is formalized in `sheaf_induction.v`.

Overall, the project contains 7914 lines, and it could be reduced a bit by improving the way Coq tries to rewrite and apply lemmas automatically. The `coqwc` tool counts 1611 lines of specifications (definitions, lemmas, theorems, propositions) and 5403 lines of proof script. This constitutes a significant amount of work but the part dedicated to sheaves and sheafification is only 2200 lines of proof script, which seems quite reasonable and encouraging, because it suggests that homotopy type theory provides a convenient tool to formalize some part of the theory of higher topoi.

5.3.2 Limitations of the formalization

In the formalization, we had to use the `type-in-type` option, to handle the universe issues we faced.

Universes are used in type theory to ensure consistency by checking that definitions are well-stratified according to a certain hierarchy. Universe polymorphism [ST14] supports generic definitions over universes, reusable at different levels. Although the presence of universe polymorphism is mandatory for our formalization, its implementation is still too rigid to allow a complete formalization of our work for the following reasons.

If Coq handles cumulativity on `Type` natively, it is not the case for the Σ -type Type_n , which require propositional resizing. This issue could be solved by adding an axiom of cumulativity for Type_n with an explicit management of universes. But as it would not have any computational content, such a solution would really complicate the proofs as the axiom would appear everywhere cumulativity is needed and it would need explicit annotations for universe levels everywhere in the formalization.

One issue with universe polymorphism lies in the management of recursive definitions. Indeed, the following recursive definition of sheafification

$$\begin{aligned} \circ : \forall (n : \text{nat}), \text{Type}_n &\rightarrow \text{Type}_n \\ \circ_{-1} (T) &\stackrel{\text{def}}{=} \neg\neg T \\ \circ_{n+1} (T) &\stackrel{\text{def}}{=} \sum_{u : T \rightarrow \text{Type}_n^\circ} \circ_{-1} \left\| \sum_{a : T} u = (\lambda t, \circ_n(a = t)) \right\| \end{aligned}$$

is not allowed. This is because Coq forces the universe of the first Type_n occurring in the definition to be the same for every n , whereas the universe of the first Type_{n+1} occurring in \circ_{n+1} should be at least one level higher as the one of Type_n occurring in \circ_n because of the use of Σ -type over $T \rightarrow \text{Type}_n^\circ$ and equality on the return type of \circ_n . Thus, the induction step presented in this paper has been formalized, but the complete recursive sheafification can not be defined for the moment. Note that the same increasing in the universe levels occurs in the Rezk completion for categories [AKS15]. In the definition of the completion, they use the Yoneda embedding and representable functors, which is similar to our use of characteristic functions.

This restriction in our formalization may be solved by generalizing the management of universe polymorphism for recursive definition or by the use of general “resizing axiom” which is still under discussion in the community.

5.4 Forcing in type theory

Try to write something smart about connections with forcing

Conclusion and future works

I clearly need to write a conclusion here

Write a conclusion

What remains to be done

Write a better abstract	i
Write the acknowledgments	iii
Go on!	1
Check plugin	2
These references were added without even opening them. Maybe a check would be good	15
Check post or pre	18
New type theory: Enhance me	27
Code the plugin, then plug it here	30
Formalization of colimits: what could we say ?...	41
Define groupoid objects in HoTT, then write it here	42
Post or Pre?	50
Try to write something smart about connections with forcing	61
Write a conclusion	63

Nomenclature

$(x : A) \rightarrow (B a)$	Dependent product over B , page 5
$p \cdot q$	Concatenation of paths, page 8
$\text{fib}_f(b)$	Fiber of f over b , page 6
idpath or 1	Constant path, page 7
idpath_x or 1_x	Constant path over x , page 7
p^{-1}	Inverse of path, page 8
\mathbb{N}	Type of naturals, page 6
$\mathbf{1}$	Unit type, page 4
$\prod_{a:A} B a$	Dependent product over B , page 5
ΣA	Suspension of a type, page 14
\simeq	Equivalence of types, page 10
$\sum_{a:A} B a$	Dependent sum over B , page 6
$\ \cdot\ $	n -truncation of types, page 18
$ \cdot _n$	n -truncation of terms or arrows, page 18
$\mathbf{0}$	Empty type, page 4
$A + B$	Coproduct of types, page 5
$a : A$	Judgement “ a is of type A ”, page 3
$a =_A b$	Type of paths from a to b in A , page 7
$a = b$	Type of paths from a to b , page 7
$A \times B$	Product of types, page 5
$A \rightarrow B$	Type of arrows from A to B , page 5

Bibliography

- [AGS12] Steve Awodey, Nicola Gambino, and Kristina Sojakova. “Inductive types in homotopy type theory”. In: *Proceedings of the 2012 27th Annual IEEE/ACM Symposium on Logic in Computer Science*. IEEE Computer Society. 2012, pp. 95–104 (cit. on p. 6).
- [AKL15] Jeremy Avigad, Krzysztof Kapulkin, and Peter LeFanu Lumsdaine. “Homotopy limits in type theory”. In: *Mathematical Structures in Computer Science* (Published online: 19 january 2015) (cit. on p. 31).
- [AKS15] Benedikt Ahrens, Krzysztof Kapulkin, and Michael Shulman. “Univalent Categories and the Rezk completion”. In: *Mathematical Structures in Computer Science* (Published online: 19 january 2015) (cit. on p. 60).
- [BL11] Andrej Bauer and Peter LeFanu Lumsdaine. *A Coq proof that Univalence Axioms implies Functional Extensionality*. 2011. URL: <https://github.com/andrejbauer/Homotopy/raw/master/OberwolfachTutorial/univalence.pdf> (cit. on p. 11).
- [Bou16] Simon Boulier. *Colimits in HoTT*. Blog post. 2016. URL: <http://homotopytypetheory.org/2016/01/08/colimits-in-hott/> (cit. on pp. 31, 38).
- [CD66] P.J. Cohen and M. Davis. *Set theory and the continuum hypothesis*. WA Benjamin New York, 1966 (cit. on p. 44).
- [Göd40] Kurt Gödel. *The Consistency of the Axiom of Choice and of the Generalized Continuum Hypothesis with the Axioms of Set Theory*. Princeton University Press, 1940 (cit. on p. 44).
- [Hof97] Martin Hofmann. “Syntax and semantics of dependent types”. In: *Extensional Constructs in Intensional Type Theory*. Springer, 1997, pp. 13–54 (cit. on p. 3).
- [HoTT] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. IAS: <http://homotopytypetheory.org/book>, 2013 (cit. on pp. 2–4, 7, 9, 14, 17, 24, 36, 45, 52, 58).
- [HoTT/Coq] The Univalent Foundations Program. *Coq HoTT library*. URL: <https://github.com/HoTT/HoTT/> (cit. on pp. 2, 13, 27, 28, 57, 59).

- [How80] William A. Howard. “The formulas-as-types notion of construction”. In: *To H. B. Curry: Essays on Combinatory Logic, Lambda Calculus, and Formalism*. Ed. by J. P. Seldin and J. R. Hindley. Reprint of 1969 article. Academic Press, 1980, pp. 479–490 (cit. on p. 4).
- [HS96] Martin Hofmann and Thomas Streicher. “The Groupoid Interpretation of Type Theory”. In: *In Venice Festschrift*. Oxford University Press, 1996, pp. 83–111 (cit. on p. 10).
- [HTS] Vladimir Voevodsky. *A simple type system with two identity types*. Started in 2013, in progress (cit. on p. 8).
- [HTT] Jacob Lurie. *Higher topos theory*. Annals of mathematics studies. Princeton, N.J., Oxford: Princeton University Press, 2009 (cit. on pp. 2, 41).
- [Hut11] Michael Hutchings. *Introduction to higher homotopy groups and obstruction theory*. 2011. URL: <https://math.berkeley.edu/~hutching/> (cit. on p. 15).
- [JTS12] Guilhem Jaber, Nicolas Tabareau, and Matthieu Sozeau. “Extending type theory with forcing”. In: *Logic in Computer Science (LICS), 2012 27th Annual IEEE Symposium on*. IEEE. 2012, pp. 395–404 (cit. on pp. 28, 30).
- [Kra15] Nicolai Kraus. “Truncation Levels in Homotopy Type Theory”. PhD thesis. University of Nottingham, June 2015 (cit. on p. 18).
- [Kun] Kenneth Kunen. *Set theory*. Ed. by North-Holland (cit. on p. 21).
- [Lic] Dan Licata. *Another proof that univalence implies function extensionality*. Blog post. URL: <http://homotopytypetheory.org/2014/02/17/another-proof-that-univalence-implies-function-extensionality/> (cit. on p. 11).
- [LS13] Daniel R. Licata and Michael Shulman. “Calculating the Fundamental Group of the Circle in Homotopy Type Theory”. In: *LICS 2013: Proceedings of the Twenty-Eighth Annual ACM/IEEE Symposium on Logic in Computer Science*. 2013 (cit. on p. 12).
- [m31] Andrej Bauer. *Andromeda implementation* (cit. on p. 8).
- [Mar98] Per Martin-Löf. “An intuitionistic theory of types”. In: *Twenty-five years of constructive type theory* 36 (1998), pp. 127–172 (cit. on p. 4).
- [MM92] Saunders MacLane and Ieke Moerdijk. *Sheaves in Geometry and Logic*. Springer-Verlag, 1992 (cit. on pp. 37, 43–45, 47).
- [Mou+15] Leonardo Mendonça de Moura et al. “The Lean Theorem Prover (System Description)”. In: *Automated Deduction - CADE-25 - 25th International Conference on Automated Deduction, Berlin, Germany, August 1-7, 2015, Proceedings*. 2015, pp. 378–388 (cit. on p. 1).

- [MV98] Fabien Morel and Vladimir Voevodsky. “ A^1 -homotopy theory of schemes”. In: (1998) (cit. on p. 45).
- [Nor07] Ulf Norell. “Towards a practical programming language based on dependent type theory”. PhD thesis. Chalmers University of Technology, 2007 (cit. on p. 1).
- [NPS01] B Nordström, K Petersson, and JM Smith. *Martin-Löf’s type theory, Handbook of logic in computer science: Volume 5: Logic and algebraic methods*. Oxford University Press, Oxford, 2001 (cit. on p. 4).
- [RS13] Egbert Rijke and Bas Spitters. “Sets in homotopy type theory”. In: *arXiv preprint arXiv:1305.3835* (2013) (cit. on p. 17).
- [Shu15] Mike Shulman. *Module for Modalities*. Blog post. 2015. URL: <http://homotopytypetheory.org/2015/07/05/modules-for-modalities/> (cit. on p. 28).
- [ST14] Matthieu Sozeau and Nicolas Tabareau. “Universe Polymorphism in Coq”. In: *Interactive Theorem Proving*. 2014 (cit. on p. 60).
- [Str93] Thomas Streicher. “Investigations Into Intensional Type Theory”. Habilitationsschrift. LMU München, 1993 (cit. on p. 10).
- [The12] The Coq development team. *Coq 8.4 Reference Manual*. Inria. 2012. URL: <http://coq.inria.fr/distrib/V8.4p13/refman/> (cit. on p. 1).
- [Van16] Floris Van Doorn. “Constructing the Proposition Truncation using Non-recursive HITs”. In: *Certified Proofs and Programs* (2016) (cit. on p. 37).
- [Voe11] Vladimir Voevodsky. *Resising Rules - their use and semantic justification*. 2011 (cit. on p. 28).
- [Voe14] Vladimir Voevodsky. *Univalent Foundations*. 2014 (cit. on p. 1).
- [WX10] Guozhen Wang and Zhouli Xu. “A Survey of Computations of Homotopy Groups of Spheres and Cobordisms”. In: (2010) (cit. on p. 15).