

STAT 305 Assignment 3

Name: Kevin Nguyen
Student Number: 97703045

Question 1

1. We have $Y_i \sim Pois(\mu)$, therefore we have

$$\begin{aligned} f_{Y_i|M}(y_i \mid \mu) &= \frac{e^{-\mu} \mu^{y_i}}{y_i!} \\ \therefore f_{Y_1, \dots, Y_n|M}(y_1, \dots, y_n \mid \mu) &= \prod_{i=1}^n \frac{e^{-\mu} \mu^{y_i}}{y_i!} \\ &= \frac{e^{-n\mu} \mu^{\sum_{i=1}^n y_i}}{\prod_{i=1}^n y_i!} \end{aligned}$$

We also have the μ takes on the prior distribution of an exponential with prior mean 10 (so $\lambda = 1/10$), therefore

$$\begin{aligned} p_M(\mu) &= \frac{1}{10} e^{-\frac{1}{10} \cdot \mu} \\ \therefore p_{M|Y_1, \dots, Y_n}(\mu \mid y_1, \dots, y_n) &= \frac{\frac{1}{10} e^{-\frac{1}{10} \cdot \mu} \cdot \frac{e^{-n\mu} \mu^{\sum_{i=1}^n y_i}}{\prod_{i=1}^n y_i!}}{f_{Y_1, \dots, Y_n}(y_1, \dots, y_n)} \\ &\propto \frac{1}{10} e^{-\frac{1}{10} \cdot \mu} \cdot \frac{e^{-n\mu} \mu^{\sum_{i=1}^n y_i}}{\prod_{i=1}^n y_i!} \\ &\propto e^{-\frac{1}{10} \cdot \mu} \cdot e^{-n\mu} \mu^{\sum_{i=1}^n y_i} \\ &= e^{-(\frac{1}{10} + n)\mu} \cdot \mu^{\sum_{i=1}^n y_i} \\ &\sim Gamma\left(\sum_{i=1}^n y_i + 1, \frac{1}{10} + n\right) \end{aligned}$$

2. Following the formula given, we have

$$\begin{aligned}
P(Y_{1897} = 0 \mid Y) &= E[P(Y_{1987} = 0 \mid \mu) \mid Y] \\
&= E\left[\frac{e^{-\mu}\mu^0}{0!} \mid Y\right] \\
&= E[e^{-\mu} \mid Y] \\
&= \int_{-\infty}^{\infty} e^{-\mu} \cdot p_{M|Y}(\mu \mid y) d\mu \\
&= \int e^{-\mu} \cdot \frac{\beta^\alpha}{T(\alpha)} \mu^{\alpha-1} e^{-\beta\mu}
\end{aligned}$$

$$\text{where } \alpha = \sum_{i=1}^n y_i + 1 \text{ and } \beta = \frac{1}{10} + n$$

notice this is $E(e^{tX})$ or MGF of a Gamma when $t = -1$

$$\begin{aligned}
&= \left(\frac{\beta}{\beta - (-1)}\right)^\alpha \\
&= \left(\frac{\frac{1}{10} + 16}{\frac{1}{10} + 16 + 1}\right)^{57+1} \\
&= \left(\frac{16.1}{17.1}\right)^{58} \\
&= 0.03034774367 \\
&\approx 0.03
\end{aligned}$$

Question 2

1. We have the posterior density

$$\begin{aligned}
 f_{\Pi|Y}(\pi | y) &= \frac{f_{\Pi}(\pi) \cdot f_{Y|\Pi}(y | \pi)}{f_Y(y)} \\
 &= \frac{\left[\sum_{i=1}^3 \frac{1}{3} \text{Beta}(\pi, \alpha_i, \beta_i) \right] \cdot \binom{8197}{y} \pi^y (1 - \pi)^{8197-y}}{\int \sum_{i=1}^3 \frac{1}{3} \text{Beta}(\pi, \alpha_i, \beta_i) \cdot \binom{8197}{y} \pi^y (1 - \pi)^{8197-y}} \\
 &= \frac{\left[\sum_{i=1}^3 \frac{1}{3} \text{Beta}(\pi, \alpha_i, \beta_i) \cdot \pi^y (1 - \pi)^{8197-y} \right] \cdot \binom{8197}{y}}{\sum_{i=1}^3 \int \frac{1}{3} \text{Beta}(\pi, \alpha_i, \beta_i) \cdot \binom{8197}{y} \pi^y (1 - \pi)^{8197-y}} \\
 &= \frac{\left[\sum_{i=1}^3 \frac{1}{3} \frac{1}{B(\alpha_i, \beta_i)} \pi^{\alpha_i+y-1} (1 - \pi)^{\beta_i+8197-y-1} \right] \cdot \binom{8197}{y}}{\sum_{i=1}^3 \int \frac{1}{3} \frac{1}{B(\alpha_i, \beta_i)} \pi^{\alpha_i+y-1} (1 - \pi)^{\beta_i+8197-y-1} \cdot \binom{8197}{y}} \\
 &= \frac{\sum_{i=1}^3 \left[\frac{1}{3} \text{Beta}(\pi, \alpha_i + y, \beta_i + 8197 - y) \right] \cdot \binom{8197}{y}}{\frac{1}{3} \frac{1}{B(\alpha_i, \beta_i)} \binom{8197}{y} \sum_{i=1}^3 \int \pi^{\alpha_i+y-1} (1 - \pi)^{\beta_i+8197-y-1}} \\
 &= \frac{\sum_{i=1}^3 \text{Beta}(\pi, \alpha_i + y, \beta_i + 8197 - y)}{\sum_{i=1}^3 \frac{1}{B(\alpha_i, \beta_i)} B(\alpha_i + y, \beta_i + 8197 - y)} \\
 &= \sum_{i=1}^3 \pi'_i \cdot \text{Beta}(\pi, \alpha'_i, \beta'_i)
 \end{aligned}$$

where $\pi'_i = \sum_{i=1}^3 \frac{1}{B(\alpha_i, \beta_i)} B(\alpha_i + y, \beta_i + 8197 - y)$,

$\alpha'_i = \alpha_i + y$ and

$\beta'_i = \beta_i + 8197 - y \quad \square$

2. Using the posterior from the previous question,

$$\begin{aligned}
f_{\Pi|Y}(\pi \mid y = 51) &= \frac{\sum_{i=1}^3 \text{Beta}(\pi, \alpha_i + 51, \beta_i + 8146)}{\sum_{i=1}^3 \frac{1}{B(\alpha_i, \beta_i)} B(a_i + 51, \beta_i + 8146)} \\
&= \frac{\text{Beta}(\pi, 52, 8156) + \text{Beta}(\pi, 52, 8147) + \text{Beta}(\pi, 61, 8147)}{\frac{B(52, 8156)}{B(1, 10)} + \frac{B(52, 8147)}{B(1, 1)} + \frac{B(61, 8147)}{B(10, 1)}} \\
&= \frac{\text{Beta}(\pi, 52, 8156) + \text{Beta}(\pi, 52, 8147) + \text{Beta}(\pi, 61, 8147)}{3 \left[\frac{1}{3} \frac{B(52, 8156)}{B(1, 10)} + \frac{1}{3} \frac{B(52, 8147)}{B(1, 1)} + \frac{1}{3} \frac{B(61, 8147)}{B(10, 1)} \right]} \\
&= \frac{\text{Beta}(\pi, 52, 8156) + \text{Beta}(\pi, 52, 8147) + \text{Beta}(\pi, 61, 8147)}{3 \cdot Z} \\
&= \frac{\sum_{i=1}^3 \text{Beta}(\pi, \alpha_i + 51, \beta_i + 8146)}{3Z}
\end{aligned}$$

Question 3

1. Using the law of total probability, we have

$$\begin{aligned}
P(L(Y) \leq \Pi \leq R(Y)) &= E[P(L(Y) \leq \Pi \leq R(y) \mid Y)] \\
&= E[0.95] && \text{by definition of Credible Interval} \\
&= 0.95
\end{aligned}$$

2. Since π is randomly generated from the prior and y is also randomly generated based on π so both Π and Y are random variables, and thus, as showed above, the probability of the random variable Π falling within the left and right bound ($L(Y)$ and $R(Y)$, respectively) is 95%. Therefore, it's calibrated