

Estimates of Wenatchee Steelhead Redds and Spawners

Spawn Year 2022

Kevin See^{1,*}

January 17, 2023

Abstract

This report contains estimates of total steelhead redds in the Wenatchee, after accounting for observer error. It also includes estimates of spawners, as well as prespawn mortality.

Contents

1	Introduction	2
2	Methods	2
2.1	Net Error Model	2
2.2	Data	3
2.3	Estimating Redds	3
2.4	Estimating Spawners	4
2.5	Prespawn Mortality	4
3	Results	6
3.1	Redd estimates	6
3.2	Spawner Estimates	6
3.3	Prespawn Mortality	10
4	Discussion	10
5	Acknowledgements	10
6	References	11
A	Sex Errors at Priest Rapids	12

¹ Washington Department of Fish & Wildlife

* Correspondence: Kevin See <Kevin.See@dfw.wa.gov>

1 Introduction

Redd counts are an established method to provide an index of adult spawners (Gallagher et al. 2007). In the Wenatchee subbasin, index reaches are surveyed weekly during the steelhead spawning season (Mar 08, 2022 - Jun 01, 2022) and non-index reaches are surveyed once during the peak spawning period. The goal of this work is to:

1. Predict observer net error, using the model described in Murdoch et al. (2018).
2. Use estimates of observer net error rates and the mean survey interval to estimate the number of redds in each index reach, using a Gaussian area under the curve (GAUC) technique described in Millar et al. (2012) and Murdoch et al. (2018).
3. Estimate the total number of redds in the non-index reaches by adjusting the observed counts with the estimated net error where possible.
4. Sum the total number of estimated redds for the entire Wenatchee subbasin.
5. Translate the estimated number of redds into estimates of spawners, by origin.

2 Methods

2.1 Net Error Model

The net error (NE) for a reach i is defined as

$$NE_i = \frac{F_i}{V_i}$$

where F_i is the number of redds the surveyor reported and V_i is the true number of redds in the reach. Therefore, if we have an estimate of net error (\hat{NE}_i), we can calculate the true number of redds based on that estimate and the number of redds the surveyor reported:

$$V_i = \frac{F_i}{\hat{NE}_i} \quad (1)$$

The model for observer net error is fully described in Murdoch et al. (2018). It uses covariates of the log of observer experience, mean discharge, the observed redd density and mean thalweg CV as a proxy for channel complexity. After normalizing these covariates, the model coefficients are shown in Table 1. The response, net error, is scaled such that estimates of net error less than 1 suggest more errors of omission, while estimates greater than 1 suggest more errors of commission. An estimate of net error equal to one would indicate the observed count equals the true number of redds.

Table 1: Net error model covariates and coefficients.

Covariate	Estimate	Std. Error
(Intercept)	0.682	0.039
Obs. Redd Density	0.277	0.053
Mean Thalweg CV	-0.169	0.043
Mean Discharge	0.116	0.048
Log Surveyor Exp.	0.115	0.042

2.2 Data

Redd counts were conducted on a nearly weekly basis for index reaches, and once during the peak spawning period for non-index reaches. The covariates in the observer error model of Murdoch et al. (2018) were collected during each survey. The covariate of mean thalweg CV was calculated based on all measurements taken within a reach across years (assuming this covariate does not vary through time within a reach). They were compared with the covariates contained in the model data set, as well as the estimates of net error (Figure 1).




Figure 1: Net error covariate values. Colors correspond to either the original study or the reaches where the model was used in this report.

Those covariates in the observer error model were collected during each survey in 2022, and predictions of net error were made for each survey. From these survey specific estimates of net error, a mean and standard error of net error was calculated for each reach. The standard deviation was calculated by taking the square root of the sum of the squared standard errors for all predictions within a reach.

2.3 Estimating Redds

Use of the GAUC methodology was limited to index reaches with a minimum of two reds and at least three weeks with at least one new redd found. For those reaches, we used the method described in Millar et al. (2012) and Murdoch et al. (2018). The GAUC model was developed with spawner counts in mind. As it is

usually infeasible to mark every individual spawner, only total spawner counts can be used, and an estimate of average stream life must be utilized to translate total spawner days to total unique spawners. However, in adapting this for redd surveys, we note that individual redds can be marked, and therefore the GAUC model can be fit to new redds only. The equivalent of stream life is thus the difference between survey numbers, which can be fixed at 1. We applied the average net error from all the surveys when redds were visible, so as to not bias the estimate from early weeks when no redds were found, and the observed redd density was zero.

For non-index reaches, which were surveyed only once during peak spawning, the estimate of total redds was calculated by dividing the observed redds by the estimate of net error associated with that survey (Eq. (1)). This assumes that no redds were washed out before the non-index survey, and that no new redds appeared after that survey. As the number of redds observed in the non-index reaches ranged from 0 to 1, any violation of this assumption should not affect the overall estimates very much. Any index reaches that did not meet the thresholds described above were treated as non-index reaches, and the total observed redds in those reaches were divided by the mean estimate of net error for each reach.

When summing reach-scale estimates to obtain estimates at the stream scale, an attempt was made to incorporate the fact that the reaches within a stream are not independent. Estimates of correlation between the reaches within a stream were made based on weekly observed redds. This method may not be perfect, since spawners could use certain reaches preferentially at different times in the season, but it may be the best we can do. Because correlations are often quite high between reaches, this is a better alternative than to naively assume the standard errors between reaches are independent of one another. These correlation estimates were combined with estimates of standard error at the reach scale to calculate a covariance matrix for the reaches within each stream, which was used when summing estimates of total redds to estimate the standard error at the stream scale. Failure to incorporate the correlations between reaches would result in an underestimate of standard error at the stream scales. Different streams (and therefore reaches in different streams) were assumed to be independent.

2.4 Estimating Spawners

Estimates of escapement to various tributaries in the Wenatchee were made using a Dam Adult Branch Occupancy Model (DABOM) (Waterhouse et al. 2020) based on PIT tag detections of fish tagged at Priest Rapids dam. All fish that escaped to the various tributaries were assumed to be spawners (i.e. prespaw mortality only occurs in the mainstem). Reaches below the PIT tag arrays in some tributaries were surveyed for redds, but we assumed there was no observer error in those reaches.

To convert estimates of redds in mainstem areas into estimates of natural and hatchery spawners, the estimates of redds were multiplied by a fish per redd (FpR) estimate and then by the proportion of hatchery or wild fish. The fish per redd estimate was based on PIT tags from the branching patch-occupancy model observed to move into the lower or upper Wenatchee (below or above Tumwater dam), but not into tributaries upstream of Tumwater. FpR was calculated as the ratio of male to female fish, plus 1. Reaches W1 - W7 are below Tumwater, while reaches W8 - W10 are above Tumwater. Similarly, the proportion of hatchery and natural origin fish was calculated from the same group of PIT tags for areas above and below Tumwater. For reaches in tributaries below the tributary PIT tag array, FpR and proportion hatchery were calculated based on observed PIT tags moving into the lower regions of each tributary.

For 2022, we noticed a discrepancy in the sex calls at Priest Rapids and the sex calls from broodstock collection at Wells Hatchery and in the Wenatchee. Therefore, we developed a correction factor that accounts for the error rate in female and male calls independently, and adjusts the fish / redd estimate accordingly. The error rate of sex calls at Priest are shown in Appendix A in Table 10.

2.5 Prespawn Mortality

After translating estimates of redds to estimates of spawners by origin, we can then compare the spawner estimates to escapement estimates made using PIT tags, and estimate a prespaw mortality rate. Taking the

Table 2: Known number of fish removed at dams or due to harvest, by origin.

Source	Hatchery	Natural
Dryden	16	0
Tumwater	32	51

total PIT-tag based escapement estimate, by origin, to the Wenatchee, we then subtract any fish removed at Tumwater or Dryden for broodstock or surplus, as well as any deaths due to harvest (Table 2), and then subtract the total estimate of spawners, including the tributaries, to provide an estimate of how many fish succumbed to prespawn mortality. Dividing that number by the total escapement estimate provides an estimate of the prespawn mortality rate, by origin, across the entire Wenatchee population.

If either origin had a higher estimates of spawners compared to escapement, we fixed our prespawn mortality estimate at 0, reflecting a very low level of prespawn mortality. There is uncertainty in both the escapement and spawner estimates, which could explain why these types of scenarios could arise.

Table 3: Estimates of mean net error and redds for each reach.

River	Reach	Type	Net Error	Net Error CV	Observed Redds	Estimated Redds	Std. Err. Redds	Redds CV
Chiwawa	C1	Tributary	1	0	0	0	0	-
Nason	N1	Tributary	1	0	0	0	0	-
Peshastin	P1	Tributary	1	0	0	0	0	-
Wenatchee	W1	Non-Index	1	0	0	0	0	-
Wenatchee	W10	Non-Index	1	0	1	1	0	0
Wenatchee	W10	Index	0.674	0.161	4	7	5.5	0.786
Wenatchee	W2	Index	1	0	0	0	0	-
Wenatchee	W3	Non-Index	1	0	0	0	0	-
Wenatchee	W4	Non-Index	1	0	0	0	0	-
Wenatchee	W5	Non-Index	1	0	0	0	0	-
Wenatchee	W6	Index	1.238	0.19	4	3	1.5	0.506
Wenatchee	W8	Non-Index	1	0	0	0	0	-
Wenatchee	W8	Index	0.588	0.177	2	3	0.6	0.201
Wenatchee	W9	Non-Index	1	0	0	0	0	-
Wenatchee	W9	Index	0.668	0.183	4	6	1.2	0.197

Table 4: Estimate of redds in mainstem Wenatchee, above and below Tumwater Dam, and any tributaries surveyed.

River	Location	Index	# Reaches	Observed Redds	Estimated Redds	Std. Err. Redds	Redds CV
Wenatchee	Mainstem Below Tumwater	N	4	0	0	0	-
Wenatchee	Mainstem Below Tumwater	Y	2	4	3	1.5	0.506
Wenatchee	Mainstem Above Tumwater	N	3	1	1	0	0
Wenatchee	Mainstem Above Tumwater	Y	3	10	16	5.7	0.354
Chiwawa	Tributary	Tributary	1	0	0	0	-
Nason	Tributary	Tributary	1	0	0	0	-
Peshastin	Tributary	Tributary	1	0	0	0	-
Total	-	-	15	15	20	5.9	0.293

3 Results

3.1 Redd estimates

The estimated net error, observed redds and estimates of redds at the reach scale are shown in Table 3. Plots of the new redds and the GAUC fit to those data are shown in Figure 2. The results are summarized by major areas and population scale in Table 4.

3.2 Spawner Estimates

Demographic data about sex and origin were derived from the PIT tags detected within each area (Table 5. Parameter estimates for fish / redd and proportion hatchery based on this PIT tag data are shown in Table 6.

Combining PIT tag-based estimates of spawners in the tributaries with adjusted redd-based estimates of spawners in the mainstem areas, Table 7 shows all of them, broken down by reach or tributary and origin. Table 8 shows a summary of spawners by origin across various areas including the Wenatchee mainstem and each tributary.




Figure 2: Plots of observed redd counts (black dots) through time for each qualifying index reach, and the fitted curve from the GAUC model (blue line) with associated uncertainty (gray).

Table 5: Number of PIT tags detected in each area by sex and origin.

Location	Sex	Origin	n tags
Below TUM (mainstem)	F	H	2
Below TUM (mainstem)	F	W	3
Below TUM (mainstem)	M	H	1
Above TUM (mainstem)	F	H	4
Above TUM (mainstem)	F	W	8
Above TUM (mainstem)	M	H	3
Above TUM (mainstem)	M	W	5
Tributaries	F	H	3
Tributaries	F	W	22
Tributaries	M	H	2
Tributaries	M	W	6
Peshastin	F	W	7
Peshastin	M	H	1
Nason	F	W	4
Nason	M	W	1
Chiwawa	F	W	1
Chiwawa	M	W	1

Table 6: Fish per redd and hatchery / natural origin proportion estimates.

Area	Fish / redd	FpR Std. Error	Prop. Hatchery	Prop Std. Error
Below TUM (mainstem)	3.000	0.431	0.500	0.204
Above TUM (mainstem)	3.333	0.391	0.350	0.107
Tributaries	2.750	0.324	0.152	0.062
Peshastin	2.667	0.350	0.125	0.117
Nason	2.500	0.278	0.000	0.000
Chiwawa	2.000	1.414	0.000	0.000

Table 7: Estimates (CV) of spawners by reach or tributary and origin.

River	Area	Type	Natural	Hatchery
Wenatchee	W1	Non-Index	0 (-)	0 (-)
Wenatchee	W2	Index	0 (-)	0 (-)
Wenatchee	W3	Non-Index	0 (-)	0 (-)
Wenatchee	W4	Non-Index	0 (-)	0 (-)
Wenatchee	W5	Non-Index	0 (-)	0 (-)
Wenatchee	W6	Index	5 (0.67)	5 (0.67)
Wenatchee	W8	Index	7 (0.28)	4 (0.38)
Wenatchee	W8	Non-Index	0 (-)	0 (-)
Wenatchee	W9	Index	13 (0.28)	7 (0.38)
Wenatchee	W9	Non-Index	0 (-)	0 (-)
Wenatchee	W10	Index	15 (0.81)	8 (0.85)
Wenatchee	W10	Non-Index	2 (0.2)	1 (0.33)
Icicle	Icicle	DABOM	79 (0.35)	19 (0.67)
Peshastin	P1	Tributary	0 (-)	0 (-)
Peshastin	Peshastin	DABOM	70 (0.37)	14 (0.89)
Mission	Mission	DABOM	15 (0.84)	13 (0.87)
Chumstick	Chumstick	DABOM	22 (0.65)	0 (-)
Chiwaukum	Chiwaukum	DABOM	21 (0.69)	10 (0.94)
Chiwawa	C1	Tributary	0 (-)	0 (-)
Chiwawa	Chiwawa	DABOM	25 (0.78)	0 (-)
Nason	N1	Tributary	0 (-)	0 (-)
Nason	Nason	DABOM	41 (0.47)	0 (-)
Little Wenatchee	Little Wenatchee	DABOM	0 (-)	0 (-)
White River	White River	DABOM	0 (-)	0 (-)
Total	Total	-	314 (0.17)	81 (0.31)

Table 8: Estimates (CV) of spawners by area and origin.

River	Location	Natural	Hatchery
Wenatchee	Below TUM (mainstem)	5 (0.6)	5 (0.6)
Wenatchee	Above TUM (mainstem)	37 (0.35)	20 (0.38)
Icicle	Icicle	79 (0.35)	19 (0.68)
Peshastin	Peshastin	70 (0.37)	14 (0.88)
Mission	Mission	15 (0.82)	13 (0.88)
Chumstick	Chumstick	22 (0.66)	0 (-)
Chiwaukum	Chiwaukum	21 (0.7)	10 (0.98)
Chiwawa	Chiwawa	25 (0.78)	0 (-)
Nason	Nason	41 (0.46)	0 (-)
Little Wenatchee	Little Wenatchee	0 (-)	0 (-)
White River	White River	0 (-)	0 (-)
Total	-	314 (0.17)	81 (0.31)

Table 9: Wenatchee prespawn mortality estimates. Includes estimates (standard error) of escapement, spawners, rate of prespawn mortality, and standard error of this rate, separated by origin.

Origin	Escapement	Spawners	Prespawn Mortality	SE	CV
Natural	355 (60)	315 (54)	0.113	0.214	1.896
Hatchery	95 (36)	81 (25)	0.145	0.419	2.892

3.3 Prespawn Mortality

The estimates of overall prespawn mortality within the Wenatchee population, by origin, are shown in Table 9.

4 Discussion

Most of the covariates collected in 2022 were within the range of those in the model data set from Murdoch et al. (2018), leading to estimates of net error in a very similar range to the model dataset (Figure 1). However, some reaches did not meet the minimum thresholds of number of observed redds or number of weeks with at least one new redd observed, so we used the GAUC method in only three reaches.

Hatchery origin steelhead PIT-tagged at Priest Rapids dam were not detected in a number of tributaries this year, leading to estimates of zero hatchery spawners in those tributaries (Table 8). Because these estimates are based on a sample of the entire Upper Columbia steelhead run, it's possible a handful of steelhead may have spawned in those tributaries, but it's potentially more likely that the estimate of zero steelhead spawners in those areas is correct.

5 Acknowledgements

The data for this report was collected by Washington Department of Fish and Wildlife, and funded by Chelan County Public Utility District.

6 References

- Gallagher, S. P., P. K. J. Hahn, and D. H. Johnson. 2007. Salmonid field protocols handbook: Techniques for assessing status and trends in salmon and trout populations. Pages 197–234 *in* D. H. Johnson, editor. American Fisheries Society, Bethesda, Maryland.
- Millar, R. B., S. McKechnie, and C. E. Jordan. 2012. Simple estimators of salmonid escapement and its variance using a new area-under-the-curve method. *Canadian Journal of Fisheries and Aquatic Sciences* 69(6):1002–1015.
- Murdoch, A. R., C. J. Herring, C. H. Frady, K. See, and C. E. Jordan. 2018. Estimating observer error and steelhead redd abundance using a modified gaussian area-under-the-curve framework. *Canadian Journal of Fisheries and Aquatic Sciences* (999):1–10.
- Waterhouse, L., J. White, K. See, A. Murdoch, and B. X. Semmens. 2020. A bayesian nested patch occupancy model to estimate steelhead movement and abundance. *Ecological Applications*.

Table 10: Error rate in sex calls at Priest Rapids.

Brood Year	Sex	N Tags	N True	N False	Perc False	Lwr CI	Upr CI
2022	Female	42	20	22	0.524	0.377	0.666
2022	Male	18	18	0	0.000	0.000	0.176

A Sex Errors at Priest Rapids

Fish collected for broodstock at Wells Hatchery and within the Wenatchee had a known sex. Based on PIT tag detections of this broodstock collection, those sexes could be compared with the sex call at Priest when the fish was initially tagged. Assuming that this error rate is systematic across the entire run within a brood year, we combined all tags with this kind of comparison. Table 10 shows the sample sizes and results.

These error rates were used to adjust the counts of male and female tags within particular spawning areas before re-calculating the fish/redd estimate. For example, the adjusted number of males would be the initial number of males, minus the number of initial males times the error rate for male identification, plus the number of initial females times the error rate for female identification.