

*Homework Notes:* Add information here on your study group, number of hours you spent on the homework, and other relevant information.

### Problem 1

You can write aligned equations as follows:

$$\begin{aligned} a &\sim p(a) \\ b &\sim p(b). \end{aligned}$$

**(7 points)** Pairwise independence does not imply mutual independence. Two random variables,  $X_i$ ,  $i=1,2$  are independent if  $P(X_i | X_j) = P(X_i)$ , for  $i,j=1,2$ ,  $i \neq j$  and therefore

$$\begin{aligned} P(X_i, X_j) &= P(X_j)P(X_i) \\ P(X_i | X_j) &= P(X_i)P(X_j) \end{aligned}$$

Now, given  $n$  random variables, we say that there are mutually independent if  $P(X_i | X_S) = P(X_i)$  for all subsets  $S$  of  $\{1, 2, \dots, n\}$  which do not contain  $i$ , and therefore  $P(X_1, \dots, X_n) = P(X_1) \cdots P(X_n)$

You can write inline equations:  $a \sim p(b)$ , or one line equations:

$$a \sim p(a).$$

- (1) **Show that pairwise independence between all pairs of variables  $(X_i, X_j)$ , does NOT imply mutual independence. Note: it is enough to give an example.**

Here is an example obtained from:

<http://mnstats.morris.umn.edu/introstat/stat2611/independence.html>

SUPPOSE A BOX CONTAINS 4 TICKETS LABELLED BY  
331 323 233 333

LET US CHOOSE ONE TICKET AT RANDOM, AND CONSIDER THE RANDOM EVENTS

A1=1 OCCURS AT THE FIRST PLACE

A2=1 OCCURS AT THE SECOND PLACE

A3=1 OCCURS AT THE THIRD PLACE

$$P(A1)=1/2 \quad P(A2)=1/2 \quad P(A3)=1/2$$

$$A1A2=112 \quad A1A3=121 \quad A2A3=211$$
$$P(A1A2)=P(A1A3)=P(A2A3)=1/4.$$

So we conclude that the three events A1, A2, A3 are pairwise independent.

However

$$A1A2A3=f$$
$$P(A1A2A3)=0$$
$$P(A1)P(A2)P(A3)=(1/2)^3$$

$$\therefore P(A1A2A3) \neq P(A1)P(A2)P(A3)$$

(2) **Show mutual independence implies pairwise independence.**

For four events A, B, C, D to be mutually independent, by definition:

$$P(ABCD) = P(A)P(B)P(C)P(D),$$
$$P(ABC) = P(A)P(B)P(C),$$
$$P(ABD) = P(A)P(B)P(D),$$
$$P(ACD) = P(A)P(C)P(D),$$
$$P(BCD) = P(B)P(C)P(D),$$
$$P(AB) = P(A)P(B), \quad P(AC) = P(A)P(C), \quad P(AD) = P(A)P(D),$$
$$P(BC) = P(B)P(C), \quad P(BD) = P(B)P(D), \quad P(CD) = P(C)P(D).$$

Last two conditions must satisfy in order for these events to be pairwise independent.

**Problem 2**

(8 points) Let X and Y be two discrete random variables which are identically distributed but not necessarily independent.

Define

$$R = 1 - H(Y|X) / H(X)$$

(a) **Show that  $R = I(X,Y) / H(X)$**

$$R = 1 - H(Y|X)/H(X) = (H(X) - H(Y|X)) / H(X)$$
$$I(X,Y) / H(X) = 1 - H(X|Y) / H(X)$$

Because X and Y are identically distributed,

$$H(Y|X) = H(X|Y) \quad R = 1 - H(Y|X)/H(X)$$

$$= (H(X) - H(Y|X)) / H(X)$$

$$= (H(X) - H(X|Y)) / H(X)$$

$$= I(X, Y) / H(X)$$

(b) **Show that  $0 \leq R \leq 1$**

$$R = 1 - H(X|Y)/H(X)$$

$$0 \leq H(X|Y) \leq H(X)$$

$$\therefore 0 \leq H(X|Y)/H(X) \leq 1 \quad 0 \leq 1 - H(X|Y)/H(X) \leq 1$$

$$0 \leq R \leq 1$$

(c) **When is  $R = 0$ ?**

$$R = 0 \text{ iff } I(X, Y) = 0$$

X and Y are independent

(d) **When is  $R = 1$ ?**

$$R = 1 \text{ iff } H(X|Y) = 0$$

X is a function of Y