



ScienceDirect

Contents lists available at sciencedirect.com Journal homepage: www.elsevier.com/locate/jval

ISPOR Report

Defining Patient Engagement in Research: Results of a Systematic Review and Analysis: Report of the ISPOR Patient-Centered Special Interest Group



Rachel L. Harrington, PhD, ^{1,8} Maya L. Hanna, MPH, PhD, ² Elisabeth M. Oehrlein, MS, PhD, ³ Rob Camp, BS, ⁴ Russell Wheeler, BSc, ⁵ Clarissa Cooblall, MPH, ⁶ Theresa Tesoro, MSN, ⁶ Amie M. Scott, MPH, ⁷ Rainald von Gizycki, MA, PhD, ⁸ Francis Nguyen, PharmD, ⁹ Asha Hareendran, MA, PhD, ¹⁰ Donald L. Patrick, MSPH, PhD, ¹¹ Eleanor M. Perfetto, MS, PhD³

¹National Committee for Quality Assurance, Washington, DC, USA; ²Boehringer Ingelheim Pharmaceuticals Inc., Ridgefield, CT USA; ³National Health Council, Washington, DC, USA; ⁴Community Advisory Board Programme, EURORDIS, Barcelona, Spain; ⁵Leber's Hereditary Optic Neuropathy, Merusac, France; ⁶Scientific & Health Policy Initiatives, ISPOR, Lawrenceville, NJ, USA; ⁷Consulting, New York, NY, USA; ⁸PRO RETINA Deutschland e.V., Aachen, Germany; ⁹Bayer Healthcare Pharmaceuticals, Jersey City, NJ, USA; ¹⁰Evidera, London, England, UK; ¹¹University of Washington, Seattle, Washington, USA

ABSTRACT

Objectives: Lack of clarity on the definition of "patient engagement" has been highlighted as a barrier to fully implementing patient engagement in research. This study identified themes within existing definitions related to patient engagement and proposes a consensus definition of "patient engagement in research."

Methods: A systematic review was conducted to identify definitions of patient engagement and related terms in published literature (2006-2018). Definitions were extracted and qualitatively analyzed to identify themes and characteristics. A multistakeholder approach, including academia, industry, and patient representation, was taken at all stages. A proposed definition is offered based on a synthesis of the findings.

Results: Of 1821 abstracts identified and screened for eligibility, 317 were selected for full-text review. Of these, 169 articles met inclusion criteria, from which 244 distinct definitions were extracted for analysis. The most frequently defined terms were: "patient-centered" (30.5%), "patient engagement" (15.5%), and "patient participation" (13.4%). The majority of definitions were specific to the healthcare delivery setting (70.5%); 11.9% were specific to research. Among the definitions of "patient engagement," the most common themes were "active process," "patient involvement," and "patient as participant." In the research setting, the top themes were "patient as partner," "patient involvement," and "active process"; these did not appear in the top 3 themes of nonresearch definitions.

Conclusion: Distinct themes are associated with the term "patient engagement" and with engagement in the "research" setting. Based on an analysis of existing literature and review by patient, industry, and academic stakeholders, we propose a scalable consensus definition of "patient engagement in research."

Keywords: definition, health economics, outcomes research, patient centered, patient centric, patient engagement, patient research.

VALUE HEALTH. 2020; 23(6):677-688

Introduction

Over the past few decades, the concepts of "patient engagement" and "patient-centeredness" have risen to prominence across the healthcare ecosystem. Rather than serving merely as research subjects, the patient community increasingly has the capacity to partner or lead in research. At a systems level, meaningful involvement of patients and increased understanding of patient experiences, goals, and needs through partnerships in the research process leads to more relevant and effective research. Better understanding and incorporation of patient

perspectives through early and continuous partnerships has the potential to enhance the impact of research. This, in turn, results in better, more patient-relevant outcomes because of easier recruitment, retention of study participants, and improved interpretation, which may contribute to faster and broader access to new and better treatment options. ^{4,5} At an individual level, data that are generated with patient engagement can ensure the availability of patient-relevant information, enhancing active patient participation in their healthcare decision making. ⁶

The role of patients as partners in health technology development has increased over the past decade, driven by the

expectation of regulatory bodies, health technology assessment (HTA) authorities, and patients themselves that patients' needs and priorities be understood.^{7–10} This patients-as-partners model has also been taken up by industry and others in the recognition that it can lead to better decisions and outcomes, more efficient processes, improved patient recruitment and participation, and shorter lead times.⁷ These developments are reflected in the adoption of laws and policies related to patient-focused drug development in both the United States and the European Union; changes to regulatory standards on clinical outcome assessments; and the growing global impact of patient advocates and citizen scientists on regulatory decision making. 11-16 These efforts are complementary to the growing role of patient engagement and centricity in healthcare research and delivery. Research organizations have begun to incorporate patients across all levels of decision making, including agenda setting, project planning, implementation, and evaluation.¹⁷ In the United States, this has been influenced by the establishment of the Patient-Centered Outcomes Research Institute, the Agency for Healthcare Research and Quality, and healthcare quality efforts. 18,19 In Europe, this has been reflected in the work of organizations, such as the European Patients' Academy and the European Organization for Rare Diseases, to champion the empowerment of patients to participate as leaders, partners, and decision makers in the development of health technologies.^{20,21} Active efforts to promote engagement of patients and the public in health research and technology assessment are also ongoing in other regions including, but not limited to, Latin America and Asia. 22,23

There is broad agreement that patient engagement should be meaningful, impactful, and measurable; however, terms related to patient engagement and patient-centeredness are used inconsistently across stakeholders and contexts.^{7,24,25} These terms are rarely defined and may be used synonymously with related, but distinct, concepts. For example, patient engagement is frequently used interchangeably with patient activation, patient involvement, and patient-centered care.¹⁹ A horizon scan of terms and definitions related to patient engagement identified 24 terms, each with multiple definitions across healthcare sectors (eg, government agencies, payers, and patient advocacy groups).²⁶ Additionally, these terms and definitions may be context specific, with variable meanings. As such, it is possible that a high-quality definition of patient engagement in the context of healthcare delivery context may not always be appropriate for use in the context of research.

Health economics and outcomes research (HEOR) stake-holders, including researchers, clinicians, patients, payers, regulators, and industry, are seeking to meaningfully partner with patients to incorporate patient input into their research.^{27–29}

Clarity on what patient centeredness and patient engagement mean in the context of health economics and outcomes research is increasingly relevant for members of the International Society for Pharmacoeconomics and Outcomes Research (ISPOR) and similar research organizations.²⁹ To ensure clear and consistent communication, an ISPOR standard definition for these terms is desirable for use in ISPOR communications, task forces, and other Society initiatives. Such a definition would also support efforts to identify and measure quality patient engagement activities.


To this end, the Patient Engagement in Research Working Group of the ISPOR Patient-Centered Special Interest Group (PCSIG) set out to conduct a comprehensive review of existing definitions of patient engagement, patient-centeredness, and related concepts, with the goal of generating a robust, evidence-based, scalable definition of patient engagement fit for use in the context of health research.

Methods

A multistep data extraction, review, and analysis process was undertaken to generate a final definition (Fig. 1). It was determined from the outset that, to ensure an adequate identification and coding process, a multistakeholder approach representing the perspectives of researchers (both academic and industry) and patients was required. To this end, a study team representing all of these groups was maintained across design, methods selection, analysis, and interpretation. It included both the systematic review and qualitative analysis research teams, as well as the stakeholders who reviewed the final definition.

Before initiating the systematic review, a targeted web search was first conducted to identify relevant terms, definitions, and frameworks in grey literature (eg, web-based documents from patient advocacy groups and umbrella patient advocacy organizations, as well as health authorities and regulators). Grey literature was deemed important for this scoping review given the possible underrepresentation of these stakeholders in the published literature. This literature was identified through either a recommendation from the authors or a general text search using Google and Google Scholar to identify relevant definitions from other web sources. The search was limited to the English language. As a result, definitions for patient engagement and patient-centeredness represented by public and private organizations globally were recorded and new patient-related terms were identified for inclusion in the systematic review search criteria. These definitions were also used to externally validate findings from the systematic review and to support further definition development.

Figure 1. Diagram depicting the multistep process followed to achieve final proposed definition of "patient engagement in research."



Systematic Review

We conducted a PubMed and EMBASE database search for titles and abstracts that included the terms "patient engagement," "patient-centeredness," or related patient terms in published literature from January 1, 2006, through December 31, 2016 (see Appendix 1 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2 020.01.019). The choice of 2006 as the start of the review period was based on an examination of the prevalence of the terms of interest in the peer-reviewed literature, which identified an increase in the mid-2000s. In addition, because terms have evolved over time, the intent was to focus on contemporary usage and meanings. To reflect the evolving nature of this topic, an update of the review specific to the "patient engagement" and "research" search criteria was conducted for the period between January 1, 2017, and December 31, 2018. Results of this extended search were integrated into the complete analysis. Published literature was limited to the English language to accurately interpret and code the definitions. To manage the anticipated high volume of articles, the full review team consisted of 16 PCSIG members who were assigned in pairs to independently screen each abstract for relevance. Any disagreements were resolved by consensus. A third researcher resolved any remaining discrepancies.

Criteria for selecting abstracts for full-text review were established a priori and defined as follows.

Inclusion Criteria:

- 1. At least 1 definition of a term was in the abstract.
- The purpose/objective of the article was to define the term or introduce, examine, or analyze a conceptual model or framework.
- 3. The article was a systematic review of the term definition or concept.
- 4. It was an opinion article about the term's use or importance in healthcare or health research.
- 5. It was unclear whether to include it without a full-text review.
- 6. The publication year was 2006 or later.

Exclusion Criteria:

- 1. No patient terms were defined or framework was referenced.
- 2. It was not healthcare or health-research related.
- 3. The term was used to describe a treatment approach or intervention in a specified population (eg, colorectal surgery) without defining the term.
- 4. The term was used in the context of improved communications or infrastructure of an organization through information technology (eg, patient portals, electronic health records [EHRs], devices, equipment, web-based tools) without defining the term.

Abstracts designated for full-text article review were then assessed independently by 2 data abstractors. During full-text review, the same inclusion and exclusion criteria were applied to identify the final analytic set. Identified terms and their definitions were categorized by whether the definition was adapted or adopted from another referenced source or the article authors provided an original definition. Abstractors also recorded the context in which the definitions were used, (ie, healthcare delivery, research, both, or other). If the article provided a framework for the definition, this was also recorded. A single article could contribute more than 1 definition.

To account for definitions that were adopted multiple times from the same referenced source, either the source reference of the original definition or the first published record of the definition in the study period was retained in the final analysis. This deduplication process resulted in additional exclusions.

Qualitative Content Analysis

Definitions for each patient-related term abstracted from a full-text article review underwent qualitative content analysis (QCA).³⁰ The full review team consisted of 6 individuals. Reviewers, assigned in pairs, assessed each definition, with a third reviewer available to resolve any discrepancies. Analysis was conducted as an iterative process, with the coding framework (ie, codebook) derived inductively from the definition content.³¹ Microanalysis, a detailed form of open coding used to generate ideas, was performed on a random 30% sample of definitions to extract and consolidate concepts and develop a preliminary codebook.³² This initial codebook was piloted on a subset of definitions and was extended and modified based on reviewer feedback to generate a final coding dictionary (Table 1; see also Appendix 2 in Supplemental Materials found at https://doi.org/1 0.1016/j.jval.2020.01.019). Complete coding on the full definition set was then conducted.

The stakeholders involved in engagement (the "who") were differentiated based on their identified role, either as the party responsible for conducting engagement (the conductor) or the party targeted to be engaged (the target). Stage of activity (the "when") was conceptualized as a timeline. Stages were identified (from earliest to latest) as: "Strategy & Priority Setting," "Design & Planning," "Conduct & Operation," "Dissemination & Communication," and "In Practice/In Use." Additionally, the stage could be identified as "All" should a definition refer to all stages across the timeline.

Upon finishing coding and reaching saturation for each patient term (no new concepts identified within the term), the resulting codes were discussed with the full research team to examine the similarities and dissimilarities of coded content within and between patient terms. As a result, similar or overlapping codes were grouped into meaningful attributes, such as "communication: accessible," "decision making: shared" and "empowerment" (for further detail, see Appendices 2 and 3 in Supplemental Materials). Patient terms with near-identical attributes were consolidated into one patient term.

Descriptive analyses, including frequency and relative ranks of coded concepts, were conducted overall by patient term and by primary coding categories (eg, context of use, target audience of engagement, stage of activity). Definition coding was conducted using Microsoft Excel and analysis was performed using SAS® version 9.4.

Definition Review and Consensus

After completion of QCA, a preliminary definition of "patient engagement in research" was generated based on the most prevalent coded characteristics of definitions of patient engagement and definitions in the research setting. This preliminary definition was then reviewed by the ISPOR PCSIG Leadership Group to ensure conceptual clarity, minimize repetition, and supplement with relevant context from other defined terms. This revised definition was proposed to the full PCSIG (n = 447 internationally) and to the participants of the past 10 ISPOR Patient Representatives Roundtables (n = 157), including North America, Europe, and Latin America. A final consensus definition was derived based on stakeholder feedback.

Results

Systematic Review

A total of 1882 articles were identified in the search (PUBMED: 1405; EMBASE: 334; Other: 8). After deduplication, 1821 unique abstracts were screened for relevance, and ultimately 317 full-text

Table 1. Qualitative analysis codebook structure.

| Category | Content | | No. of Allowed [*] Definition | Terms per | No. of Terms in Code List [†] |
|-------------------------------|---|-----|---|-----------|---|
| Defined Term (What) | The search term that produced the definition | Yes | | 1 | 12 |
| Characteristics (How) | The defined process, activities or expectations | Yes | 8 | 3 | 51 |
| Rationale (Why) | The defined goals of the activity | No | 3 | 3 | 51 |
| Setting (Where) | The defined setting or context of use | No | 2 | 2 | 7 |
| Conductor of Engagement (Who) | The defined conductor ("engager") the activity | No | Ē | 3 | 7 |
| Target of Engagement (Who) | The defined target ("engagee") the activity | No | 3 | 3 | 6 |
| Stage of Activity (When) | The defined point in the process at which the activity occurs | No | 3 | 3 | 6 |

^{*}Required indicates at least one term must be coded in the field for the definition to be included in the analysis and considered "informative."

articles were reviewed for definitions. Out of a final set of 169 articles, 244 distinct definitions across all search terms were identified for analysis (Fig. 2). The most frequently referenced "adopted" definition was from the Institute of Medicine, followed by the one by Stewart et al.^{34,35} The majority of articles were published between 2012 and 2015, with 2012 being the year with the highest number of publications offering definitions (Table 2). The search term contributing the most articles was "patient-centered" (26.8%), followed by "patient engagement" (16.7%).

Qualitative Content Analysis

Qualitative content analysis was conducted on 244 definitions. Of those, 17.2% were extracted from articles identified by the search term "patient engagement," 43.4% identified by the search term "patient-centered," and 39.3% split among other terms. The most frequent setting of the activity, overall, was healthcare at 70.5%, with the research setting identified in only 11.9% of definitions.

The 3 most common characteristics, overall, were "patient as participant, "patient involvement," and "incorporate patient: preferences" and "relationship: patient-provider" (tied). Although the top 10 characteristics were similar across defined terms, relative rankings differed (Table 3). The most prevalent characteristics among definitions of patient engagement were "process: active," "patient involvement," and "patient as participant." The top patient-centered definition characteristics were "incorporate patient: preferences," "incorporate patient: perspective," and "relationship: patient-provider." There were also differences between the settings (Table 4). The top 3 characteristics of definitions in the research setting were "patient involvement," "patient as partner," and "process: active." In healthcare, they were "relationship: patient-provider," "patient as participant," and "incorporate patient: preferences." Four characteristics appeared in the overall top 10 and were never coded among definitions in the research setting: "individualized approach: care," "patient in control," "patient as person," and "respect."

Overall, "patients" were the most common target of engagement (83.7%). Among these definitions, the target was further defined in a majority of cases. The most common characterization of patients was "patient community, in general" (38.8%), "individual patient" (29.6%), and "patient advocacy/advocates" (1.5%). Family or caregivers were identified as a target of the

activity in only 8.2% of definitions (n = 19 and n = 1, respectively). When possible, conductors of engagement were identified within definitions. The most common were healthcare practitioners (35.7%) and the healthcare system (22.1%). Patients, traditionally a "target" of engagement, were identified as the party responsible for conducting engagement activities in 17.2% of definitions.

The most common stage of activity identified among definitions was "In Practice/In Use" (the latest stage). Only 5.7% of definitions identified activities at the "Strategy & Priority Setting" stage; however, those that fell into the patient engagement term definition and research setting had the highest proportion compared with other terms or settings (9.5% and 10.3%, respectively). Among definitions in the research setting, 34.5% identified activity across "All Stages," compared with only 2.9% of definitions in the Healthcare setting.

Generally, the proportion of definitions in which a characteristic could not be identified (ie, "Undetermined") was less than 20%: "Setting of Activity (13.1%), "Target of Engagement" (13.9%), and "Stage of Activity" (19.3%). "Conductor of Activity," on the other hand, could not be determined in 36.1% of definitions.


We also examined the overlap of concepts between the term "patient engagement" and the research setting. The term "active process" was ranked first among concepts related to the defined term "patient engagement" and third among defined terms in the research setting. Similarly, "meaningful to patient" was ranked fifth and seventh, respectively. Although the overall occurrence of engagement across "All Stages" was low (6.6%), it was the most prevalent stage of engagement among research setting definitions (34.5%) and the highest among the definitions of "patient engagement" (11.9%), compared with other terms.

Proposed Definition of Patient Engagement in Research

A working definition was constructed based on cross-tabulated frequencies and shared ranking of coded terms in the patient engagement and research categories. The authors made modifications to this QCA-constructed definition, informed by results from the grey literature scan and by review of prevalent themes unique to the defined term "patient engagement" or the setting of research, excluding overlapping themes. ^{36–39} This revised definition was then provided to the full PCSIG membership, as well as to members of the 4 regional ISPOR Patient Representatives

[†]Reflects number of high-level terms in each code list. Refer to Appendix 2 in Supplemental Materials for additional information.

Figure 2. Identification and selection of articles into systematic review and analysis attrition diagram. *Refer to Appendix 1 in Supplemental Material for details on exclusion reasons and counts.



Roundtables (see Appendix 4 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2020.01.019). Comments were received from 24 reviewers, who self-identified as representing the following perspectives: patient representatives (42%), academicians/researchers (29%), industry (12.5%), health technology

assessment (HTA) bodies (8%), regulatory/government (4%), and other (4%). The following regions were represented: North America (32%), Latin America (32%), Europe (24%), Asia (8%). The PCSIG Leadership Group then reviewed all comments and feedback, from which a final consensus definition was derived.

Table 2. Characteristics of articles identified by systematic review.

| | N | 6 1 B 3 | 5.6 |
|-----------------------|------------------|-----------------------------|-----------------------------|
| Search Terms | No. of Articles* | Average year of publication | References |
| Overall | 169 | 2012 | - |
| Patient-Centered | 49 | 2012 | 19,51–98 |
| Patient Engagement | 30 | 2014 | 2,13,19,99–125 |
| Patient Involvement | 25 | 2011 | 91,126–149 |
| Patient Participation | 24 | 2011 | 55,66,69,140,150–169 |
| Patient Empowerment | 21 | 2011 | 52,158,170–188 |
| Patient-Centeredness | 18 | 2012 | 13,70,79,94,106,179,189–200 |
| Person-Centered | 10 | 2012 | 88,201–209 |
| Patient Focused | 4 | 2012 | 13,203,210,211 |
| Patient-Centric | 3 | 2009 | 212–214 |
| Consumer Involvement | 2 | 2008 | 63,215 |
| Patient Input | 1 | 2011 | 216 |
| Patient Oriented | 1 | 2013 | 217 |
| Consumer Engagement | 1 | 2010 | 218 |

Note. For purposes of analysis and presentation of results, the following terms were combined: Patient Focused & Patient-Focused Drug Development & Patient-Focused Outcomes Research; Patient-Centered & Patient-Centered Outcomes Research

^{*}A single article could contribute multiple definitions. The overall total reflects the total number of distinct articles. The term-specific number of articles reflects the number of distinct articles contributing a definition for that term.

Table 3. Identified characteristics of definitions, overall and by defined term.

| Definitions, Overall | Overall | | | Patient Engagement | | | Patient-Centered* | | | Other | | |
|--|--------------------------------------|--------------------------------------|--|---------------------------------|------------------------------------|---|---------------------------------|------------------------------------|---|--------------------------------------|--|--|
| | Rank | n | % | Rank | n | % | Rank | n | % | Rank | n | % |
| | | 244 | | | 42 | 17.2 | | 106 | 43.4 | | 96 | 39.3 |
| Top Characteristics of Activity | | | | | | | | | | | | |
| patient as participant | 1 | 43 | 17.6 | 3 | 9 | 21.4 | 8 | 13 | 12.3 | 1 | 21 | 21.9 |
| patient involvement | 2 | 36 | 14.8 | 2 | 10 | 23.8 | 8 | 13 | 12.3 | 4 | 13 | 13.5 |
| incorporate patient: preferences | 3 | 35 | 14.3 | 6 | 4 | 9.5 | 1 | 26 | 24.5 | 11 | 5 | 5.2 |
| relationship: patient-provider Decision making: shared | 3 4 | 35 33 | 14.3 13.5 | 6 5 | 4 6 | 9.5 14.3 | 3 8 | 21 13 | 19.8 12.3 | 6 3 | 10 14 | 10.4 14.6 |
| patient as partner | 5 | 32 | 13.1 | 4 | 8 | 19.0 | 7 | 14 | 13.2 | 6 | 10 | 10.4 |
| incorporate patient: perspective | 6 | 27 | 11.1 | 9 | 1 | 2.4 | 2 | 23 | 21.7 | 13 | 3 | 3.1 |
| process: active | 6 | 27 | 11.1 | 1 | 12 | 28.6 | 19 | 2 | 1.9 | 4 | 13 | 13.5 |
| individualized approach: care | 7 | 22 | 9.0 | 9 | 1 | 2.4 | 4 | 18 | 17.0 | 13 | 3 | 3.1 |
| information sharing | 8 | 20 | 8.2 | 7 | 3 | 7.1 | 17 | 4 | 3.8 | 4 | 13 | 13.5 |
| patient in control | 8 | 20 | 8.2 | 8 | 2 | 4.8 | 19 | 2 | 1.9 | 2 | 16 | 16.7 |
| understanding | 8 | 20 | 8.2 | 9 | 1 | 2.4 | 5 | 16 | 15.1 | 13 | 3 | 3.1 |
| patient as person | 9 | 17 | 7.0 | 9 | 1 | 2.4 | 6 | 15 | 14.2 | 15 | 1 | 1.0 |
| Decision making: informed/competent | 10 | 16 | 6.6 | 7 | 3 | 7.1 | 15 | 6 | 5.7 | 9 | 7 | 7.3 |
| empowerment | 10 | 16 | 6.6 | 7 | 3 | 7.1 | 20 | 1 | 0.9 | 5 | 12 | 12.5 |
| process: adaptive/responsive | 10 | 16 | 6.6 | n/a | 0 | 0.0 | 10 | 11 | 10.4 | 11 | 5 | 5.2 |
| respect | 10 | 16 | 6.6 | 9 | 1 | 2.4 | 9 | 12 | 11.3 | 13 | 3 | 3.1 |
| Stage of Activity ^T 5) In Practice/In Use Undetermined 2) Design & Planning All | 1 2 3 4 | 156 47 30 16 | 63.9 19.3 12.3 6.6 | 1 3 2 4 | 22 9 10 5 | 52.4 21.4 23.8 11.9 | 1 2 3 5 | 84 14 11 3 | 79.2 13.2 10.4 2.8 | 1 2 3 4 | 50 24 9 8 | 52.1 25.0 9.4 8.3 |
| Strategy & Priority Setting Conduct & Operation Dissemination & Communication | 5 6 7 | 14 13 6 | 5.7 5.3 2.5 | 5 5 6 | 4 4 2 | 9.5 9.5 4.8 | 4 7 6 | 4 1 2 | 3.8 0.9 1.9 | 5 4 6 | 6 8 2 | 6.3 8.3 2.1 |
| Setting of Activity Healthcare Undetermined Research | 1 2 3 | 172 32 29 | 70.5 13.1 11.9 | 1 3 2 | 27 2 12 | 64.3 4.8 28.6 | 1 2 3 | 88 12 5 | 83.0 11.3 4.7 | 1 2 3 | 57 18 12 | 59.4 18.8 12.5 |
| Other | 4 | 11 | 4.5 | 4 | 1 | 2.4 | 4 | 1 | 0.9 | 4 | 9 | 9.4 |
| Target of Engagement Patient Undetermined Family Public Provider Stakeholder (otherwise undefined) Caregiver | 1 2 3 4 5 6 7 | 206 34 19 15 9 8 | 84.4 13.9 7.8 6.1 3.7 3.3 0.4 | 1 2 3 5 6 4 7 | 38 6 5 2 1 3 0 | 90.5 14.3 11.9 4.8 2.4 7.1 0.0 | 1 3 2 6 4 5 7 | 99 8 10 2 4 3 | 93.4 7.5 9.4 1.9 3.8 2.8 0.9 | 1 2 4 3 4 5 6 | 69 20 4 11 4 2 0 | 71.9 20.8 4.2 11.5 4.2 2.1 0.0 |
| Conductor of Engagement Undetermined Healthcare Practitioner Healthcare System Patient Researcher Provider Organization Government Industry / Commercial | 1 2 3 4 5 6 7 8 | 88 87 54 42 21 5 4 | 36.1 35.7 22.1 17.2 8.6 2.0 1.6 1.2 | 1 5 4 3 2 6 7 | 13 8 9 10 11 2 0 | 31.0 19.0 21.4 23.8 26.2 4.8 0.0 0.0 | 2 1 3 4 5 6 7 | 33 56 28 7 4 3 0 | 31.1 52.8 26.4 6.6 3.8 2.8 0.0 0.0 | 1 3 4 2 5 8 6 7 | 42 23 17 25 6 0 4 3 | 43.8 24.0 17.7 26.0 6.3 0.0 4.2 3.1 |

 $\it Note.$ Percentage represents percent of definitions in which the term appears.

As a result of this process, we propose the following definition of "patient engagement in research":

The active, meaningful, and collaborative interaction between patients and researchers across all stages of the research process, where research decision making is guided by patients' contributions as partners, recognizing their specific experiences, values, and expertise.

It is important to clarify key concepts within this definition. First, the term "patients" is used inclusively to represent

individual patients, their families, and their caregivers, in addition to patient representatives and advocacy organizations. This is in line with other work in this space. 40,41 The term "researchers" is also used broadly to refer to multiple contributors within the process, such as research producers (those directly conducting the research activity) and decision-making consumers (such as payers and regulators). Research funders, through their role in priority setting and protocol review, should also be held accountable for ensuring engagement is

^{*}Combination of patient: centered, centeredness, centric, and person-centered

[†]Note that a single definition could have multiples stages, targets, and conductors, so percentages may add up to more than 100%.

Table 4. Identified characteristics of definitions, overall and by setting of activity.

| | Overall | | | Research | | | Healthcare | | | Other* | | |
|-------------------------------------|---------|-----|------|----------|----|------|------------|-----|------|--------|----|------|
| Definitions, Overall | Rank | | % | Rank | | % | Rank | | % | Rank | | % |
| | | 244 | - | | 29 | 11.9 | | 172 | 70.5 | | 43 | 17.6 |
| Top Characteristics of Activity | | | | | | | | | | | | |
| patient as participant | 1 | 43 | 17.6 | 4 | 5 | 17.2 | 2 | 53 | 30.8 | 1 | 21 | 48.8 |
| patient involvement | 2 | 36 | 14.8 | 2 | 8 | 27.6 | 4 | 37 | 21.5 | 1 | 21 | 48.8 |
| incorporate patient: preferences | 3 | 35 | 14.3 | 7 | 2 | 6.9 | 3 | 52 | 30.2 | 6 | 10 | 23.3 |
| relationship: patient-provider | 3 | 35 | 14.3 | 7 | 2 | 6.9 | 1 | 54 | 31.4 | 5 | 11 | 25.6 |
| decision making: shared | 4 | 33 | 13.5 | 7 | 2 | 6.9 | 6 | 32 | 18.6 | 2 | 19 | 44.2 |
| patient as partner | 5 | 32 | 13.1 | 1 | 10 | 34.5 | 7 | 30 | 17.4 | 3 | 15 | 34.9 |
| incorporate patient: perspective | 6 | 27 | 11.1 | 5 | 4 | 13.8 | 5 | 34 | 19.8 | 7 | 9 | 20.9 |
| process: active | 6 | 27 | 11.1 | 3 | 6 | 20.7 | 8 | 29 | 16.9 | 5 | 11 | 25.6 |
| individualized approach: care | 7 | 22 | 9.0 | 9 | 0 | 0.0 | 8 | 29 | 16.9 | 6 | 10 | 23.3 |
| information sharing | 8 | 20 | 8.2 | 6 | 3 | 10.3 | 12 | 19 | 11.0 | 3 | 15 | 34.9 |
| patient in control | 8 | 20 | 8.2 | 9 | 0 | 0.0 | 12 | 19 | 11.0 | 3 | 15 | 34.9 |
| understanding | 8 | 20 | 8.2 | 8 | 1 | 3.4 | 11 | 20 | 11.6 | 4 | 13 | 30.2 |
| patient as person | 9 | 17 | 7.0 | 9 | 0 | 0.0 | 10 | 21 | 12.2 | 7 | 9 | 20.9 |
| decision making: informed/competent | 10 | 16 | 6.6 | 8 | 1 | 3.4 | 13 | 18 | 10.5 | 9 | 7 | 16.3 |
| empowerment | 10 | 16 | 6.6 | 7 | 2 | 6.9 | 10 | 21 | 12.2 | 10 | 6 | 14.0 |
| process: adaptive/responsive | 10 | 16 | 6.6 | 8 | 1 | 3.4 | 12 | 19 | 11.0 | 8 | 8 | 18.6 |
| respect | 10 | 16 | 6.6 | 9 | 0 | 0.0 | 9 | 23 | 13.4 | 11 | 5 | 11.6 |
| Most Prevalent Stage of Activity | | | | | | | | | | | | |
| 5) In Practice/In Use | 1 | 156 | 63.9 | 3 | 4 | 13.8 | 1 | 141 | 82.0 | 2 | 11 | 25.6 |
| Undetermined | 2 | 47 | 19.3 | 3 | 4 | 13.8 | 2 | 18 | 10.5 | 1 | 25 | 58.1 |
| 2) Design & Planning | 3 | 30 | 12.3 | 2 | 9 | 31.0 | 2 | 18 | 10.5 | 3 | 3 | 7.0 |
| All | 4 | 16 | 6.6 | 1 | 10 | 34.5 | 5 | 5 | 2.9 | 5 | 1 | 2.3 |
| 1) Strategy & Priority Setting | 5 | 14 | 5.7 | 4 | 3 | 10.3 | 3 | 8 | 4.7 | 3 | 3 | 7.0 |
| 3) Conduct & Operation | 6 | 13 | 5.3 | 3 | 4 | 13.8 | 4 | 7 | 4.1 | 4 | 2 | 4.7 |
| 4) Dissemination & Communication | 7 | 6 | 2.5 | 3 | 4 | 13.8 | 6 | 2 | 1.2 | 6 | 0 | 0.0 |
| Target of Engagement | | | | | | | | | | | | |
| Patient | 1 | 206 | 84.4 | 1 | 26 | 89.7 | 1 | 156 | 90.7 | 1 | 24 | 55.8 |
| Undetermined | 2 | 34 | 13.9 | 4 | 3 | 10.3 | 3 | 16 | 9.3 | 2 | 15 | 34.9 |
| Family | 3 | 19 | 7.8 | 5 | 1 | 3.4 | 2 | 17 | 9.9 | 4 | 1 | 2.3 |
| Public | 4 | 15 | 6.1 | 2 | 5 | 17.2 | 5 | 4 | 2.3 | 3 | 6 | 14.0 |
| Provider | 5 | 9 | 3.7 | 5 | 1 | 3.4 | 4 | 8 | 4.7 | 5 | 0 | 0.0 |
| Stakeholder (otherwise undefined) | 6 | 8 | 3.3 | 3 | 4 | 13.8 | 6 | 3 | 1.7 | 4 | 1 | 2.3 |
| Caregiver | 7 | 1 | 0.4 | 5 | 1 | 3.4 | 7 | 0 | 0.0 | 5 | 0 | 0.0 |
| Conductor of Engagement | | | | | | | | | | | | |
| Undetermined | 1 | 88 | 36.1 | 3 | 5 | 17.2 | | 51 | 29.7 | | 32 | 74.4 |
| Healthcare Practitioner | 2 | 87 | 35.7 | 4 | 2 | 6.9 | | 84 | 48.8 | | 1 | 2.3 |
| Healthcare System | 3 | 54 | 22.1 | 4 | 2 | 6.9 | | 46 | 26.7 | | 6 | 14.0 |
| Patient | 4 | 42 | 17.2 | 2 | 6 | 20.7 | | 28 | 16.3 | | 8 | 18.6 |
| Researcher | 5 | 21 | 8.6 | 1 | 20 | 69.0 | | 0 | 0.0 | | 1 | 2.3 |
| Provider Organization | 6 | 5 | 2.0 | 6 | 0 | 0.0 | | 5 | 2.9 | | 0 | 0.0 |
| Government | 7 | 4 | 1.6 | 6 | 0 | 0.0 | | 2 | 1.2 | | 2 | 4.7 |
| Industry / Commercial | 8 | 3 | 1.0 | 5 | 1 | 3.4 | | 1 | 0.6 | | 1 | 2.3 |
| maasay / commercial | U | J | 1.4 | J | | ٦.┯ | | ' | 0.0 | | | ۷.5 |

^{*}Combination of patient *centered, *centeredness, *centric, and person-centered

appropriately defined. Different patient and researcher representatives may be involved at different stages in the research process. For example, funding bodies may be more involved during priority setting stages whereas payers may be more involved in implementation and dissemination.

By incorporating "all stages of the research process," we intend to cover the full spectrum of research activities, including planning, conduct, and dissemination. The phrase "all stages" is used to indicate that patient engagement may be possible, and should be considered, at any point in a study. Nevertheless, it is not meant to function as a mandate that engagement must happen at all stages of a single study, which may be neither feasible nor appropriate. By characterizing

engagement as "active, meaningful, and collaborative," the definition reflects the importance of substantive, versus superficial, interactions. 42,43

Finally, it is essential to describe what is meant by "patients' contributions." This concept is not intended to reflect patient-level clinical information collected as part of a research study, (ie, the patient as a passive "data point" contributor). Rather, patient contributions include capturing and utilizing patient perspectives, preferences, experiences, opinions, and inputs into the research itself. These may be captured formally (eg, preference elicitation methods, focus groups, advisory board membership), or informally (eg, through routine interactions that evolve as part of research partnerships).

Discussion

We propose a standardized definition of "patient engagement in research" derived from the results of our analysis, further informed by the results of an environmental scan and by the review of multiple stakeholder groups. It is intended to be aspirational and represent what full patient engagement in research should look like. The goal of engagement should be a partnership, where patient contributions are given equal weight to those of other contributors to the research team. We recognize that external constraints may present challenges at first, and that different forms of engagement may be needed to support different types of research activities. Results of our review emphasize the importance of inclusion, respect, and equality in the interactions between patients and researchers.

Previous studies have highlighted the (possibly inappropriate) interchangeable use of different terms related to patient engagement. ^{19,44} This was substantiated by our analysis, where a clear distinction emerged between the terms "patient-centered" and "patient engagement." "Patient-centered" was more focused on the healthcare setting in a patient-provider context. Compared with other defined terms, "patient engagement" was most strongly associated with an active, involved process. We also found evidence of comprehensive engagement across stages in the research setting compared with a healthcare setting.

Our study benefits from a number of strengths. A multistakeholder approach was taken at all stages, from study design through execution and dissemination, with a focus on transparency and collaboration. This is the first study to conduct a systematic review or qualitative analysis of the literature in this space. Rubrics and frameworks can be critical for measurement. Nevertheless, the lack of a clear, concise definition has presented a barrier to uptake as stakeholders, particularly on the researcher side, struggle to implement patient engagement in research in their organizations. We conducted an analysis of concepts unique to research to allow for a targeted definition, rather than relying on concepts more appropriate to the provision of patient-level healthcare. The proposed definition is concise and can be used without reference to external frameworks or conceptual models. Our definition aligns closely with ongoing complementary initiatives by researchers, health authorities, and patient organizations. This includes distinguishing between "patient-provided information" (eg, patient preferences, feedback on design) and "patient medical information" (eg, clinical endpoints or measures), the incorporation of "specific knowledge and expertise" brought by patients to scientific discussion, and the importance of engagement throughout the entire research process. 15,25,29,41,45

Results of our review are subject to limitations. This review was limited to English language literature. Although the volume of articles suggests this represents a large share of the work in this area, there is a possibility that unique concepts not present in the English literature were missed. Moreover, the volume of articles and definitions to be reviewed, and challenges related to the availability of study team members who volunteered their time, meant it took a considerably longer time to execute the study than originally anticipated. As a result, the literature review (which included publications available as of December 31, 2018) does not reflect more recent publications. Nevertheless, the previously highlighted consistency of our results with contemporary work in this space suggests consistency in themes. Given the large scope of the published literature review, full qualitative content analysis was not conducted on sources from the nonpublished literature (eg, patient advocacy websites and white papers). The final

proposed definition, however, was reviewed against the grey literature identified by the complementary web search. Finally, QCA coding terms were defined based on a detailed microanalysis of the review-identified definitions. It is possible that coding terms were identified that reflected the reviewer's understanding, but not the author's original intent, for a given definition. The discrete nature of the coding may also not reflect the full nuances of the field.

As a next step, results of this work could be used to inform companion definitions to the one proposed for "patient engagement in research." In particular, a definition of "patient-centered research" would help provide context for use of these separate terms and explore alignment between concepts.

Conclusions

The ability for patient engagement to improve research is only as good as how it is defined and operationalized and will be limited if stakeholder understanding of engagement is convoluted or misunderstood. Based on our analysis from a systematic literature review, we offer a standardized definition of "patient engagement in research" that can be adopted by various stakeholders across the research continuum.

Several next steps should be taken to continue moving understanding and implementation of patient engagement in research forward. Most pressingly, there is a need to determine how to operationalize, implement, and measure the success of patient-engagement initiatives and to learn from any challenges. Such an activity must account for what "good" engagement looks like from both the researcher and patient perspectives. This work has already begun among some stakeholder groups, including professional and research organizations, who play an important role in shaping the dialogue and scientific standards in their fields. 45-48 For example, the Patient-Centered Outcomes Research Institute Engagement Rubric provides examples of how to operationalize the role of "patient as partner." Revisiting activities based on a standardized definition and understanding of engagement would assist in moving the process forward. The time and resources needed for patients to engage in research must be considered and respected, with appropriate support (financial and otherwise) and training provided. Finally, field-specific gaps in the uptake of engagement should be identified, and work should be undertaken to resolve them. ⁴⁹ For instance, the lack of relevance of research to patients has been noted as a barrier to patient use of comparative effectiveness research.⁵⁰ This includes failure to communicate in lay language.

These future activities must not occur in siloes. Many stakeholders in this space are contributing to the evolving science and philosophy of patient engagement. This breadth of contributions brings many benefits but also the risk of fragmentation. Collaboration between contributors across stakeholder perspectives, regions, and organizations can lead to a more robust and consistent path forward. Finally, it should be noted that patient engagement in research is not a static concept. Although our proposed definition is aspirational, there remains room for further development as the research environment continues to evolve. Patients and patient organizations have demonstrated a strong desire and growing capability to become drivers and leaders in research. The outlook for patient engagement in research may look different in the near future. It is incumbent upon researchers to consider the impact of this change and not only ask how patients can inform their research agenda, but also how they can support the research agenda of patients directly.

Acknowledgments

This article was developed as a collaborative effort of members of the ISPOR Patient-Centered Special Interest Group (SIG). We thank all reviewers who commented during our roundtable discussions, forums, and open meeting presentations at the ISPOR conferences. We especially thank the individuals who submitted written comments on drafts of our manuscript. Both oral and written feedback significantly improved the quality of this report and contributed to the consensus nature of this ISPOR SIG manuscript. The following representatives from the ISPOR Patient Representatives Roundtables and ISPOR Patient-Centered SIG submitted feedback: Suzanne Schrandt, Robert McBurney, Jill Bell, Nathalie Bere, Cindy Coney, Nigel Cook, Marcy Fitz-Randolph, Elizabeth Franklin, Zachary Hallinan, Dominique Hamerlijnck, Abner Lobao, Gurmit Sandhu, Yu Wang, Durhane Wong-Rieger, Alyce Adams Patrizia Allegra, Michael Cangelosi, Tamryn Gray, Hanna Gyllensten, Yu Heng Kwan, Ilya Ivlev, Sarah Knight, Yvonne Lis, Joey Mattingly, Celine Pribil, and Ajaree Rayanakorn. We also would like to thank the following individuals for their support with reviewing, and data cleaning and management: Violeta Astratinei, Aida Kuzacan, O'Mareen Spence, Shannon Marrow, Hager Elgendi, and Vrushali Patil.

Source of financial support

None of the authors received support for their participation in this SIG. All authors volunteered their time for the discussion, research, and writing of this report. This research was supported in part by ISPOR, which contributed 2 staff liaisons for this project.

Supplemental Material

Supplementary data associated with this article can be found in the online version at https://doi.org/10.1016/j.jval.2020.01.019.

REFERENCES

- Selby JV, Beal AC, Frank L. The Patient-Centered Outcomes Research Institute (PCORI) national priorities for research and initial research agenda. *JAMA*. 2012;307(15):1583–1584.
- Perfetto EM, Oehrlein EM, Boutin M, Reid S, Gascho E. Value to whom? The patient voice in the value discussion. Value Health. 2017;20(2):286–291.
- Mason NR, Sox HC, Whitlock EP. A patient-centered approach to comparative
 effectiveness research focused on older adults: lessons from the PatientCentered Outcomes Research Institute. J Am Geriatr Soc. 2019;67(1):21–28.
- Crocker JC, Ricci-Cabello I, Parker A, et al. Impact of patient and public involvement on enrolment and retention in clinical trials: systematic review and meta-analysis. BMJ. 2018;363:k4738.
- Levitan B, Getz K, Eisenstein EL, et al. Assessing the financial value of patient engagement: a quantitative approach from CTTI's patient groups and clinical trials project. *Ther Innov Regul Sci.* 2018;52(2):220–229.
- Gardiner R. The transition from "informed patient" care to "patient informed" care. Stud Health Technol Inform. 2008;137:241–256.
- Hoos A, Anderson J, Boutin M, et al. Partnering With patients in the development and lifecycle of medicines: a call for action. *Ther Innov Regul Sci.* 2015;49(6):929–939.
- 8. Pushparajah DS. Making patient engagement a reality. *Patient*. 2018;11(1): 1–8.
- Anderson M, McCleary KK. From passengers to co-pilots: patient roles expand. Sci Transl Med. 2015;7(291), 2916525-2916525.
- Institute for Patient- and Family-Centered Care. About us. http://www.ipfcc. org/about/index.html. Accessed October 1, 2018.
- Basch E, Geoghegan C, Coons SJ, et al. Patient-reported outcomes in cancer drug development and US regulatory review: perspectives from industry, the Food and Drug Administration, and the patient. JAMA Oncol. 2015;1(3):375– 270
- 12. Furlong P, Bridges JFP, Charnas L, et al. How a patient advocacy group developed the first proposed draft guidance document for industry for submission to the U.S. Food and Drug Administration. *Orphanet J Rare Dis.* 2015;10:82.
- Perfetto EM, Burke L, Oehrlein EM, Epstein RS. Patient-focused drug development: a new direction for collaboration. *Med Care*. 2015;53(1): 9–17.
- 14. U.S. Department of Health and Human Services Food and Drug Administration. Guidance for industry: patient-reported outcome measures: use in

- medical product development to support label claims. https://www.fda.gov/downloads/drugs/guidances/ucm193282.pdf. Accessed April 23, 2018.
- European Medicines Agency. Partners & networks patients and consumers. http://www.ema.europa.eu/ema/index.jsp?curl=pages/partners_and_networks/general/general_content_000317.jsp&mid=WC0b01ac0580035 00c. Accessed May 3, 2018.
- Food and Drug Administration Center for Drug Evaluation and Research. FDA
 patient-focused drug development guidance series for enhancing the
 incorporation of the patient's voice in medical product development and
 regulatory decision making. https://www.fda.gov/drugs/developmentapproval
 process/ucm610279.htm. Accessed October 22, 2018.
- Innovative Medicines Initiative. Get involved patients. IMI Innovative Medicines Initiative. https://www.imi.europa.eu/get-involved/patients. Accessed May 3, 2018.
- Frank L, Basch E, Selby JV. The PCORI perspective on patient-centered outcomes research. JAMA. 2014;312(15):1513–1514.
- Carman KL, Dardess P, Maurer M, et al. Patient and family engagement: a framework for understanding the elements and developing interventions and policies. Health Aff (Millwood). 2013;32(2):223–231.
- What is EUPATI? EUPATI. https://www.eupati.eu/what-is-eupati/. Accessed July 23, 2018.
- EURORDIS. What we do. https://www.eurordis.org/what-we-do. Accessed October 1, 2018.
- Troya MI, Bartlam B, Chew-Graham CA. Involving the public in health research in Latin America: making the case for mental health. Rev Panam Salud Pública. 2018;42.
- Yang JC-L. Patient involvement in HTA in Taiwan. Presented at the ISPOR 22nd Annual International meeting. https://www.ispor.org/docs/default-source/presentations/1071.pdf?sfvrsn=35927285_1. Accessed December 9, 2019.
- **24.** Concannon TW, Meissner P, Grunbaum JA, et al. A new taxonomy for stakeholder engagement in patient-centered outcomes research. *J Gen Intern Med.* 2012;27(8):985–991.
- Goodyear-Smith F. Collective enquiry and reflective action in research: towards a clarification of the terminology. Fam Pract. 2017;34(3):268–271.
- FasterCures. Reports—expanding the science of patient input: the power of language. https://www.fastercures.org/reports/view/59. Accessed October 1, 2018.
- Harrington R, Hareendran A, Scott A, et al. Patient engagement in outcomes research: a snapshot from the ISPOR community. Value Outcomes Spotlight. 2016;2(5).
- Berger ZD, Brito JP, Ospina NS, et al. Patient centred diagnosis: sharing diagnostic decisions with patients in clinical practice. BMJ. 2017;359: j4218.
- Haywood K, Lyddiatt A, Brace-McDonnell SJ, Staniszewska S, Salek S. Establishing the values for patient engagement (PE) in health-related quality of life (HRQoL) research: an international, multiple-stakeholder perspective. Qual Life Res. 2017;26(6):1393–1404.
- Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. Qual Health Res. 2005;15(9):1277–1288.
- **31.** Bradley EH, Curry LA, Devers KJ. Qualitative data analysis for health services research: developing taxonomy, themes, and theory. *Health Serv Res.* 2007;42(4):1758–1772.
- Corbin J, Strauss A. Basics of Qualitative Research (3rd Ed.): Techniques and Procedures for Developing Grounded Theory. Thousand Oaks, CA: SAGE Publications; 2018.
- Domecq JP, Prutsky G, Elraiyah T, et al. Patient engagement in research: a systematic review. BMC Health Serv Res. 2014;14(1):89.
- Institute of Medicine (US) Committee on Quality of Health Care in America. Crossing the Quality Chasm: A New Health System for the 21st Century. Washington, DC: National Academies Press; 2001.
- Stewart M. Patient-Centered Medicine: Transforming the Clinical Method. Oxford, UK: Radcliffe Publishing; 2003.
- Patient-Centered Outcomes Research Institute. Engagement rubric for applicants. https://www.pcori.org/sites/default/files/Engagement-Rubric.pdf.
 Accessed March 1, 2018.
- Wicks P, Hixson J. The Patient engagement pill: lessons from epilepsy. https://www.healthaffairs.org/do/10.1377/hblog20130207.027850/full/. Accessed March 1, 2018.
- European Patients Forum. Value+ Toolkit. http://www.eu-patient.eu/ whatwedo/projects/valueplus. Accessed March 1, 2018.
- U.S. Food and Drug Administration. Plan for issuance of patient-focused drug development guidance under 21st Century Cures Act Title III Section 3002. https://www.fda.gov/downloads/ForIndustry/UserFees/PrescriptionDrug UserFee/UCM563618.pdf. Accessed March 1, 2018.
- Patient-Centered Outcomes Research Institute. what we mean by engagement. https://www.pcori.org/engagement/what-we-mean-engagement. Accessed February 28, 2018.
- 41. Hung A, Baas C, Bekelman J, Fitz-Randolph M, Mullins CD. Patient and stakeholder engagement in designing pragmatic clinical trials. In: Birnbaum HG, Greenberg PE, eds. *Decision Making in a World of Comparative Effectiveness Research*. Singapore, Singapore: Adis; 2017:137–152.
- National Health Council. The patient voice in value: The National Health Council patient-centered value model rubric. http://www.

nationalhealthcouncil.org/sites/default/files/Value-Rubric.pdf. Accessed

- Buckman-Garner S, Gill D, Larkindale J, Anderson M. Collaboration 2.0: exploring the future of research-by-collaboration in medical R&D. Presented at the FasterCures webinar. https://www.fastercures.org/events/webinars/. Accessed October 22, 2018.
- 44. FasterCures. Frameworks & principles. http://www.fastercures.org/programs/patients-count/patient-engagement-library/frameworks-and-principles/. Accessed March 26, 2018.
- National Health Council. Genetic Alliance. Patient-focused drug development

 recommended language for use in guidance document development.
 https://nationalhealthcouncil.org/wp-content/uploads/2019/12/NHC-GA Feb2017.pdf. Accessed February 28, 2018.
- Drug Information Association. DIA considerations guide to implementing patient-centric initiatives in health care product development. http://engage. diaglobal.org/PatientEngagementConsiderationsGuide.html. Accessed May 7, 2018.
- Patient-Centered Outcomes Research Institute. PCORI methodology standards. https://www.pcori.org/research-results/about-our-research/researchmethodology/pcori-methodology-standards. Accessed May 7, 2018.
- 48. Heath N, Castillo WC, Majid T, et al. Patient engagement in observational pharmacoepidemiology research: where are we, where do we need to be, and what are the steps for getting there? *Pharmacoepidemiol Drug Saf.* 2016;25(Suppl 3):47.
- Forsythe LP, Frank LB, Workman TA, Hilliard T, Harwell D, Fayish L. Patient, caregiver and clinician views on engagement in comparative effectiveness research. J Comp Eff Res. 2017;6(3):231–244.
- Law E, Harrington R, Alexander GC, Saha S, Oehrlein E, Perfetto EM. Increasing uptake of comparative effectiveness and patient-centered outcomes research among stakeholders: insights from conference discussion. J Comp Eff Res. 2018;7(2):181–191.
- Abrahams E, Foti M, Kean MA. Accelerating the delivery of patient-centered, high-quality cancer care. Clin Cancer Res. 2015;21(10):2263–2267.
- 52. Knier S, Stichler JF, Ferber L, Catterall K. Patients' perceptions of the quality of discharge teaching and readiness for discharge. *Rehabil Nurs*. 2015;40(1):30–30
- 53. Van Royen P, Beyer M, Chevallier P, et al. The research agenda for general practice/family medicine and primary health care in Europe. Part 3. Results: person centred care, comprehensive and holistic approach. Eur J Gen Pract. 2010;16(2):113–119.
- Li J, Porock D. Resident outcomes of person-centered care in long-term care: a narrative review of interventional research. *Int J Nurs Stud.* 2014;51(10):1395–1415.
- Holzmueller CG, Wu AW, Pronovost PJ. A framework for encouraging patient engagement in medical decision making. J Patient Saf. 2012;8(4):161– 164.
- Balbale SN, Morris MA, LaVela SL. Using photovoice to explore patient perceptions of patient-centered care in the Veterans Affairs Health Care System. Patient. 2014;7(2):187–195.
- 57. Lindstrom I, Gaston-Johansson F, Danielson E. Documentation of patients' participation in care at the end of life. *Nurs Ethics*. 2006;13(4):394–403.
- Govindarajan P, Larkin GL, Rhodes KV, et al. Patient-centered integrated networks of emergency care: consensus-based recommendations and future research priorities. Acad Emerg Med. 2010;17(12):1322–1329.
- Redfern J, Christopher M, Wolfe CDA. Risk management after stroke: the limits of a patient-centred approach. *Health Risk Soc.* 2007;8(2):123–141.
- Greenhalgh T, Snow R, Ryan S, Rees S, Salisbury H. Six "biases" against patients and carers in evidence-based medicine. BMC Med. 2015;13, 200-015-0437-x.
- 61. Hasse GL. Patient-centered care in adult trauma intensive care unit. *J Trauma Nurs*. 2013;20(3):163–165.
- 62. Nazario RJ. Medical humanities as tools for the teaching of patient-centered care. *J Hosp Med*. 2009;4(8):512–514.
- 63. Gagliardi AR, Lemieux-Charles L, Brown AD, Sullivan T, Goel V. Barriers to patient involvement in health service planning and evaluation: an exploratory study. *Patient Educ Couns*. 2008;70(2):234–241.
- **64.** Robinson JH, Callister LC, Berry JA, Dearing KA. Patient-centered care and adherence: definitions and applications to improve outcomes. *J Am Acad Nurse Pract*. 2008;20(12):600–607.
- **65.** Van Berckelaer A, DiRocco D, Ferguson M, Gray P, Marcus N, Day S. Building a patient-centered medical home: obtaining the patient's voice. *J Am Board Fam Med.* 2012;25(2):192–198.
- Cegala DJ, Post DM. The impact of patients' participation on physicians' patient-centered communication. *Patient Educ Couns*. 2009;77(2):202–208.
- 67. Gerrits T. The ambiguity of patient-centred practices: the case of a Dutch fertility clinic. *Anthropol Med*. 2014;21(2):125–135.
- **68.** Lawrence M, Kinn S. Defining and measuring patient-centred care: an example from a mixed-methods systematic review of the stroke literature. *Health Expect.* 2012;15(3):295–326.
- Koberich S, Farin E. A systematic review of instruments measuring patients' perceptions of patient-centred nursing care. *Nurs Inq.* 2015;22(2): 106–120.
- Lord L, Gale N. Subjective experience or objective process: understanding the gap between values and practice for involving patients in designing patientcentred care. J Health Organ Manag. 2014;28(6):714–730.

- Tham JS, Letendre MC. Health care decision making: cross-cultural analysis of the shift from the autonomous to the relational staff. New Bioeth. 2014;20(2):174–185.
- Grenness C, Hickson L, Laplante-Levesque A, Davidson B. Patient-centred care: a review for rehabilitative audiologists. *Int J Audiol*. 2014;53(Suppl 1):S60–S67.
- 73. van de Bovenkamp HM, Trappenburg MJ. Reconsidering patient participation in guideline development. *Health Care Anal.* 2009;17(3):198–216.
- Lavallee DC, Wicks P, Alfonso Cristancho R, Mullins CD. Stakeholder engagement in patient-centered outcomes research: high-touch or hightech? Expert Rev Pharmacoecon Outcomes Res. 2014;14(3):335–344.
- Beaver K, Craven O, Witham G, et al. Patient participation in decision making: views of health professionals caring for people with colorectal cancer. J Clin Nurs. 2007;16(4):725–733.
- Dancet EA, Ameye L, Sermeus W, et al. The ENDOCARE questionnaire (ECQ): a valid and reliable instrument to measure the patient-centeredness of endometriosis care in Europe. *Hum Reprod Oxf Engl*. 2011;26(11):2988–2999.
- Luxford K, Safran DG, Delbanco T. Promoting patient-centered care: a qualitative study of facilitators and barriers in healthcare organizations with a reputation for improving the patient experience. *Int J Qual Health Care*. 2011;23(5):510–515.
- Masi AT, Vincent A. A historical and clinical perspective endorsing personcentered management of fibromyalgia syndrome. Curr Rheumatol Rev. 2015;11(2):86–95.
- Norfolk T, Birdi K, Walsh D. The role of empathy in establishing rapport in the consultation: a new model. *Med Educ*. 2007;41(7):690–697.
- 80. Prey JE, Woollen J, Wilcox L, et al. Patient engagement in the inpatient setting: a systematic review. J Am Med Inform Assoc. 2014;21(4):742–750.
- Macdonald MT, Heilemann MV, MacKinnon NJ, et al. Confirming delivery: understanding the role of the hospitalized patient in medication administration safety. Qual Health Res. 2014;24(4):536–550.
- Mathers SA, Chesson RA, Proctor JM, McKenzie GA, Robertson E. The use of patient-centered outcome measures in radiology: a systematic review. *Acad Radiol*. 2006;13(11):1394–1404.
- 83. McMurray A, Chaboyer W, Wallis M, Johnson J, Gehrke T. Patients' perspectives of bedside nursing handover. *Coll R Coll Nurs Aust*. 2011;18(1):19–26.
- Constand MK, MacDermid JC, Dal Bello-Haas V, Law M. Scoping review of patient-centered care approaches in healthcare. BMC Health Serv Res. 2014;14, 271-6963-14-271.
- Dwamena F, Holmes-Rovner M, Gaulden CM, et al. Interventions for providers to promote a patient-centred approach in clinical consultations. Cochrane Database Syst Rev. 2012;12:CD003267.
- **86.** Lacy NL, Backer EL. Evidence-based and patient-centered care: results from an STFM group project. *Fam Med.* 2008;40(6):417–422.
- Fredericks S, Lapum J, Schwind J, Beanlands H, Romaniuk D, McCay E. Discussion of patient-centered care in health care organizations. *Qual Manag Health Care*. 2012;21(3):127–134.
- Robb G, Seddon M. Effective Practice Informatics and Quality (EPIQ). Quality improvement in New Zealand healthcare. Part 6: keeping the patient front and centre to improve healthcare quality. N Z Med J. 2006;119(1242):U2174.
- 89. Goodwin P. Partnership in action. Nurs Times. 1998;94(29):36.
- Wuyts WA, Peccatori FA, Russell AM. Patient-centred management in idiopathic pulmonary fibrosis: similar themes in three communication models. Eur Respir Rev. 2014;23(132):231–238.
- 91. Marshall A, Kitson A, Zeitz K. Patients' views of patient-centred care: a phenomenological case study in one surgical unit. *J Adv Nurs*. 2012;68(12):2664–2673.
- Kitson A, Marshall A, Bassett K, Zeitz K. What are the core elements of patient-centred care? A narrative review and synthesis of the literature from health policy, medicine and nursing. J Adv Nurs. 2013;69(1):4–15.
- Macdonald M. Patient safety: examining the adequacy of the 5 rights of medication administration. Clin Nurse Spec CNS. 2010;24(4):196–201.
- 94. Groene O, Lombarts MJ, Klazinga N, Alonso J, Thompson A, Sunol R. Is patient-centredness in European hospitals related to existing quality improvement strategies? Analysis of a cross-sectional survey (MARQuIS study). Qual Saf Health Care. 2009;18(Suppl 1):i44–i50.
- Rathert C, Williams ES, McCaughey D, Ishqaidef G. Patient perceptions of patient-centred care: empirical test of a theoretical model. *Health Expect*. 2015;18(2):199–209.
- Tzeng HM, Yin CY. Patient engagement in hospital fall prevention. Nurs Econ. 2015;33(6):326–334.
- Islam KM, Opoku ST, Apenteng BA, et al. Engaging patients and caregivers in patient-centered outcomes research on advanced stage lung cancer: insights from patients, caregivers, and providers. J Cancer Educ. 2014;29(4):796–801.
- **98.** Uijen AA, Schers HJ, Schellevis FG, van den Bosch WJ. How unique is continuity of care? A review of continuity and related concepts. *Fam Pract*. 2012;29(3):264–271.
- 99. Barello S, Graffigna G. Engaging patients to recover life projectuality: an Italian cross-disease framework. *Qual Life Res.* 2015;24(5):1087–1096.
- Beal A, Hernandez S, Doty M. Latino access to the patient-centered medical home. J Gen Intern Med. 2009;24(Suppl 3):514–520.
- 101. Berger Z, Flickinger TE, Pfoh E, Martinez KA, Dy SM. Promoting engagement by patients and families to reduce adverse events in acute care settings: a systematic review. BMJ Qual Saf. 2014;23(7):548–555.

- **102.** Blanton K. The Patient engagement prescription. *N C Med J.* 2015;76(3):157–
- Brennan PF, Valdez R, Alexander G, et al. Patient-centered care, collaboration, communication, and coordination: a report from AMIA's 2013 Policy Meeting. J Am Med Inform Assoc. 2015;22(e1):e2–e6.
- **104.** Bright FA, Kayes NM, Worrall L, McPherson KM. A conceptual review of engagement in healthcare and rehabilitation. *Disabil Rehabil*. 2015;37(8):643–654.
- Collier MS. Culture change to promote patient engagement: it isn't how... it's who. N. C. Med. J. 2015;76(3):168–170.
- **106.** Collins SA, Gazarian P, Stade D, et al. Clinical workflow observations to identify opportunities for nurse, physicians and patients to share a patient-centered plan of care. *AMIA Annu Symp Proc.* 2014;2014:414–423.
- Cunningham P. Patient engagement during medical visits and smoking cessation counseling. JAMA Intern Med. 2014;174(8):1291–1298.
- Dent M, Pahor M. Patient involvement in Europe–a comparative framework. *J Health Organ Manag*. 2015;29(5):546–555.
- Esmail L, Moore E, Rein A. Evaluating patient and stakeholder engagement in research: moving from theory to practice. *J Comp Eff Res.* 2015;4(2): 133–145.
- Forsythe LP, Szydlowski V, Murad MH, et al. A systematic review of approaches for engaging patients for research on rare diseases. J Gen Intern Med. 2014;29(Suppl 3):S788–S800.
- French J, Purificacion S, Brown E, et al. Each cancer journey begins with one shared step: patient engagement and radiation therapy. *Int J Radiat Oncol Biol Phys.* 2015;93(3):E502.
- Grande SW, Faber MJ, Durand MA, Thompson R, Elwyn G. A classification model of patient engagement methods and assessment of their feasibility in real-world settings. *Patient Educ Couns*. 2014;95(2):281–287.
- 113. Haywood K, Brett J, Salek S, et al. Patient and public engagement in health-related quality of life and patient-reported outcomes research: what is important and why should we care? Findings from the first ISOQOL patient engagement symposium. *Qual Life Res.* 2015;24(5):1069–1076.
- Hibbard JH, Greene J. What the evidence shows about patient activation: better health outcomes and care experiences; fewer data on costs. *Health Aff* (Millwood). 2013;32(2):207–214.
- 115. Irizarry T, DeVito Dabbs A, Curran CR. Patient portals and patient engagement: a state of the science review. J Med Internet Res. 2015;17(6):e148.
- Kates N, Hutchison B, O'Brien P, Fraser B, Wheeler S, Chapman C. Framework for advancing improvement in primary care. *Healthc Pap.* 2012;12(2):8–21.
- Kovacs Burns K, Bellows M, Eigenseher C, Gallivan J. "Practical" resources to support patient and family engagement in healthcare decisions: a scoping review. BMC Health Serv Res. 2014;14:175.
- Marlett N, Shklarov S, Marshall D, Santana MJ, Wasylak T. Building new roles and relationships in research: a model of patient engagement research. Qual Life Res. 2015;24(5):1057–1067.
- Nease RF, Glave Frazee S, Zarin L, Miller SB. Choice architecture is a better strategy than engaging patients to spur behavior change. *Health Aff (Mill-wood)*. 2013;32(2):242–249.
- Patel N, Rajasingam D. User engagement in the delivery and design of maternity services. Best Pract Res Obstet Gynaecol. 2013;27(4):597–608.
- 121. Pomey MP, Ghadiri DP, Karazivan P, Fernandez N, Clavel N. Patients as partners: a qualitative study of patients' engagement in their health care. Plos One. 2015;10(4), e0122499.
- 122. Rogers S, Howard C. Patient and community engagement: the approach at university health network. *Healthc Q Tor Ont.* 2014;17(3):70–73.
- 123. Simmons LA, Wolever RQ, Bechard EM, Snyderman R. Patient engagement as a risk factor in personalized health care: a systematic review of the literature on chronic disease. *Genome Med.* 2014;6(2):16.
- 124. Hahn DL, Hoffmann AE, Felzien M, LeMaster JW, Xu J, Fagnan LJ. Tokenism in patient engagement. *Fam Pract*. 2017;34(3):290–295.
- Bélisle-Pipon J-C, Rouleau G, Birko S. Early-career researchers' views on ethical dimensions of patient engagement in research. BMC Med Ethics. 2018;19(1):21.
- 126. Aaen Geest T, Wetzels R, Wensing M, Cohen Castel O, Olesen F, Improve Research Group. Evaluation of an intervention designed to enhance involvement of older patients in their own care. Eur J Gen Pract. 2006;12(1):3–9.
- Baart IL, Abma TA. Patient participation in fundamental psychiatric genomics research: a Dutch case study. Health Expect. 2011;14(3):240–249.
- 128. Bailey S, Boddy K, Briscoe S, Morris C. Involving disabled children and young people as partners in research: a systematic review. *Child Care Health Dev.* 2015;41(4):505–514.
- **129.** Baker A. Patient involvement in a professional body: reflections and commentary. *J Health Organ Manag.* 2007;21(4-5):460–469.
- Bastiaens H, Van Royen P, Pavlic DR, Raposo V, Baker R. Older people's preferences for involvement in their own care: a qualitative study in primary health care in 11 European countries. *Patient Educ Couns*. 2007;68(1):33–42.
- 131. Cleemput I, Christiaens W, Kohn L, Leonard C, Daue F, Denis A. Acceptability and perceived benefits and risks of public and patient involvement in health care policy: a Delphi survey in Belgian stakeholders. *Value Health*. 2015;18(4):477–483.
- de Wit MP, Berlo SE, Aanerud GJ, et al. European League Against Rheumatism recommendations for the inclusion of patient representatives in scientific projects. Ann Rheum Dis. 2011;70(5):722–726.

- Entwistle VA, Watt IS. Patient involvement in treatment decision-making: the case for a broader conceptual framework. *Patient Educ Couns*. 2006;63(3):268–278.
- 134. Forbat L, Hubbard G, Kearney N. Patient and public involvement: models and muddles. *J Clin Nurs*. 2009;18(18):2547–2554.
- Frith J, Bashir AS, Elliott CS, Newton JL. Public and patient research priorities for orthostatic hypotension. *Age Ageing*. 2014;43(6):865–868.
- 136. Gagnon MP, Desmartis M, Gagnon J, et al. Introducing the patient's perspective in hospital health technology assessment (HTA): the views of HTA producers, hospital managers and patients. Health Expect. 2014;17(6):888–900.
- 137. Gagnon MP, Gagnon J, St-Pierre M, et al. Involving patients in HTA activities at local level: a study protocol based on the collaboration between researchers and knowledge users. BMC Health Serv Res. 2012;12:14.
- Groene O, Sunol R, Klazinga NS, et al. Involvement of patients or their representatives in quality management functions in EU hospitals: implementation and impact on patient-centred care strategies. *Int J Qual Health Care*. 2014;26(Suppl 1):81–91.
- Happ MB, Swigart VA, Tate JA, Hoffman LA, Arnold RM. Patient involvement in health-related decisions during prolonged critical illness. Res Nurs Health. 2007;30(4):361–372.
- Haywood K, Marshall S, Fitzpatrick R. Patient participation in the consultation process: a structured review of intervention strategies. *Patient Educ Couns*. 2006;63(1-2):12–23.
- 141. Holzel LP, Kriston L, Harter M. Patient preference for involvement, experienced involvement, decisional conflict, and satisfaction with physician: a structural equation model test. BMC Health Serv Res. 2013;13:231.
- 142. Kasper J, Heesen C, Kopke S, Fulcher G, Geiger F. Patients' and observers' perceptions of involvement differ. Validation study on inter-relating measures for shared decision making. PloS One. 2011;6(10), e26255.
- 143. Martin HM, Navne LE, Lipczak H. Involvement of patients with cancer in patient safety: a qualitative study of current practices, potentials and barriers. *BMJ Qual Saf.* 2013;22(10):836–842.
- 144. McCaffery KJ, Smith SK, Wolf M. The challenge of shared decision making among patients with lower literacy: a framework for research and development. Med Decis Making. 2010;30(1):35–44.
- 145. O'Brien MA, Charles C, Whelan TJ, Ellis PM, Gafni A, Lovrics P. Women's perceptions of their involvement in treatment decision making for early stage breast cancer. Support Care Cancer. 2013;21(6):1717–1723.
- **146.** Petit-Zeman S, Philpots E, Denegri S. "Natural ground" for medical research charities: public and patient involvement in research funding. *J Ambul Care Manage*. 2010;33(3):249–256.
- 147. Rise MB, Solbjor M, Lara MC, Westerlund H, Grimstad H, Steinsbekk A. Same description, different values. How service users and providers define patient and public involvement in health care. Health Expect. 2013;16(3):266–276.
- **148.** Tambuyzer E, Pieters G, Van Audenhove C. Patient involvement in mental health care: one size does not fit all. *Health Expect.* 2014;17(1):138–150.
- 149. Williamson L. Patient and citizen participation in health: the need for improved ethical support. Am J Bioeth. 2014;14(6):4–16.
- **150.** Aasen EM, Kvangarsnes M, Heggen K. Perceptions of patient participation amongst elderly patients with end-stage renal disease in a dialysis unit. *Scand J Caring Sci.* 2012;26(1):61–69.
- 151. Abma TA. Patients as partners in a health research agenda setting: the feasibility of a participatory methodology. Eval Health Prof. 2006;29(4): 424–439.
- **152.** Cegala DJ. An exploration of factors promoting patient participation in primary care medical interviews. *Health Commun.* 2011;26(5): 427–436.
- **153.** Cegala DJ, Chisolm DJ, Nwomeh BC. Further examination of the impact of patient participation on physicians' communication style. *Patient Educ Couns*. 2012;89(1):25–30.
- **154.** Elberse JE, Caron-Flinterman JF, Broerse JE. Patient-expert partnerships in research: how to stimulate inclusion of patient perspectives. *Health Expect*. 2011;14(3):225–239.
- Eldh AC, Ekman I, Ehnfors M. Conditions for patient participation and nonparticipation in health care. Nurs Ethics. 2006;13(5):503–514.
- **156.** Eldh AC, Ekman I, Ehnfors M. Considering patient non-participation in health care. *Health Expect.* 2008;11(3):263–271.
- Eldh AC, Ekman I, Ehnfors M. A comparison of the concept of patient participation and patients' descriptions as related to healthcare definitions. Int J Nurs Terminol Classif. 2010;21(1):21–32.
- Goss C, Mosconi P, Renzi C, Deledda G. Participation of patients and citizens in healthcare decisions in Italy. Z Evidenz Fortbild Qual Gesundhwes. 2011;105(4):277–282.
- 159. Heggland LH, Mikkelsen A, Ogaard T, Hausken K. Measuring patient participation in surgical treatment decision-making from healthcare professionals' perspective. J Clin Nurs. 2014;23(3-4):482–491.
- Heggland LH, Hausken K. Patient participation, decision-makers and information flow in surgical treatment. J Clin Nurs. 2014;23(9-10):1430–1444.
- 161. Helgesen AK, Larsson M, Athlin E. Patient participation in special care units for persons with dementia: A losing principle? Nurs Ethics. 2014;21(1):108–118.
- 162. Kim H, Suh EE, Lee HJ, Yang HK. The effects of patient participation-based dietary intervention on nutritional and functional status for patients

with gastrectomy: a randomized controlled trial. *Cancer Nurs*. 2014;37(2):E10–E20.

- 163. Kvangarsnes M, Torheim H, Hole T, Ohlund LS. Intensive care unit nurses' perceptions of patient participation in the acute phase of chronic obstructive pulmonary disease exacerbation: an interview study. J Adv Nurs. 2013;69(2):425–434.
- 164. Larsson IE, Sahlsten MJ, Sjostrom B, Lindencrona CS, Plos KA. Patient participation in nursing care from a patient perspective: a Grounded Theory study. Scand J Caring Sci. 2007;21(3):313–320.
- Lyttle DJ, Ryan A. Factors influencing older patients' participation in care: a review of the literature. Int J Older People Nurs. 2010;5(4):274–282.
- 166. Nordin C, Gard G, Fjellman-Wiklund A. Being in an exchange process: experiences of patient participation in multimodal pain rehabilitation. *J Rehabil Med.* 2013;45(6):580–586.
- **167.** Thorarinsdottir K, Kristjansson K. Patients' perspectives on person-centred participation in healthcare: a framework analysis. *Nurs Ethics*. 2014;21(2):129–147.
- 168. Vahdat S, Hamzehgardeshi L, Hessam S, Hamzehgardeshi Z. Patient involvement in health care decision making: a review. *Iran Red Crescent Med* I. 2014:16(1), e12454.
- 169. van der Weijden T, van Veenendaal H, Timmermans D. Shared decision-making in the Netherlands-current state and future perspectives. *Z Arztl Fortbild Qualitatssich*. 2007;101(4):241–246.
- 170. Aardoom JJ, Dingemans AE, Boogaard LH, Van Furth EF. Internet and patient empowerment in individuals with symptoms of an eating disorder: a crosssectional investigation of a pro-recovery focused e-community. Eat Behav. 2014;15(3):350–356.
- Alpay LL, Henkemans OB, Otten W, Rovekamp TA, Dumay AC. E-health applications and services for patient empowerment: directions for best practices in The Netherlands. *Telemed J E Health*. 2010;16(7):787–791.
- 172. Ammenwerth E, Schnell-Inderst P, Hoerbst A. Patient empowerment by electronic health records: first results of a systematic review on the benefit of patient portals. *Stud Health Technol Inform.* 2011;165:63–67.
- 173. Aujoulat I, d'Hoore W, Deccache A. Patient empowerment in theory and practice: polysemy or cacophony? *Patient Educ Couns*. 2007;66(1):13–20.
- Barr PJ, Scholl I, Bravo P, Faber MJ, Elwyn G, McAllister M. Assessment of patient empowerment–a systematic review of measures. *PloS One*. 2015;10(5), e0126553.
- Bridges JF, Jones C. Patient-based health technology assessment: a vision of the future. Int J Technol Assess Health Care. 2007;23(1):30–35.
- Christensen M, Hewitt-Taylor J. Empowerment in nursing: paternalism or maternalism? Br J Nurs Mark Allen Publ. 2006;15(13):695–699.
- 177. Christensen M, Hewitt-Taylor J. Patient empowerment: does it still occur in the ICU? *Intensive Crit Care Nurs*. 2007;23(3):156–161.
- **178.** de-Graft A. Barriers to patient empowerment in sub-Saharan Africa: a social psychological evaluation. *Diabetes Res Clin Pract.* 2014;103. S7-S7.
- Holmstrom I, Roing M. The relation between patient-centeredness and patient empowerment: a discussion on concepts. *Patient Educ Couns*. 2010;79(2):167–172.
- Koch S. İmproving quality of life through eHealth the patient perspective. *Stud Health Technol Inform.* 2012;180:25–29.
- 181. Lejbkowicz I, Caspi O, Miller A. Participatory medicine and patient empowerment towards personalized healthcare in multiple sclerosis. Expert Rev Neurother. 2012;12(3):343–352.
- **182.** Loukanova S, Molnar R, Bridges JF. Promoting patient empowerment in the healthcare system: highlighting the need for patient-centered drug policy. *Expert Rev Pharmacoecon Outcomes Res.* 2007;7(3):281–289.
- 183. Macq J, Torfoss T, Getahun H. Patient empowerment in tuberculosis control: reflecting on past documented experiences. Trop Med Int Health. 2007;12(7):873–885.
- 184. McCarley P. Patient empowerment and motivational interviewing: engaging patients to self-manage their own care. Nephrol Nurs J. 2009;36(4):409–413.
- 185. Schulz PJ, Nakamoto K. Health literacy and patient empowerment in health communication: the importance of separating conjoined twins. *Patient Educ Couns*. 2013;90(1):4–11.
- 186. Te Boveldt N, Vernooij-Dassen M, Leppink I, Samwel H, Vissers K, Engels Y. Patient empowerment in cancer pain management: an integrative literature review. *Psychooncology*, 2014;23(11):1203–1211.
- 187. Teunissen T, Visse M, de Boer P, Abma TA. Patient issues in health research and quality of care: an inventory and data synthesis. *Health Expect*. 2013;16(4):308–322.
- 188. Wang J, Thombs BD, Schmid MR. The Swiss Health Literacy Survey: development and psychometric properties of a multidimensional instrument to assess competencies for health. *Health Expect*. 2014;17(3):396–417.
- Beach MC, Keruly J, Moore RD. Is the quality of the patient-provider relationship associated with better adherence and health outcomes for patients with HIV? J Gen Intern Med. 2006;21(6):661–665.
- Bernabeo E, Holmboe ES. Patients, providers, and systems need to acquire a specific set of competencies to achieve truly patient-centered care. Health Aff (Millwood). 2013;32(2):250–258.

- 191. Bjorkman IK, Bernsten CB, Sanner MA. Care ideologies reflected in 4 conceptions of pharmaceutical care. Res Soc Adm Pharm. 2008;4(4):332– 342
- Hauer KE, Boscardin C, Gesundheit N, Nevins A, Srinivasan M, Fernandez A. Impact of student ethnicity and patient-centredness on communication skills performance. Med Educ. 2010;44(7):653–661.
- Hommel I, van Gurp PJ, Tack CJ, et al. Perioperative diabetes care: room for improving the person centredness. *Diabet Med.* 2015;32(4): 561–568.
- 194. Ishikawa H, Hashimoto H, Kiuchi T. The evolving concept of "patient-centeredness" in patient-physician communication research. Soc Sci Med. 2013;96:147–153.
- 195. Kofahl C, Trojan A, Knesebeck O, Nickel S. Self-help friendliness: a German approach for strengthening the cooperation between self-help groups and health care professionals. Soc Sci Med. 2014;123:217–225.
- 196. Levack WM, Dean SG, Siegert RJ, McPherson KM. Navigating patient-centered goal setting in inpatient stroke rehabilitation: how clinicians control the process to meet perceived professional responsibilities. *Patient Educ Couns*. 2011;85(2):206–213.
- Scholl I, Zill JM, Harter M, Dirmaier J. An integrative model of patientcenteredness - a systematic review and concept analysis. PloS One. 2014;9(9), e107828.
- 198. Wiig S, Storm M, Aase K, et al. Investigating the use of patient involvement and patient experience in quality improvement in Norway: rhetoric or reality? BMC Health Serv Res. 2013;13(206).
- 199. Zill JM, Scholl I, Harter M, Dirmaier J. Which dimensions of patient-centeredness matter? results of a web-based expert Delphi survey. PloS One. 2015;10(11), e0141978.
- 200. Wijma AJ, Bletterman AN, Clark JR, et al. Patient-centeredness in physiotherapy: what does it entail? A systematic review of qualitative studies. *Physiother Theory Pract*. 2017;33(11):825–840.
- 201. Bolster D, Manias E. Person-centred interactions between nurses and patients during medication activities in an acute hospital setting: qualitative observation and interview study. *Int J Nurs Stud.* 2010;47(2):154–165.
- 202. American Geriatrics Society Expert Panel on Person-Centered Care. Person-Centered care: a definition and essential elements. J Am Geriatr Soc. 2016;64(1):15–18.
- 203. Kirkendall AM, Waldrop D, Moone RP. Caring for people with intellectual disabilities and life-limiting illness: merging person-centered planning and patient-centered, family-focused care. J Soc Work End Life Palliat Care. 2012;8(2):135–150.
- Lindberg J, Kreuter M, Person LO, Taft C. Patient participation in rehabilitation questionnaire (PPRQ)-development and psychometric evaluation. Spinal Cord. 2013;51(11):838–842.
- **205.** Olander EK, Berg M, McCourt C, Carlstrom E, Dencker A. Person-centred care in interventions to limit weight gain in pregnant women with obesity a systematic review. *BMC Pregnancy Childbirth*. 2015;15(50).
- 206. Rosvik J, Brooker D, Mjorud M, Kirkevold O. What is person-centred care in dementia? Clinical reviews into practice: the development of the VIPS practice model. Rev Clin Gerontol. 2013;23(2):155–163.
- Slater L. Person-centredness: a concept analysis. Contemp Nurse. 2006;23(1):135–144.
- Westphal EC, Alkema G, Seidel R, Chernof B. How to get better care with lower costs? See the person, not the patient. J Am Geriatr Soc. 2016;64(1):19– 21
- 209. Nasser M, van Weel C, van Binsbergen JJ, van de Laar FA. Generalizability of systematic reviews of the effectiveness of health care interventions to primary health care: concepts, methods and future research. Fam Pract. 2012;29(Suppl 1):i94–i103.
- Bellows J, Young S, Chase A. Person-focused care at Kaiser Permanente. Perm J. 2014;18(1):90–91.
- Irwin RS, Richardson ND. Patient-focused care: using the right tools. Chest. 2006;130(1 Suppl):73S-82S.
- 212. Lenert L. Transforming healthcare through patient empowerment. Stud Health Technol Inform. 2010;153:159–175.
- Robbins DA, Curro FA, Fox CH. Defining patient-centricity: opportunities, challenges, and implications for clinical care and research. *Ther Innov Regul Sci.* 2003;47(3):349–355.
- **214.** Frank L, Forsythe L, Ellis L, et al. Conceptual and practical foundations of patient engagement in research at the patient-centered outcomes research institute. *Qual Life Res.* 2015;24(5):1033–1041.
- 215. Happell B, Roper C. Consumer participation in mental health research: articulating a model to guide practice. *Australas Psychiatry*. 2007;15(3): 237–241.
- 216. Barham L. Public and patient involvement at the UK National Institute for Health and Clinical Excellence. *Patient*. 2011;4(1):1–10.
- 217. Duncan B, Ables AZ. Do drug treatment POEMs report data in clinically useful ways? *J Fam Pract*. 2013;62(2):E1–E5.
- 218. Gruman J, Rovner MH, French ME, et al. From patient education to patient engagement: implications for the field of patient education. *Patient Educ Couns*. 2010;78(3):350–356.