

A generalized-means choice model for regularity violations

Khaled Boughanmi – Kamel Jedidi – Rajeev Kohli
Columbia University

THE REGULARITY ASSUMPTION

The probability of choosing an alternative does not increase when other alternatives are added to a choice set

VIOLATION OF REGULARITY

- ▶ Choice complexity
 - ▶ Redelmeier and Shafir (1995), Iyengar and Lepper (2000)

VIOLATION OF REGULARITY

- ▶ Choice complexity
 - ▶ Redelmeier and Shafir (1995), Iyengar and Lepper (2000)
- ▶ Context effects
 - ▶ Huber et. al. (1982), Simonson (1989), Sedikides, Ariely and Olsen (1999), Tenori, Osherson, Hasher and May (2001), Kivetz et. al. (2004), Rooderkerk et. al. (2011)

VIOLATION OF REGULARITY

- ▶ Choice complexity
 - ▶ Redelmeier and Shafir (1995), Iyengar and Lepper (2000)
- ▶ Context effects
 - ▶ Huber et. al. (1982), Simonson (1989), Sedikides, Ariely and Olsen (1999), Tenori, Osherson, Hasher and May (2001), Kivetz et. al. (2004), Rooderkerk et. al. (2011)
- ▶ Menu dependence, herd behavior, chooser dependence, responsibility
 - ▶ Kant (1788), Adam Smith (1790), Banerjee (1992), Sen (1993), Redelmeier and Shafir (1995), Sen (1997)

IS THE GLASS HALF FULL OR HALF EMPTY?



HOW DO CONSUMERS VALUE THINGS?

Different valuation models have been considered:

- ▶ Linear Additive Model : $V = \sum_{j=1}^m \beta_j x_j$

HOW DO CONSUMERS VALUE THINGS?

Different valuation models have been considered:

- ▶ Linear Additive Model : $V = \sum_{j=1}^m \beta_j x_j$
- ▶ “Award” : $V = \max(\beta_j x_j)$

HOW DO CONSUMERS VALUE THINGS?

Different valuation models have been considered:

- ▶ Linear Additive Model : $V = \sum_{j=1}^m \beta_j x_j$
- ▶ “Award” : $V = \max(\beta_j x_j)$
- ▶ “Deny” : $V = \min(\beta_j x_j)$

HOW DO CONSUMERS VALUE THINGS?

Different valuation models have been considered:

- ▶ Linear Additive Model : $V = \sum_{j=1}^m \beta_j x_j$
- ▶ “Award” : $V = \max(\beta_j x_j)$
- ▶ “Deny” : $V = \min(\beta_j x_j)$
- ▶ Prospect theory : V defined
 - (i) on deviations from a reference point x_j^*
 - (ii) concave for gains and convex for losses
 - (iii) steeper for losses than gains

where:

- ▶ $x_j, j = 1, \dots, m$ are product attributes
- ▶ $x_j^*, j = 1, \dots, m$ are attributes references
- ▶ β_j attribute weights

OBJECTIVES

- ▶ Propose a generalized mean value function
- ▶ Show how to use it for modeling choice
- ▶ Show how it captures violation of regularity
- ▶ Illustrate the model on empirical choice data

THE VALUE FUNCTION

- Let the value of attribute j be

$$v_j = e^{\beta_j(x_j - x_j^*)}$$

THE VALUE FUNCTION

- ▶ Let the value of attribute j be

$$v_j = e^{\beta_j(x_j - x_j^*)}$$

- ▶ Then the generalized mean value of an alternative is:

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

where r is a parameter that governs the averaging process

SPECIAL CASES

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

$$r = 1 : \quad V_1 = \frac{1}{m} \sum_{j=1}^m v_j \quad \text{Arithmetic mean}$$

SPECIAL CASES

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

$$r = 1 : \quad V_1 = \frac{1}{m} \sum_{j=1}^m v_j \quad \text{Arithmetic mean}$$

$$r = 0 :$$

SPECIAL CASES

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

$$r = 1 : \quad V_1 = \frac{1}{m} \sum_{j=1}^m v_j \quad \text{Arithmetic mean}$$

$$r = 0 : \quad V_0 = \prod_{j=1}^m v_j^{\frac{1}{m}} \quad \text{Geometric mean}$$

SPECIAL CASES

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

$$r = 1 : \quad V_1 = \frac{1}{m} \sum_{j=1}^m v_j \quad \text{Arithmetic mean}$$

$$r = 0 : \quad V_0 = \prod_{j=1}^m v_j^{\frac{1}{m}} \quad \text{Geometric mean}$$

$$r \rightarrow -\infty : \quad V_{-\infty} = \min(v_j) \quad \text{Minimum}$$

SPECIAL CASES

$$V_r = \left(\frac{1}{m} \sum_{j=1}^m (v_j)^r \right)^{\frac{1}{r}}$$

$$r = 1 : \quad V_1 = \frac{1}{m} \sum_{j=1}^m v_j \quad \text{Arithmetic mean}$$


$$r = 0 : \quad V_0 = \prod_{j=1}^m v_j^{\frac{1}{m}} \quad \text{Geometric mean}$$

$$r \rightarrow -\infty : \quad V_{-\infty} = \min(v_j) \quad \text{Minimum}$$

$$r \rightarrow +\infty : \quad V_{+\infty} = \max(v_j) \quad \text{Maximum}$$


VALUE SURFACE

(c) Geometric mean $r=0$




VALUE SURFACE

(a) Minimum



VALUE SURFACE

(e) Maximum



THE CHOICE MODEL

$$U_r(i) = V_r(i)e^\epsilon$$

- ▶ $i \in C = \{1 \dots n\}$ denotes alternatives
- ▶ V_r is the systematic value
- ▶ $v_{ij} = e^{\beta_j(x_{ij} - x_j^*)}$
- ▶ ϵ is an extreme value error

THE CHOICE MODEL

$$U_r(i) = V_r(i)e^{\epsilon}$$

- ▶ $i \in C = \{1 \dots n\}$ denotes alternatives
- ▶ V_r is the systematic value
- ▶ $v_{ij} = e^{\beta_j(x_{ij} - x_j^*)}$
- ▶ ϵ is an extreme value error

$$\mathbb{P}(i \in C|r) = \frac{V_r(i)}{V_r(1) + \dots + V_r(n)}$$

MNL IS A SPECIAL CASE WHEN $r \rightarrow 0$ (GEOMETRIC MEAN)

$$V_r(i) = \left(\frac{1}{m} \sum_{j=1}^m e^{r\beta_j(x_{ij}-x_j^*)} \right)^{\frac{1}{r}} \xrightarrow{r \rightarrow 0} \prod_{j=1}^m e^{\frac{\beta_j}{m}(x_{ij}-x_j^*)} = e^{\sum_{j=1}^m \frac{\beta_j}{m}(x_{ij}-x_j^*)}$$

MNL IS A SPECIAL CASE WHEN $r \rightarrow 0$ (GEOMETRIC MEAN)



$$V_r(i) = \left(\frac{1}{m} \sum_{j=1}^m e^{r\beta_j(x_{ij}-x_j^*)} \right)^{\frac{1}{r}} \xrightarrow{r \rightarrow 0} \prod_{j=1}^m e^{\frac{\beta_j}{m}(x_{ij}-x_j^*)} = e^{\sum_{j=1}^m \frac{\beta_j}{m}(x_{ij}-x_j^*)}$$

The Logit model is equivalent to a geometric averaging process:


$$\mathbb{P}(i \in C | r = 0) = \frac{e^{\sum_{j=1}^m \tilde{\beta}_j x_{ij}}}{e^{\sum_{j=1}^m \tilde{\beta}_j x_{1j}} + \dots + e^{\sum_{j=1}^m \tilde{\beta}_j x_{nj}}}$$

where $\tilde{\beta}_j \leftarrow \frac{\beta_j}{m}$


ILLUSTRATING VIOLATION OF REGULARITY

 $r_1 = 0$  r_2


CHOICE PROBABILITIES



CHOICE PROBABILITIES




CHOICE PROBABILITIES




EMPIRICAL ILLUSTRATION

- ▶ Data on choice of digital cameras (Roederkerk et. al. (2011)) of 154 subjects
- ▶ Independent variables include picture quality (from 2MP to 6MP) and optical zoom (from $\times 2$ to $\times 10$)
- ▶ Choice sets varying in size from three to five alternatives plus a no-choice option

VIOLATION OF REGULARITY



ACTUAL AND PREDICTED SHARES OF THE NO-CHOICE OPTION



MODEL PERFORMANCE STATISTICS

Model	WAIC	LOO
Multinomial Logit	6553	-3279
Common r	3904	-1963
Varying r	3882	-1956

HYPER PARAMETERS ESTIMATES


Parameter	Label	Multinomial logit	Constant averaging	General averaging
No choice	β_0	10.71 (9.91,11.52)		
Reference points				
Picture quality	PQ^*		3.77 (3.61,3.90)	3.73 (3.58,3.89)
Optical zoom	OZ^*		3.45 (3.24,3.84)	3.45 (3.14,3.76)
Sensitivities				
Megapixels	β_1	1.82 (1.70,4.56)	4.56 (4.02,5.16)	4.42 (4.00,4.92)
Optical zoom	β_2	0.67 (0.62,0.73)	2.51 (2.18,2.89)	2.43 (2.15,2.72)
Averaging rules				
Invariant	r		-0.43 (-0.59,-0.31)	
3 alternatives	r_3			-0.51 (-0.71,-0.39)
4 alternatives	r_4			-0.69 (-1.48,-0.37)
5 alternatives	r_5			-3.62 (-6.66,-0.88)

Notes. Significant coefficients are highlighted in boldface ($p \leq 0.05$).


The 95% posterior intervals for the parameters are shown in parentheses.

FITTED VALUE SURFACES


(a) General averaging: 3 options




(b) General averaging: 4 options



(c) General averaging: 5 options





FITTED VALUE FUNCTION




Optical zoom is fixed at its reference point


IMPLICATIONS




IMPLICATIONS: MULTINOMIAL LOGIT




IMPLICATIONS: MULTINOMIAL LOGIT



IMPLICATIONS: GENERALIZED-MEANS MODEL



IMPLICATIONS: GENERALIZED-MEANS MODEL



CONCLUSION

- ▶ Proposed a generalized mean value function
- ▶ Showed how to use it for modeling choice
- ▶ Showed how it captures violation of regularity
- ▶ Illustrated the model on empirical choice data

IS THE GLASS HALF FULL OR HALF EMPTY?



IS THE GLASS HALF FULL OR HALF EMPTY?



- ▶ Optimists: Half-full
- ▶ Pessimists: Half-Empty
- ▶ $V_r(\text{Glass}) = \left(\frac{v_{\text{Full}}^r + v_{\text{Empty}}^r}{2} \right)^{\frac{1}{r}}$

EXPECTED IN-SAMPLE AND OUT-OF-SAMPLE HIT RATES BY CHOICE-SET SIZE

Model	3 options		4 options		5 options	
	In	Out	In	Out	In	Out
Multinomial Logit	0.46	0.45	0.39	0.38	0.38	0.38
Common r	0.71	0.67	0.67	0.62	0.65	0.62
Varying r	0.71	0.67	0.68	0.63	0.69	0.63