

deeplearning.ai

## One hidden layer Neural Network

## Gradient descent for neural networks

## Gradient descent for neural networks

Porometrs: 
$$(n^{(1)}, n^{(2)})$$
  $(n^{(2)}, n^{(2)})$   $(n^{(2)}, n^{(2)}$ 

## Formulas for computing derivatives

Formal bobadquin;  

$$S_{CIJ} = P_{CIJ}(S_{CIJ}) = e(S_{CIJ})$$

$$S_{LSJ} = P_{LSJ}(S_{LSJ}) = e(S_{LSJ})$$

$$S_{LSJ} = P_{LSJ}(S_{LSJ}) = e(S_{LSJ})$$

$$S_{LSJ} = P_{LSJ}(S_{LSJ}) = e(S_{LSJ})$$

Back propagation:

$$\begin{aligned}
&\mathcal{L}^{[2]} = \mathcal{A}^{[2]} - \mathcal{L} \\
&\mathcal{L}^{[1]} = \mathcal{A}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{$$

**Andrew Ng**