

Chapter 9

Introduction to Graphs

Discrete Structures for Computing

TÀI LIỆU SƯU TẬP

Huynh Tuong Nguyen, Tran Tuan Anh, Nguyen Ngoc Le
Faculty of Computer Science and Engineering
University of Technology - VNUHCM
{htnguyen;trtanh}@hcmut.edu.vn

Introduction to Graphs

Huynh Tuong Nguyen,
Tran Tuan Anh, Nguyen
Ngoc Le



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs
and Graph
Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Contents

1 Graph definitions

Terminology
Special Graphs

2 Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

3 Exercise

Graph
Bipartite graph
Isomorphism



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Course outcomes

Course learning outcomes	
L.O.1	Understanding of logic and discrete structures L.O.1.1 – Describe definition of propositional and predicate logic L.O.1.2 – Define basic discrete structures: set, mapping, graphs
L.O.2	Represent and model practical problems with discrete structures L.O.2.1 – Logically describe some problems arising in Computing L.O.2.2 – Use proving methods: direct, contrapositive, induction L.O.2.3 – Explain problem modeling using discrete structures
L.O.3	Understanding of basic probability and random variables L.O.3.1 – Define basic probability theory L.O.3.2 – Explain discrete random variables
L.O.4	Compute quantities of discrete structures and probabilities L.O.4.1 – Operate (compute/ optimize) on discrete structures L.O.4.2 – Compute probabilities of various events, conditional ones, Bayes theorem



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Motivations

The need of the graph

- Representation/Storing
- Searching/sorting
- Optimization

Its applications

- Electric circuit/board
- Chemical structure
- Networking
- Map, geometry, ...

- Graph theory is useful for analysing “things that are connected to other things”.
- Some difficult problems become easy when represented using a graph.



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Graph


just only one type : or un or directed

Definition

A graph (đồ thị) G is a pair of (V, E) , which are:


- V – nonempty set of **vertices** (nodes) (đỉnh)
- E – set of **edges** (cạnh)

A graph captures abstract relationships between vertices.



Undirected graph

$$1 \rightarrow 2 = 2 \rightarrow 1$$



Directed graph

have direct to connect ele

$$1 \rightarrow 2 \neq 2 \rightarrow 1$$



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Undirected Graph (Đồ thị vô hướng)

Definition (Simple graph (đơn đồ thị))

- Each edge connects two different vertices, and
- No two edges connect the same pair of vertices

An edge between two vertices u and v is denoted as $\{u, v\}$



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Undirected Graph

Definition (Multigraph (đa đồ thị))

Graphs that may have multiple edges connecting the same vertices.

An unordered pair of vertices $\{u, v\}$ are called **multiplicity m** (bội m) if it has m different edges between.



BACHKHOACNCP.COM



Contents

Graph definitions

- Terminology
- Special Graphs

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise


- Graph
- Bipartite graph
- Isomorphism

Undirected Graph

Definition (Pseudograph (giả đồ thị))

Are multigraphs that have

- **loops** (*khuyên*)— edges that connect a vertex to itself



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Directed Graph

Definition (Directed Graph (đồ thị có hướng))

A directed graph G is a pair of (V, E) , in which:

- V – nonempty set of vertices
- E – set of directed edges (*cạnh có hướng*, arcs)

A directed edge **start** at u and **end** at v is denoted as (u, v) .



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Terminologies For Undirected Graph

Neighborhood

In an undirected graph $G = (V, E)$,

- two vertices u and $v \in V$ are called **adjacent** (*liền kề*) if they are **end-points** (*điểm đầu mút*) of edge $e \in E$, and
- e is **incident with** (*cạnh liên thuộc*) u and v
- e is said to **connect** (*cạnh nối*) u and v ;

The degree of a vertex

The **degree of a vertex** (*bậc của một đỉnh*), denoted by $\deg(v)$ is the **number of edges incident with it, except that a loop contributes twice to the degree of that vertex.**

- **isolated** vertex (*đỉnh cô lập*): vertex of degree **0**
- **pendant** vertex (*đỉnh treo*): vertex of degree **1**



loop = twice \Rightarrow caculate the edges in v



Example

Example

What are the degrees and neighborhoods of the vertices in these graphs?



Solution

In G , $\deg(a) = 2$, $\deg(b) = \deg(c) = \deg(f) = 4$, $\deg(d) = 1$, ...

Neighborhoods of these vertices are

$$N(a) = \{b, f\}, N(b) = \{a, c, f, e\}, \dots$$

In H , $\deg(a) = 4$, $\deg(b) = \deg(e) = 6$, $\deg(c) = 1$, ...

Neighborhoods of these vertices are

$$N(a) = \{b, d, e\}, N(b) = \{a, b, c, d, e\}, \dots$$



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Basic Theorems

Theorem (The Handshaking Theorem)

Let $G = (V, E)$ be an undirected graph with m edges. Then

$$2m = \sum_{v \in V} \deg(v)$$

(Note that this applies even if multiple edges and loops are present.)

Theorem

An undirected graph has an even number of odd-degree vertices.



Prove that ...

hw

...

If the number of vertices in an undirected graph is an odd number, then there exists an even-degree vertex.

...

If the number of vertices in an undirected graph is an odd number, then the number of vertices with even degree is odd.

...

If the number of vertices in an undirected graph is an even number, then the number of vertices with even degree is even.



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Neighborhood

In an directed graph $G = (V, E)$,

- u is said to be **adjacent to** (nối tới) v and v is said to be **adjacent from** (được nối từ) u if (u, v) is an arc of G , and
- u is called **initial vertex** (đỉnh đầu) of (u, v)
- v is called **terminal** (đỉnh cuối) or **end vertex** of (u, v)
- the initial vertex and terminal vertex of a loop are the same.

The degree of a vertex

In a graph G with directed edges:

- **in-degree** (bậc vào) of a vertex v , denoted by $\deg^-(v)$, is the number of arcs with v as their terminal vertex.
- **out-degree** (bậc ra) of a vertex v , denoted by $\deg^+(v)$, is the number of arcs with v as their initial vertex.

Note: a loop at a vertex contributes **1** to both the in-degree and the out-degree of this vertex.



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Basic Theorem

Theorem

Let $G = (V, E)$ be a graph with directed edges. Then

$$\sum_{v \in V} \deg^-(v) = \sum_{v \in V} \deg^+(v) = |E|. \quad = \text{sum of edges}$$



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph


Bipartite graph

Isomorphism


Complete Graphs

A complete graph (đồ thị đầy đủ) on n vertices, K_n , is a simple graph that contains **exactly one edge** between each pair of distinct vertices.

$\deg(v)$ always equal $n-1$, $n \geq 1$



K_5



K_4



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph


Isomorphism

Cycles

A cycle (đồ thị vòng) C_n , $n \geq 3$, consists of n vertices v_1, v_2, \dots, v_n and edges $\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}$, and $\{v_n, v_1\}$. deg always equal 2, has n edges



C_5



C_4



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism


Wheels

We obtain a wheel (đồ thị hình bánh xe) W_n when we add an additional vertex to a cycle C_n , for $n \geq 3$, and connect this new vertex to each of the n vertices in C_n .


has $2n$ edges

$\deg v = n$

$\deg \text{ xung quanh} = 3$



W_5



W_4

BACHKHOACNCP.COM



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism


Exercise

Graph

Bipartite graph

Isomorphism

An n -dimensional hypercube (*khối n chiều*), Q_n , is a graph that has vertices representing the 2^n bit strings of length n . Two vertices are adjacent iff the bit strings that they represent differ in exactly one bit position.



Q_n is bipartite

What's about Q_4 ? has 16 vertices and $16 \cdot 4 / 2$ edges



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Applications of Special Graphs

- Local networks topologies
 - Star, ring, hybrid
- Parallel processing
 - Linear array
 - Mesh network

TÀI LIỆU SƯU TẬP
BỞI HCMUT-CNCP

BACHKHOACNCP.COM



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism

Exercise hw

Exercise (5)

Give the number of edges in function of number of vertices in a complete graph K_n . $(n-1)*n/2$

Exercise (6)

Give an undirected simple graph $G = (V, E)$ with $|V| = n$, show that

- a $\forall v \in V, \deg(v) < n$,
- b there does not exist simultaneously both a vertex of degree 0 and a vertex of degree $(n-1)$ with $n \geq 2$,
- c deduce that there are at least two vertices of the same degree.

$$0+1+2+\dots+n-1$$

Exercise (7)

Is it possible that each person has exactly 3 friends in the same group of 9 people ? $3*9=27$ ko chia het cho 2



Bipartite Graphs

Definition


A simple graph G is called bipartite (đồ thị phân đôi) if its vertex set V can be partitioned into two disjoint sets V_1 and V_2 such that every edge in the graph connects a vertex in V_1 and a vertex in V_2 (so that no edge in G connects either two vertices in V_1 or two vertices in V_2)

ne u do thi con la tam giac thi ko phan doi


Example

C_6 is bipartite

C_n is bipartite if n is even



C_6



C_n la do thi vong da hc o tren

BACHKHOACNCP.COM



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph


Isomorphism

Complete Bipartite Graphs

Definition

A complete bipartite $K_{m,n}$ is a graph that

- has its vertex set partitioned into **two subsets** of m and n vertices, respectively,
- with an edge between two vertices iff one vertex is in the first subset and the other is in the second one



$K_{3,3}$

mọi đỉnh dc nối với all đỉnh ở subset kia



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph


Bipartite graph

Isomorphism

Bipartite graphs

Example (Bipartite graphs?)

- C_6
- C_n
- K_3
- K_n
- the following graph



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism


New Graph From Old

Definition


A **subgraph** (đồ thị con) of a graph $G = (V, E)$ is a graph $H = (W, F)$ where $W \subseteq V$ and $F \subseteq E$.

Definition


The **union** (hợp) of two simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is a simple graph with vertex set $V_1 \cup V_2$ and edge set $E_1 \cup E_2$. The union of G_1 and G_2 is denoted by $G_1 \cup G_2$.



G_1



G_2



$G_1 \cup G_2$

BACHKHOACNCP.COM



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism


Exercise

Graph

Bipartite graph

Isomorphism

Planar Graphs



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph


Bipartite graph

Isomorphism

Planar Graphs

Definition

- A graph is called **planar** (*phẳng*) if it can be drawn in the plane **without any edges crossing**.
- Such a drawing is called **planar representation** (*biểu diễn phẳng*) of the graph.



K_4


K_4 with no crossing




Important Corollaries

Corollary

- If G is a **connected planar simple graph** with e edges and v vertices where $v \geq 3$, then $e \leq 3v - 6$.
 - If G is a **connected planar simple graph** with e edges and v vertices where $v \geq 3$, and **no circuits of length 3**, then $e \leq 2v - 4$.
- không chu trình 3 cạnh



$K_{3,3}$
Non-planar



K_5
Non-planar



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph

Isomorphism

Definition

- Given a planar graph G , an **elementary subdivision** (*phân chia sơ cấp*) is removing an edge $\{u, v\}$ and adding a new vertex w together with edges $\{u, w\}$ and $\{w, v\}$.
- Graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are called **homeomorphic** (*đồng phôi*) if they can be obtained from the same graph by a sequence of elementary subdivisions.



BACHKHOACNCP.COM



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph



Bipartite graph

Isomorphism

Kuratowski's Theorem

Theorem


A graph is nonplanar iff it contains a subgraph homeomorphic to $K_{3,3}$ or K_5 .




Exercise

Exercise

- Is K_4 planar?
- Is Q_3 planar?



K_4



Q_3



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise


Graph

Bipartite graph


Isomorphism

Adjacency Lists (Danh sách kề)

Vertex	Adjacent vertices
a	b, c, e
b	a
c	a, d, e
d	c, e
e	a, c, d



Initial vertex	Terminal vertices
a	b, c, d, e
b	b, d
c	a, c, e
d	c, e
e	b, c, d



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Adjacency Matrices

Definition


Adjacency matrix (*Ma trận kề*) A_G of $G = (V, E)$

- Dimension $|V| \times |V|$

- Matrix elements

$$a_{ij} = \begin{cases} 1 & \text{if } (v_i, v_j) \in E \\ 0 & \text{otherwise} \end{cases}$$

vo huong : doi xung qua duong cheo chinh



$$\begin{matrix} & \begin{matrix} a & b & c & d \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Examples

Example

Give the graph defined by the following adjacency matrix

	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>
<i>A</i>	0	0	1	1	0
<i>B</i>	0	1	0	1	0
<i>C</i>	1	0	0	1	0
<i>D</i>	1	1	1	0	1
<i>E</i>	0	0	0	1	0



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Adjacency Matrices

Example

Give the directed graph defined by the following adjacency matrix

	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>
<i>A</i>	1	0	1	1	0
<i>B</i>	0	0	0	0	0
<i>C</i>	1	0	0	0	0
<i>D</i>	1	1	1	0	1
<i>E</i>	1	0	0	0	0



Contents

Graph definitions

- Terminology
- Special Graphs

Representing Graphs and Graph Isomorphism

- Representing Graphs

- Graph Isomorphism

Exercise


- Graph
- Bipartite graph
- Isomorphism

Incidence Matrices

Definition

Incidence matrix (ma trận liên thuộc) M_G of $G = (V, E)$

- Dimension $|V| \times |E|$
- Matrix elements
$$m_{ij} = \begin{cases} 1 & \text{if } e_j \text{ is incident with } v_i \\ 0 & \text{otherwise} \end{cases}$$



$$\begin{matrix} & e_1 & e_2 & e_3 & e_4 \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \end{matrix}$$

BACHKHOACNCP.COM



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism


Exercise

Graph
Bipartite graph
Isomorphism

Examples

Example

Give incidence matrix according to the following graph



Contents

Graph definitions

- Terminology
- Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

- Graph Isomorphism

Exercise


- Graph
- Bipartite graph
- Isomorphism

Graph Isomorphism

Definition

$G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are **isomorphic** (đẳng cấu) if there is a **one-to-one function** f from V_1 to V_2 with the property that a and b are adjacent in G_1 iff $f(a)$ and $f(b)$ are adjacent in G_2 , for all a and b in V_1 . Such a function f is called an **isomorphism** (một đẳng cấu).

(i.e. there is a one-to-one correspondence between vertices of the two graphs that preserves the adjacency relationship.)



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism


Representing Graphs

Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Bipartie graph



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism


Exercise

Graph

Bipartie graph

Isomorphism

Isomorphism ?



Contents

Graph definitions

- Terminology
- Special Graphs


Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Isomorphism?



Contents

Graph definitions

- Terminology
- Special Graphs


Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Isomorphism?



Contents

Graph definitions

- Terminology
- Special Graphs

Representing Graphs and Graph Isomorphism

- Representing Graphs
- Graph Isomorphism

Exercise

- Graph
- Bipartite graph
- Isomorphism

Are the simple graphs with the following adjacency matrices isomorphic ?

$$① \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$② \quad \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

$$③ \quad \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$



Contents

Graph definitions

Terminology
Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs
Graph Isomorphism

Exercise

Graph
Bipartite graph
Isomorphism

Determine whether the graphs (without loops) with the incidence matrices are isomorphic.

- $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}$ $\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$
- $\begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{pmatrix}$ $\begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{pmatrix}$
- Extend the definition of isomorphism of simple graphs to undirected graphs containing loops and multiple edges.
- Define isomorphism of directed graphs



Contents

Graph definitions

Terminology

Special Graphs

Representing Graphs and Graph Isomorphism

Representing Graphs

Graph Isomorphism

Exercise

Graph

Bipartite graph

Isomorphism