

Definition 1 ($\text{LRA}(G)$). Let $G \triangleq (N, T, P, S)$ be a context-free grammar.

- (i) $G' := (N', T', P', S') \triangleq (N \uplus \{S'\}, T \uplus \{\$, \}, P \cup \{[S' \rightarrow S\$]\}, S')$.
- (ii) $V \triangleq N \uplus T$. $V' \triangleq N' \uplus T'$.
- (iii) $\beta \xrightarrow{\text{rm}}_{P'} \gamma \xleftrightarrow{\text{def}} (\beta, \gamma) \in \{(\alpha Az, \alpha \omega z) \mid [A \rightarrow \omega] \in P', \alpha \in V'^*, z \in T'^*\}$.
- (iv) $I_{\text{LR}(0)} \triangleq \{[A \rightarrow \alpha \cdot \beta] \mid [A \rightarrow \alpha \beta] \in P'\}$.
- (v) $\text{Cl}(q) =_{\mu} q \cup \{[A \rightarrow \varepsilon \cdot \omega] \mid [A \rightarrow \omega] \in P', [B \rightarrow \beta \cdot A\gamma] \in \text{Cl}(q)\}$ for $q \in \mathcal{P}(I_{\text{LR}(0)})$.
- (vi) $q_0 := \text{Cl}(\{[S' \rightarrow \varepsilon \cdot S\$]\})$.
- (vii) $\text{GOTO}(q, X) := \text{Cl}(\{[A \rightarrow \alpha X \cdot \beta] \mid [A \rightarrow \alpha \cdot X\beta] \in q\})$ for $q \in \mathcal{P}(I_{\text{LR}(0)})$ and $X \in V'$.
- (vii) $Q \triangleq \text{PT} \setminus \{\emptyset\}$ where $\text{PT} =_{\mu} \{q_0\} \cup \{\text{GOTO}(q, X) \mid q \in \text{PT}, X \in V'\}$.
- (ix) $\varepsilon : p \rightarrow q \xleftrightarrow{\text{def}} p \in Q \wedge p = q$. $X\alpha : p \rightarrow q \xleftrightarrow{\text{def}} p \in Q \wedge \alpha : \text{GOTO}(p, X) \rightarrow q$.
- (x) **Config** $\triangleq \{(\alpha : p \rightarrow q, z) \mid \alpha \in V'^*, p \in Q, q \in Q, z \in T'^*, \alpha : p \rightarrow q\}$.
- (xi) $\delta(q, X) := \text{GOTO}(q, X)$ for $q \in Q$ and $X \in V'$.
- (xii) **reduce** $(q, t) := \{[A \rightarrow \omega] \mid [A \rightarrow \omega \cdot \varepsilon] \in q\}$ for $q \in Q$ and $t \in T'$.
- (vii) Let $\vdash$ be a binary relation on the set **Config** with two introduction rules:

$$\frac{q'' = \delta(q', t)}{(\alpha : q \rightarrow q', tz) \vdash (\alpha t : q \rightarrow q'', z)} \text{ Shift}$$

$$\frac{\alpha : q \rightarrow p \quad \omega : p \rightarrow q' \quad [A \rightarrow \omega] \in \text{reduce}(q', t) \quad q'' = \delta(p, A)}{(\alpha \omega : q \rightarrow q', tz) \vdash (\alpha A : q \rightarrow q'', tz)} \text{ Reduce}(A \rightarrow \omega)$$

- (xiv) $q_f := \delta(\delta(q_0, S), \$)$.
- (xv) The language accepted by $\text{LRA}(G)$ is defined as follows:

$$L(\text{LRA}(G)) := \{z \in T^* \mid (\varepsilon : q_0 \rightarrow q_0, z\$) \vdash^* (S\$: q_0 \rightarrow q_f, \varepsilon)\}.$$

Fact 2. It is known that $L(G) = L(\text{LRA}(G))$, where $L(G) \triangleq \{z \in T^* \mid S \Rightarrow_P^* z\}$.

Theorem 3. Define **LA** : $\{(q, [A \rightarrow \omega]) \mid q \in Q, [A \rightarrow \omega \cdot \varepsilon] \in q\} \rightarrow \mathcal{P}(T')$ by

$$\mathbf{LA}(q, [A \rightarrow \omega]) := \left\{ t \in T' \mid S' \xrightarrow{\text{rm}}_{P'}^* \alpha A t z, \alpha \omega : q_0 \rightarrow q, \alpha \in V'^*, z \in T'^* \right\}. \quad (1)$$

Then, overriding **reduce** : $Q \times T' \rightarrow \mathcal{P}(P')$ of the $\text{LR}(0)$ parser $\text{LRA}(G)$ with

$$(q, t) \mapsto \{[A \rightarrow \omega] \mid [A \rightarrow \omega \cdot \varepsilon] \in q, t \in \mathbf{LA}(q, [A \rightarrow \omega])\}$$

yields an $\text{LALR}(1)$ parser if there are no conflicts, which accepts the same language.

Theorem 4. Letting $R!x \triangleq \{y \mid (x, y) \in R\}$, define relations **Read** and **Follow** from the set

$$\{(p, A) \mid p \in Q, A \in N', \delta(p, A) \neq \emptyset\}$$

to the set T' inductively as follows:

$$\frac{\delta(\delta(p, A), t) \neq \emptyset}{t \in \mathbf{Read}!(p, A)} \text{ DR} \quad \frac{\delta(p, A) = r \quad C \Rightarrow_{P'}^* \varepsilon}{\mathbf{Read}!(r, C) \subseteq \mathbf{Read}!(p, A)} \text{ reads}$$

$$\frac{t \in \mathbf{Read}!(p, A)}{t \in \mathbf{Follow}!(p, A)} \text{ Read} \quad \frac{[B \rightarrow \beta \cdot A\gamma] \in p \quad \beta : p' \rightarrow p \quad \gamma \Rightarrow_{P'}^* \varepsilon}{\mathbf{Follow}!(p', B) \subseteq \mathbf{Follow}!(p, A)} \text{ includes}$$

Then, whenever $q \in Q$ and $[A \rightarrow \omega \cdot \varepsilon] \in q$, we can compute $\mathbf{LA}(q, [A \rightarrow \omega])$ by

$$\mathbf{LA}(q, [A \rightarrow \omega]) = \{t \in T' \mid p \in Q, \omega : p \rightarrow q, \delta(p, A) \neq \emptyset, t \in \mathbf{Follow}!(p, A)\}. \quad (2)$$