

# TITLE

Cody Buntain, Christopher Natoli, Miroslav Živković

OSDC PIRE 2014, hosted at University of Amsterdam

24 July 2014

# TITLE

- item



# BUILDING THE TEST STATISTIC

$$\text{test statistic} = \frac{\sum_{t=1}^h e_t}{\quad}$$



# BUILDING THE TEST STATISTIC

$$\text{test statistic} = \frac{\sum_{t=1}^h \mathbf{e}_t^\top}{\quad} \mathbf{e}_t$$

# BUILDING THE TEST STATISTIC

$$\text{test statistic} = \frac{\sum_{t=1}^h \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{h}$$

$$\hat{\Sigma} := \frac{1}{n-1} \sum_{t=1}^n \mathbf{e}_t \mathbf{e}_t^\top$$


# BUILDING THE TEST STATISTIC

$$\text{test statistic} = \frac{\sum_{t=1}^h \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{h} - \frac{\sum_{t=1}^n \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{n}$$

# BUILDING THE TEST STATISTIC

$$\text{test statistic} = \frac{h}{\sqrt{2kn}} \left( \frac{\sum_{t=1}^h \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{h} - \frac{\sum_{t=1}^n \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{n} \right)$$

# FINDING A SINGLE CHANGEPPOINT



Cusum statistic for a time series with three changepoints.






# FINDING A SINGLE CHANGEPOINT

$$\max(\text{test statistic}) = \max \left( \frac{h}{\sqrt{2kn}} \left( \frac{\sum_{t=1}^h \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{h} - \frac{\sum_{t=1}^n \mathbf{e}_t^\top (\hat{\Sigma})^{-1} \mathbf{e}_t}{n} \right) \right)$$
$$\xrightarrow{D} \sup \{ \text{Brownian bridge} \},$$

which is a known distribution!

## FINDING MORE CHANGEPOINTS



Cusum statistic for a time series with three changepoints.

# TITLE

- item



# TITLE

- item