

# Revision of introduction to epidemiology

Dr David A McAllister

Wellcome Trust Intermediate Clinical  
Fellow

# Aim


- To review the epidemiological concepts covered in previous courses

# **BASIC EPIDEMIOLOGICAL TERMS**

# Mortality rates

- Am I worried?
  - 50 people died from flu
  - 50 people in Glasgow died from flu
  - 50 people in Glasgow died from flu in past 100 years
- Meaningful statistics need
  - A denominator population
  - A time frame

# Scotland's population 2013



# Other denominators

- Health board
- City
- Hospital
- Disease register
- Recruited to a study

**THE DENOMINATOR MUST CORRESPOND TO THE NUMERATOR**

# Two sorts of time

- Person-time
  - 10 deaths per 10,000 person years
    - 10,000 people for 1 year
    - 5,000 people for 2 years
    - 2,000 people for 5 years
- n-year follow-up
  - 5-year mortality of 10 per 10,000 people
- **Without denominator population and time death rates are meaningless**

# Incidence

- Number of new cases
  - person-time, eg 5 per 10,000 person years
  - n years of follow-up, 10-year cancer incidence rate is 1 per 10,000



# Prevalence

- Proportion of population that has disease
  - at a specified time, eg 1% of population had COPD in 2010
  - over a specified period, eg lifetime prevalence of dementia 40%

# Compare/contrast

## **Prevalence**

- A proportion
- Useful for planning services
- Depends partly on incidence

## **Incidence**


- A rate or a proportion
- Useful for identifying causes of diseases
- Occurs, by definition, only in people without the disease

# How incidence relates to prevalence



# Example - heart failure

Increased myocardial infarction  
↓  
Increased heart failure incidence



Improved long-term management

→  
Reduced death rate

# **ASSOCIATIONS BETWEEN EXPOSURES AND OUTCOMES**

# Outcomes

- Outcomes
  - death
  - hospitalisation
  - first diagnosis with a disease
  - recurrence, eg cancer
  - quality of life
  - surrogates – blood pressure, lung function


# Exposures

- Non-modifiable
  - age, sex, genotype
- Modifiable
  - smoking, weight, diet, alcohol consumption
- Interventions – a special kind of exposure
  - drug therapy
  - surgery
  - lifestyle advice

# Association

- a and b are associated when a and b are not independent
  - $P(a | b) = P(a)$
  - $P(b | a) = P(b)$
- Knowing something about a tells you nothing about b
- Knowing something about b tells you nothing about a





# Effect estimates

	Died	N	Proportion
Thrombolysis	100	1000	10%
No thrombolysis	130	1000	13%

$$RR = \frac{risk_{exposed}}{risk_{unexposed}}$$

$$ARR = risk_{unexposed} - risk_{exposed}$$

Measure	Result
Risk ratio (RR)	=10/13 = 0.77
Relative risk reduction (RRR)	100 x (1-0.77) = 23%
Risk difference (absolute risk reduction – ARR)	= 13-10 = 3%
Number needed to treat (NNT)	= 1/0.03 = 33

$$RRR = 100 \times (1 - RR)$$

$$NNT = \frac{1}{ARR}$$

# Other commonly encountered measures

- Odds ratio – commonly used estimate of risk ratio
- Rate ratio – ratio between two mortality rates, hospitalisation rates etc
- Hazard ratio – a special kind of rate ratio


# **CONFIDENCE INTERVALS**

# Confidence intervals

- Vital to reading ANY! research paper
- Formal definitions are counterintuitive
- Pragmatically
  - “A confidence interval can be thought of as a range of plausible values”
  - Values near the limits are much less plausible than those in the middle


**EXAMPLE – AS IN TABLE EFFECT ESTIMATES -  
THROMBOLYSIS**

# 95% CI represented visually




Don't worry about the units on the y-axis  
Don't worry about formal definitions!!!

# Relative risk







# Relative risk reduction



# Risk difference (absolute risk reduction)



# Number needed to treat (NNT)



# Confidence intervals

- Can be presented for any statistic/effect measure
- Represents range of plausible values
- More extreme values less likely
- Very useful in appraising published research

# **DESCRIPTIVE EPIDEMIOLOGY**

# Crude mortality

State	Deaths in 2013	Population in 2013	Crude annual mortality
New York City	48,000	8,000,000	6 per 1,000
Florida	19,000	19,000,000	10 per 1000
		Rate ratio	1.67

# WHY?

# Standardised mortality ratio (SMR)

State	Deaths	Expected deaths	SMR
New York City	48,000	50,000	96
Florida	19,000	22,000	86

- Calculate expected deaths based on
  - age-sex specific mortality rates in whole of US
  - age and sex of people in NY and Florida
- Both places have lower than expected mortality
- New York has a higher SMR than Florida

**CONFOUNDING**



# NY vs Florida was an example of confounding


- True relationship confused by a third factor
- Can deal with confounding
  - study design
  - data analysis – eg via standardisation

# Bias


- **Systematic** error in
  - what data are collected
  - how data are collected
  - how data are analysed
  - how data are interpreted
  - how data are reported
- Bias leads to wrong conclusions about
  - disease causation
  - treatment effectiveness

# Bias - examples


- **Systematic** error in
  - what data are collected - novel treatments
  - how data are collected - different follow-up
  - how data are analysed - misconduct
  - how data are interpreted - highlight certain findings
  - how data are reported - publication bias
- Causes wrong conclusions about
  - disease causation
  - treatment effectiveness



All studies





Published  
studies



**DIFFERENT TYPES OF STUDY ARE  
MORE/LESS PRONE TO  
CONFOUNDING AND BIAS**

# Hierarchy of evidence



# Case-control study

- Select CASES with a disease – eg lung cancer
- Select CONTROLS without that disease
- Find out what EXPOSURES the cases and controls had – eg smoking
- Compare exposures in cases and controls
- Identify if association

# Cohort study

- Select people without a disease (eg heart disease)
- Classify them according to an exposure eg
  - high cholesterol
  - low cholesterol
- Follow-them, eg for 10 years
- Compare RISK of disease in exposed and unexposed





# Randomised controlled trial (RCT)

- Like cohort study, except
- **ALLOCATE PEOPLE TO INTERVENTION**
  - cohort study ‘observe exposure’
  - randomised controlled trial ‘allocate intervention’
- Compare RISK of outcome in treatment and control groups

# Cross-sectional study

- Sample a population
- Estimate the proportion of people with
  - different exposures (eg smoking, alcohol consumption, income)
  - different symptoms (eg breathlessness, cough)
  - different diseases (eg angina)
- Use these data
  - to describe disease burden
  - to explore associations

# Hierarchy of evidence



# Topics covered

- Basic epidemiological terms
- Confidence intervals
- Confounding
- Bias
- Case-control and cohort studies
- The hierarchy of evidence

Questions?