

HW Chapter 1. Aug. 30th.

4) $f(t) = \frac{\sin^2 t}{t}$.

Sol: $|f(t)| \leq \left| \frac{\sin t}{t} \right| \cdot |\sin t| \leq 1 \cdot |\sin t| \leq 1$. on $t \in [0, +\infty)$. thus $\alpha_0 = 0$. $M = 1$.

denote $g(t) = \sin^2 t = \frac{1 - \cos 2t}{2}$.

$$g(t) \leftrightarrow G(p). \text{ by the table, } G(p) = \frac{1}{2} \left(\frac{1}{p} - \frac{p}{p^2+4} \right) = \frac{1}{2} \cdot \frac{p^2+4-p^2}{p(p^2+4)} = \frac{2}{p(p^2+4)}$$

by the property of integration of image.

$$f(t) = \frac{g(t)}{t} \leftrightarrow F(p) = \int_p^\infty G(\xi) d\xi = \frac{1}{2} \int_p^\infty \left(\frac{1}{\xi} - \frac{\xi}{\xi^2+4} \right) d\xi = \int_p^\infty \frac{d\xi^2}{\xi^2(\xi^2+4)} = \frac{1}{4} \ln \left(1 + \frac{4}{p^2} \right)$$

$$\text{i.e. } f(t) = \frac{g(t)}{t} \leftrightarrow F(p) = \frac{1}{4} \ln \left(1 + \frac{4}{p^2} \right), \quad p > 0$$