

Equations of Mathematical Physics

Lecturer Regina Saitova

PhD, Associate Professor at St. Petersburg State University

Autumn 2024

CONTENTS

Chapter 1. LINEAR DIFFERENTIAL OPERATORS	
Chapter 2. ONE-DIMENSIONAL WAVE EQUATION	
Chapter 3. STURM-LIOUVILLE PROBLEM	
Chapter 4. GENERALIZED FUNCTIONS	
Chapter 5. CONVOLUTION AND FOURIER TRANSFORM	
Chapter 6. HEAT CONDUCTION EQUATION	
Chapter 7. SOBOLEV SPACES. GENERALIZED SOLUTION OF THE DIRICHLET PROBLEM	
Chapter 8. EIGENVALUES AND EIGENFUNCTIONS OF THE LAPLACE OPERATOR	
Chapter 9. WAVE EQUATION	
Chapter 10. PROPERTIES OF POTENTIALS AND THEIR CALCULATION	
Chapter 11. WAVE FRONT AND SHORT-WAVE APPROXIMATION FOR HYPERBOLIC EQUATIONS	

LITERATURE

1. **Michael Reed, Barry Simon:** Methods of Modern Mathematical Physics. Academic Press New York London, 4 Volumes (1972).
2. **Robert D. Richtmyer:** Principles of Advanced Mathematical Physics. Springer-Verlag New York Heidelberg Berlin, 2 Volumes (1978).
3. **Jon Mathews, Walker R.L.:** Mathematical Methods of Physics. W.A. Benjamin, inc. New York-Amsterdam (1964).
4. **Vladimirov V.S.:** Equations of Mathematical Physics. Marcel Dekker Incorporated. 432 p. (1971).
5. **Shubin M.A.:** Lectures on Equations of Mathematical Physics. Moscow: MCCME. 303 p. (2003).

Chapter 1

LINEAR DIFFERENTIAL OPERATORS

1.1. Definition and examples

We introduce notations that are convenient when using functions of several variables and differential operators.

A multi-index α is a set of $\alpha = (\alpha_1, \dots, \alpha_n)$, where $\alpha_j \in \mathbb{Z}_+$ (that is, α_j are non-negative integers).

If α is a multi-index, then we will put $|\alpha| = \alpha_1 + \dots + \alpha_n$, $\alpha! = \alpha_1! \dots \alpha_n!$, and if vector $x = (x_1, \dots, x_n) \in \mathbb{C}^n$ is given, then $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$.

If Ω is an open subset in \mathbb{R}^n , then by $C^\infty(\Omega)$ we denote **the set of complex-valued infinitely differentiable functions** in Ω , and by $C_0^\infty(\Omega)$ we denote **the set of finite infinitely differentiable functions** in Ω , that is, such $\varphi \in C^\infty(\Omega)$ that there exists a compact $K \subset \Omega$, outside of which the function φ turns to 0 (compact K depends on the function φ).

Let's say

$$\partial_j = \frac{\partial}{\partial x_j} : C^\infty(\Omega) \rightarrow C^\infty(\Omega); \quad D_j = -i\partial_j,$$

where $i = \sqrt{-1}$; $\partial = (\partial_1, \dots, \partial_n)$; $D = (D_1, \dots, D_n)$; $\partial^\alpha = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n}$;

$$D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}.$$

Thus,

$$\partial^\alpha = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}}$$

is a mixed derivative operator, $\partial^\alpha: C^\infty(\Omega) \rightarrow C^\infty(\Omega)$, $D^\alpha = i^{-|\alpha|} \partial^\alpha$.

If $f \in C^\infty(\Omega)$, we instead $\partial^\alpha f$ will also sometimes write $f^{(\alpha)}$.

Linear differential operator is an operator

$$A: C^\infty(\Omega) \rightarrow C^\infty(\Omega)$$

$$A = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha, \quad (1.2)$$

where $a_\alpha(x) \in C^\infty(\Omega)$.

Instead D^α we can write ∂^α , but written through D^α more convenient, as will be seen from further.

Here $m \in \mathbb{Z}_+$ and we will say that A - operator order $\leq m$.

We will say that A operator order m , if it is written in the form (1.2) and there is a multi-index α , which $|\alpha|=m$ and $a_\alpha(x) \neq 0$.

Examples.

1. Laplace operator

$$\Delta = \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2} = -(D_1^2 + \dots + D_n^2).$$

2. Thermal conductivity operator

$$\frac{\partial}{\partial t} - \Delta$$

(here the number of variables is equal to $n+1$ and they are denoted by t, x_1, \dots, x_n).

3. Wave operator or d'Alambertian

$$\square = \frac{\partial^2}{\partial t^2} - \Delta.$$

4. Operator of Sturm-Liouville, determined by formula

$$Lu = \frac{d}{dx} \left(p(x) \frac{du}{dx} \right) + q(x)u,$$

where $p, q \in C^\infty((a, b))$, $n = 1$, $\Omega = (a, b)$.

Examples operators 1)-3) – are operators with constant coefficients, the operator of Sturm-Liouville has variable coefficients.

1.2. Full and main symbols

The symbol or the full symbol of the operator A of the order m is called a function of the form

$$a(x, \xi) = \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha, \quad x \in \Omega, \quad \xi \in \mathbb{R}^n, \quad (1.3)$$

and the main symbol is a function of the form

$$a_m(x, \xi) = \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha, \quad x \in \Omega, \quad \xi \in \mathbb{R}^n. \quad (1.4)$$

The symbol belongs to $C^\infty(\Omega)[\xi]$, that is, it is a polynomial of ξ_1, \dots, ξ_n with coefficients from the ring $C^\infty(\Omega)$, and the full symbol is a homogeneous polynomial of ξ_1, \dots, ξ_n degree m with coefficients from $C^\infty(\Omega)$.

Examples.

1. The Laplace operator has matching full and main symbols

$$\xi^2 = \xi_1^2 + \dots + \xi_n^2.$$

2. The thermal conductivity operator $\frac{\partial}{\partial t} - \Delta$ has a full symbol $a(t, x, \tau, \xi) = i\tau + \xi^2$, and its main symbol is $a_2(t, x, \tau, \xi) = \xi^2$.

3. The wave operator has matching full and main symbols

$$a(t, x, \tau, \xi) = a_2(t, x, \tau, \xi) = -\tau^2 + \xi^2.$$

4. The Sturm-Liouville operator has the full symbol $a(x, \xi) = -p(x)\xi^2 + ip'(x)\xi + q(x)$, and the main symbol $a_2(x, \xi) = -p(x)\xi^2$.

The symbol of the operator A is restored according to the operator by the formula

$$a(x, \xi) = e^{-ix \cdot \xi} A(e^{ix \cdot \xi}), \quad (1.5)$$

in which the operator A is applied according to x .

The notation $x \cdot \xi = x_1 \xi_1 + \dots + x_n \xi_n$ is used here.

The formula (1.5) is obtained from the ratio

$$D^\alpha e^{ix \cdot \xi} = \xi^\alpha e^{ix \cdot \xi},$$

which is easily verified by induction according to $|\alpha|$. It also follows from (1.5) that the coefficients $a_\alpha(x)$ of the operator A are uniquely determined.

It is useful to apply the operator A to a more general exponent than $e^{ix \cdot \xi}$ to the exponent $e^{i\lambda \varphi(x)}$, where $\varphi \in C^\infty(\Omega)$, λ is a parameter.

Lemma 1.1. If $f \in C^\infty(\Omega)$, then $e^{-i\lambda \varphi(x)} A(f(x) e^{i\lambda \varphi(x)})$ is a polynomial of λ degree $\leq m$ with coefficients from $C^\infty(\Omega)$, and

$$e^{-i\lambda \varphi(x)} A(f(x) e^{i\lambda \varphi(x)}) = \lambda^m f(x) a_m(x, \varphi_x) + \lambda^{m-1}(\dots) + \dots, \quad (1.6)$$

that is, the highest coefficient of this polynomial (for λ^m) is $f(x)a_m(x, \varphi_x)$, where $\varphi_x = (\frac{\partial \varphi}{\partial x_1}, \dots, \frac{\partial \varphi}{\partial x_n}) = \text{grad } \varphi$ is the gradient vector φ .

Proof: We have

$$D_j [f(x) e^{i\lambda \varphi(x)}] = \lambda (\partial_j \varphi)(x) e^{i\lambda \varphi(x)} + (D_j f) e^{i\lambda \varphi(x)},$$

whence the statement of the lemma is obtained for operators of order less than 1 (≤ 1). In the future, when finding arbitrary $D^\alpha e^{i\lambda \varphi(x)}$, it will also be necessary to differentiate products of the form $f(x) e^{i\lambda \varphi(x)}$, where $f \in C^\infty(\Omega)$. In this case, a new multiplier λ appears only when the exponent is differentiated. Therefore, it is clear that

$$D^\alpha e^{i\lambda \varphi(x)} = \lambda^{|\alpha|} \varphi_x^\alpha e^{i\lambda \varphi(x)} + \lambda^{|\alpha|-1}(\dots) + \dots,$$

whence the statement of the Lemma follows.

Corollary 1.2. Let A, B be two linear differential operators in Ω , k, l be their orders, and a_k, b_l are the main symbols. Next, let $C = A \circ B$ be the composition of these operators, and c_{k+l} be its main symbol. Then

$$c_{k+l}(x, \xi) = a_k(x, \xi) b_l(x, \xi). \quad (1.7)$$

Comment. Obviously, C is a differential operator of the order $\leq k+l$.

Proof of corollary 1.2:

$$\begin{aligned} C(e^{i\lambda \varphi(x)}) &= A(\lambda^l b_l(x, \varphi_x) e^{i\lambda \varphi(x)} + \lambda^{l-1}(\dots) + \dots) = \\ &= \lambda^{k+l} a_k(x, \varphi_x) b_l(x, \varphi_x) e^{i\lambda \varphi(x)} + \lambda^{k+l-1}(\dots) + \dots \end{aligned}$$

Hence, according to Lemma 1.1. it follows

$$c_{k+l}(x, \varphi_x) = a_k(x, \varphi_x) b_l(x, \varphi_x) \quad (1.8)$$

for any function $\varphi \in C^\infty(\Omega)$.

But then, choosing $\varphi(x) = x \cdot \xi$, we get $\varphi_x = \xi$ and (1.8) goes to

$$c_{k+l}(x, \xi) = a_k(x, \xi) b_l(x, \xi),$$

which was to be proved.

1.3. Change of variables

Let's give a diffeomorphism $\kappa: \Omega \rightarrow \Omega_1$, where Ω, Ω_1 are regions in \mathbb{R}^n . Such a diffeomorphism can be defined by a set of functions $y_1(x_1, \dots, x_n), \dots, y_n(x_1, \dots, x_n)$, $y_j \in C^\infty(\Omega)$, equal to the coordinates of point x at point $\kappa(x)$.

If $f \in C^\infty(\Omega_1)$, then we will put $\kappa^* f = f \circ \kappa$,

that is, $(\kappa^* f)(x) = f(\kappa(x))$ or $(\kappa^* f)(x) = f(y_1(x), \dots, y_n(x))$.

Essentially, $\kappa^* f$ is obtained from f by replacing variables or moving to coordinates x_1, \dots, x_n from coordinates y_1, \dots, y_n . The diffeomorphism κ has an inverse map κ^{-1} , which is also a diffeomorphism. It is clear that κ^* is a linear isomorphism of $\kappa^*: C^\infty(\Omega_1) \rightarrow C^\infty(\Omega)$, and $(\kappa^*)^{-1} = (\kappa^{-1})^*$.

Let's give the operator $A: C^\infty(\Omega) \rightarrow C^\infty(\Omega)$. Let's define the operator $A_1: C^\infty(\Omega_1) \rightarrow C^\infty(\Omega_1)$ using a commutative diagram

$$\begin{array}{ccc} C^\infty(\Omega) & \xrightarrow{A} & C^\infty(\Omega) \\ \uparrow \kappa^* & & \uparrow \kappa^* \\ C^\infty(\Omega_1) & \xrightarrow{A_1} & C^\infty(\Omega_1) \end{array}$$

that is, $A_1 = (\kappa^*)^{-1} A \kappa^* = (\kappa^{-1})^* A \kappa^*$.

In other words,

$$A_1 v(y) = \{A[v(y(x))]\} \Big|_{x=x(y)}, \quad (1.9)$$

where $y(x) = \kappa(x)$, $x(y) = \kappa^{-1}(y)$.

Thus, the A_1 operator is essentially just a written A operator in y coordinates.

If A is a linear differential operator, then from (1.9) and the formula for differentiating a complex function it is easily verified that A_1 is also a linear differential operator.

Let's find out how the main symbols of the operators A and A_1 are connected.

Reminder: *cotangent vector* or *covector* at point $x \in \Omega$ is a linear functional on the tangent space Ω to at point x (we will denote the tangent space to Ω at point x by $T_x \Omega$, and the cotangent space - the set of all cotangent vectors at point x - by $T_x^* \Omega$).

By $T^* \Omega$ we denote the union of $\bigcup_x T_x^* \Omega \cong \Omega \times \mathbb{R}^n$.

Choosing the basis of vectors $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}$ in each tangent space $T_x \Omega$, we construct a dual basis in $T_x^* \Omega$: it consists of such functionals

dx_1, \dots, dx_n that $dx_i \left(\frac{\partial}{\partial x_j} \right) = \delta_{ij}$, where δ_{ij} is the Kronecker symbol ($\delta_{ij} = 0, i \neq j$ and $\delta_{ij} = 1, i = j$).

An **example** of a tangent vector is the velocity vector of a curve $x(t): \dot{x}(0) \in T_{x(0)}\Omega$.

Its coordinates in the basis $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}$ are equal to $\frac{dx_1}{dt}|_{t=0}, \dots, \frac{dx_n}{dt}|_{t=0}$.

An **example** of a cotangent vector is the differential or gradient of the function $f \in C^\infty(\Omega)$: this is a functional on $T_x\Omega$, whose value on the tangent vector $\dot{x}(0)$ is equal to $\frac{df(x(t))}{dt}|_{t=0}$; its coordinates in the basis dx_1, \dots, dx_n are equal to $\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}$.

If the diffeomorphism $\kappa: \Omega \rightarrow \Omega_1$ is given, then there is a natural mapping (the differential of the mapping κ):

$$\kappa_*: T_x\Omega \rightarrow T_{\kappa(x)}\Omega_1$$

and a dual mapping on linear functionals

$${}^t\kappa_*: T_{\kappa(x)}^*\Omega_1 \rightarrow T_x^*\Omega.$$

Choosing the bases in $T_x\Omega$, $T_{\kappa(x)}\Omega_1$, $T_{\kappa(x)}^*\Omega_1$, $T_x^*\Omega$, respectively, $\left\{\frac{\partial}{\partial x_j}\right\}_{j=1}^n$, $\left\{\frac{\partial}{\partial y_j}\right\}_{j=1}^n$, $\{dy_j\}_{j=1}^n$, $\{dx_j\}_{j=1}^n$, we get that κ_* has a matrix called the Jacobian matrix: $(\kappa_*)_{kl} = \frac{\partial y_k}{\partial x_l}$, and ${}^t\kappa_*$ has a matrix transposed to the Jacobian matrix κ_* .

Note that the maps κ_* and ${}^t\kappa_*$ are isomorphisms at each point of $x \in \Omega$.

Next, it is convenient to set the cotangent vector at point x by specifying a pair (x, ξ) , where $\xi \in \mathbb{R}^n$, $\xi = (\xi_1, \dots, \xi_n)$ is the set of coordinates of this vector in the basis $\{dx_j\}_{j=1}^n$.

With isomorphism, $({}^t\kappa_*)^{-1}: T^*\Omega \rightarrow T^*\Omega_1$ the covector (x, ξ) corresponds to the covector (y, η) , where $y = \kappa(x)$, $\eta = ({}^t\kappa_*)^{-1}\xi$.

Theorem 1.3.

The main symbol a_m of operator A takes the same value on vector (x, ξ) as the main symbol a'_m of operator A_1 on the corresponding vector (y, η) , that is,

$$a_m(x, \xi) = a'_m(\kappa(x), ({}^t\kappa_*)^{-1}\xi). \quad (1.10)$$

In other words, the main symbol is a well-defined function on $T^*\Omega$ (independent of the choice of curved coordinates in Ω).

Proof:

The cotangent vector at point x can be written as the gradient $\varphi_x(x)$ of the function $\varphi \in C^\infty(\Omega)$ at point x .

From Lemma 1.1. it is clear that $a_m(x, \varphi_x(x))$ does not depend on the choice of coordinates. But on the other side, it is clear that this value does not depend on the choice of the function φ with a given differential at point x . Therefore, the main symbol is correctly assigned to $T^*\Omega$, which was required.

→ 1.4. Reduction of operators of the 2nd order with constant coefficients to the canonical form.

- ➔ **1.5. Characteristics. Ellipticity and hyperbolicity**
- ➔ **1.6. Characteristics and reduction to the canonical form of operators and equations of the 2nd order at $n=2$**
- ➔ **1.7. General solution of a homogeneous hyperbolic equation with constant coefficients at $n=2$**