

Chapter 12. Introduction to ordinary differential equations.

Initial-value problem (Cauchy's problem) for first-order equation:

Find a solution $y(x)$ of the equation

$$y' = f(x, y) \tag{1}$$

which satisfies the initial condition

$$y(a) = y_0, \tag{2}$$

where $f(x, y)$ is a given function in the domain $a \leq x \leq b$, $-\infty < y < \infty$.

Theorem 1.

If functions $f(x,y)$ and $\partial f(x,y)/\partial y$ are continuous and bounded in the domain $a \leq x \leq b$, $-\infty < y < \infty$, then there exists a unique solution $y(x)$ of problem (1),(2) (see proof in a course of Diff. Equations)

Euler's algorithm of calculations:

Let us divide the segment $a \leq x \leq b$ into n subsegments of length h each by the points

$$x_0=a, \quad x_1, \quad x_2, \dots, \quad x_n=b \qquad h=(b-a)/n$$

As shown in Chapter 12,

$$y'(x_i) = [y(x_{i+1}) - y(x_i)] / h + O(h)$$

$$y(x_{i+1}) - y(x_i) = h y'(x_i) - h \cdot O(h)$$

Now we replace $y'(x_i)$ by the right-hand side of equation (1):

$$y(x_{i+1}) - y(x_i) = h f(x_i, y(x_i)) - \underline{h \cdot O(h)}$$

If we omit the error $h \cdot O(h)$, then obtain formula for calculation of the approximate solution:

$$y_{i+1} = y_i + h f(x_i, y_i) \quad (3)$$
$$i=0, 1, 2, \dots, n-1$$

Theorem 2.


Values y_i of the approximate solution of problem (1), (2) at points x_i can be calculated using (3). The error of the solution is estimated by:

$$\max_{i=0,1,\dots,n} | y(x_i) - y_i | = O(h)$$

approximate solution

exact solution guaranteed by Theorem 1

**A broken line illustrates the
approximate solution
(keep in mind the linear
interpolation)**



Example.

$$y' = y \cdot \sin(3x), \quad y(0)=1, \quad 0 \leq x \leq 5$$

Scilab

```
clear
```

```
n=50 // number of subsegments
```

```
a=0
```

```
b=5
```

```
h= (b-a)/n
```

```
for i=1:n+1
```

```
  x(i)= a+ h*(i-1)
```

```
  y(i)=1
```

```
end
```

```
for i=1:n
```

```
  f=y(i)*sin(3*x(i))
```

```
  y(i+1)= y(i)+h*f
```

```
end
```

```
plot(x,y,'ob')
```

Improved Euler's algorithm


(error at 1 step is of order $O(h^3)$)

$$y' = f(x, y)$$

First, calculate $\hat{y}_{i+1} = y_i + h f(x_i, y_i)$ following (3)

Then make a more accurate step:

$$y_{i+1} = y_i + h [f(x_i, y_i) + f(x_{i+1}, \hat{y}_{i+1})]/2 \quad (4)$$



Derivation of (4) is based on the expression of second-order accuracy for y'_i at inner nodes, see Chapter 12:

$$[\ y'_i = (y_{i+1} - y_{i-1}) / (2h) + O(h^2) \]$$

In this case: $y'_{i+1/2} = (y_{i+1} - y_i) / (2h/2) + O(h^2/4)$

$$(y_{i+1} - y_i) / (2h/2) + O(h^2/4) = f(x_{i+1/2}, y_{i+1/2})$$

$$(y_{i+1} - y_i) / h + O(h^2) = [f(x_i, y_i) + f(x_{i+1}, y_{i+1})] / 2 + O(h^2)$$

$$y_{i+1} - y_i = h [f(x_i, y_i) + f(x_{i+1}, y_{i+1})] / 2 + O(h^3)$$

$$y_{i+1} - y_i = h [f(x_i, y_i) + f(x_{i+1}, \hat{y}_{i+1})] / 2 + O(h^3)$$

(details will be explained in a lecture in November)

```
n=50                                // improved Euler
a=0
b=5
h= (b-a)/n
for i=1: n+1
x(i)= a+ h*(i-1)
end
y(1)= 1
for i=1:n
f1=y(i)*sin(3*x(i))
ysimple= y(i)+h*f1
f2 = ysimple*sin(3*x(i+1))
y(i+1)=y(i)+h*0.5*(f1+f2)
end
plot(x,y,'r')
```


Scilab: `y = ode(y0, x0, x, right2)`

```
function myfunction=f(x,y)
```

```
    myfunction = y*sin(3*x)
```

```
endfunction
```

```
y0 = 1;
```


```
x0 = 0;
```

```
x = 0:0.1: 5;
```

```
y = ode(y0, x0, x, f);
```

```
plot(x,y,'k')
```

by default it uses a predictor-corrector Adams method



John Adams (photo ≈ 1870)

y =ode("rk", y0, x0, x, right2)

↗
Runge-Kutta method



Carle Runge





Martin Wilhelm Kutta

Initial-value problem for a system of two first-order equations

Problem: find a solution $y(x)$, $z(x)$ of the system

$$\begin{cases} y' = f(x, y, z) \\ z' = g(x, y, z) \end{cases}$$

which satisfies the initial condition $y(a) = y_0$, $z(a) = z_0$,
where functions f and g are given in the domain
 $a \leq x \leq b$, $-\infty < y, z < \infty$.



$$y' = f(x, y, z)$$

$$z' = g(x, y, z)$$

Euler's method:

$$\Delta y / \Delta x = f(x_i, y_i, z_i)$$

$$\Delta z / \Delta x = g(x_i, y_i, z_i)$$

$$(y_{i+1} - y_i) / h = f(x_i, y_i, z_i) \quad (z_{i+1} - z_i) / h = g(x_i, y_i, z_i)$$

$$i = 0, 1, 2, \dots, n-1$$

$$y_{i+1} = y_i + h f(x_i, y_i, z_i) \quad z_{i+1} = z_i + h g(x_i, y_i, z_i)$$

Example.

$$y' = z(x+y) \quad y(0)=0$$

$$z' = zx+3 \quad z(0)=0$$

$$x_1 = h$$

$$y_1 = 0 + h \cdot 0 = 0$$

$$z_1 = 0 + h \cdot 3 = 3h$$

$$x_2 = 2h$$

$$y_2 = y_1 + h f(x_1, y_1, z_1) = 0 + h 3h(h+0) = 3h^3$$

$$z_2 = z_1 + h g(x_1, y_1, z_1) = 3h + h(3hh+3) = 6h + 3h^3$$

Initial-value problem (Cauchy's problem) for a second-order differential equation

Formulation of the problem: find a solution $y(x)$ of equation

$$y''=f(x, y, y')$$

$$d^2y/dx^2=f(x, y, dy/dx)$$

which satisfies the initial condition $y(a)=y_0$, $y'(a)=y'_0$

where function f is given at $a \leq x \leq b$, $-\infty < y, y' < \infty$

Solution:

Let us denote $y'(x)=z(x)$.

Then the problem transforms to the above considered problem for the system of 2 equations:

$$\begin{cases} z' = f(x, y, z) \\ y' = z \end{cases}$$

with initial data:

$$z(a)=y'_0, \quad y(a)=y_0$$

Initial-value problem (Cauchy's problem) for a system of 2 second-order differential equations

$$\begin{cases} y''=f(x, y, y', z, z') \\ z''=g(x, y, y', z, z') \end{cases}$$

We introduce 2 new functions:

$$u(x)=y'(x) , \quad v(x)=z'(x)$$

and obtain the system of 4 first-order equations

$$\left\{ \begin{array}{l} u' = f(x, y, u, z, v) \\ v' = g(x, y, u, z, v) \\ y' = u \\ z' = v \end{array} \right.$$

**Example: rotation of a
small mass in a gravity field**

We denote by t (instead of x) the independent variable, which is time, in the next example.

Equations of mass rotation in the plane (y,z) :

$$\begin{cases} d^2y/dt^2 = -ky / (y^2+z^2)^{3/2} \\ d^2z/dt^2 = -kz / (y^2+z^2)^{3/2} \end{cases} \quad \text{where} \quad k=5+0.1 \cdot t$$

The system transforms to

$$\begin{cases} dy/dt = u \\ dz/dt = v \\ du/dt = -ky / (y^2+z^2)^{3/2} \\ dv/dt = -kz / (y^2+z^2)^{3/2} \end{cases}$$

Initial data: $y=0, z=1, u=1, v=1$ at $t=0$

```
clear
```

```
function f=right3(t, z)
```

```
// z(1)=y   z(2)=z   z(3)=u   z(4)=v
```

```
    k=5+0.1*t
```

```
    a=z(1)*z(1)+z(2)*z(2) ;
```

```
    b=a*sqrt(a) ;
```

```
f(1) =z(3)      ;    // right hand side of eq. 1
```

```
f(2) =z(4)      ;    // right hand side of eq. 2
```

```
f(3) = -k*z(1)/b ;    // right hand side of eq. 3
```

```
f(4) = -k*z(2)/b ;    // right hand side of eq. 4
```

```
endfunction
```

```
t=0 : 0.002 : 15
```

```
z=ode([0;1; 1;1], 0, t, right3) ;
```

```
comet( z(1,:), z(2,:))
```

Boundary-value problems for a second-order differential equation

Problem: find a solution $y(x)$ of equation

$$y''=f(x, y, y')$$

at $a \leq x \leq b$ that satisfies the boundary conditions


$$y(a)=y_a, \quad y(b)=y_b \quad \text{at opposite ends of } [a,b]$$

Shooting method

Let us assign some $y'(a)$ and solve the problem with initial data

$$y(a)=y_a, \quad y'(a)$$

As a result, we obtain some y_b at $x=b$.



Compare obtained y_n with given y_b , and calculate $|y_n - y_b|$.

We correct the value of $y'(a)$, again solve the initial value problem and in this way usually choose $y'(a)$ so as to provide $|y_n - y_b| < \varepsilon$.

Example 1. $y'' = -y - 1, \quad y(0) = y(1) = 0$

In some cases, however, the solution of boundary-value problem does not exist.

Example 2. $y'' = (y')^2 + 1 \quad 0 \leq x \leq 4 \quad y(0) = 0 \quad y(4) = 1$

(why does not Theorem 1 work in this case?)

In a particular case of the linear equation

$$y'' + p(x)y' + q(x)y = f(x) \quad (*)$$

$$y(a) = y_a, \quad y(b) = y_b$$

(where p, q, f – given functions) **the solution certainly exist and it is unique if $q \leq 0$ (see a course of ode).**

Moreover, this solution can be calculated using only two shooting. Indeed, it can be represented in the form

$$y(x) = \hat{y}(x) + C \cdot \tilde{y}(x)$$

where \hat{y} – solution of equation (*) with initial condition

$$\hat{y}(a) = y_a, \quad \hat{y}'(a) = 0 \quad (\text{first shooting});$$

\tilde{y} - solution of equation (*) with $f=0$ and initial condition $\tilde{y}(a)=0, \quad \tilde{y}'(a)=1$ (second shooting),

$$C = [y_b - \hat{y}(b)] / \tilde{y}(b)$$

Finite-Difference Method of solving boundary-value problem for the linear second-order differential equation (by transition to algebraic equations)

$$y'' + p(x)y' + q(x)y = f(x) \quad (1)$$

$$y(a) = y_a, \quad y(b) = y_b \quad (2)$$

By dividing the segment $a \leq x \leq b$ into n subsegments, we obtain as usual the nodal points

$$x_0 = a, \quad x_1, \quad x_2, \dots, \quad x_n = b, \quad h = (b - a)/n$$

We can write down equation (*) at inner points, and replace y'' , y' by $y(x_{i-1})$, $y(x_i)$, $y(x_{i+1})$ according to formulae discussed in Chapter 11:

$$\begin{aligned} & [y(x_{i+1}) - 2y(x_i) + y(x_{i-1})] / h^2 + O(h^2) + \\ & + p(x_i)[y(x_{i+1}) - y(x_{i-1})] / (2h) + O(h^2) + \\ & + q(x_i) y(x_i) = f(x_i) \quad (3) \end{aligned}$$

at $i=1, 2, \dots, n-1$.

If $y(x)$ is the solution of problem (1),(2), then it must satisfy expression (3).

Now, omitting errors $O(h^2)$, we get the system of algebraic equations for the approximate solution y_1, y_2, \dots, y_{n-1} :

$$\begin{aligned} & (y_{i+1} - 2y_i + y_{i-1})/h^2 + \\ & \quad + p(x_i)(y_{i+1} - y_{i-1})/(2h) + \\ & \quad + q(x_i) y_i = f(x_i) \end{aligned} \tag{4}$$

at $i=1, 2, \dots, n-1$.

We notice that $y_0=y_a$ and $y_n=y_b$ are known due to boundary conditions (2).

Multiplying (4) by h^2 , we obtain:

$$\begin{aligned} y_{i+1} - 2y_i + y_{i-1} + \\ + p(x_i)(y_{i+1} - y_{i-1})h/2 + \\ + h^2 q(x_i) y_i = h^2 f(x_i) \end{aligned}$$

This system can be transformed to standard form:

$$a_{11} y_1 + a_{12} y_2 = b_1 ,$$

$$a_{21} y_1 + a_{22} y_2 + a_{23} y_3 = b_2 ,$$

$$a_{32} y_2 + a_{33} y_3 + a_{34} y_4 = b_3 ,$$

$$a_{n-2,n-3} y_{n-3} + a_{n-2,n-2} y_{n-2} + a_{n-2,n-1} y_{n-1} = b_{n-2}$$

$$a_{n-1,n-2} y_{n-2} + a_{n-1,n-1} y_{n-1} = b_{n-1}$$

where a_{ij} are known coefficients,

y_i are unknowns

Usually this system can be solved using Gaussian's elimination, see Chapter 3:

$$a_{11} \mathbf{y}_1 + a_{12} \mathbf{y}_2 = b_1 ,$$

$$\tilde{a}_{22} \mathbf{y}_2 + a_{23} \mathbf{y}_3 = \beta_2 ,$$

$$\tilde{a}_{33} \mathbf{y}_3 + a_{34} \mathbf{y}_4 = \beta_3 ,$$

$$\tilde{a}_{n-2, n-2} \mathbf{y}_{n-2} + a_{n-2, n-1} \mathbf{y}_{n-1} = \beta_{n-2}$$

$$\tilde{a}_{n-1, n-1} \mathbf{y}_{n-1} = \beta_{n-1}$$

Other boundary conditions which can occur in practice:

$$(+) \quad y(a) = c_1, \quad y'(b) = c_2$$

$$(++) \quad y'(a) = c_3, \quad y(b) = c_4$$

$$(+++) \quad y(a) = c_5, \quad y(b) + c_6 y'(b) = c_7$$

where values of $c_1, c_2, c_3, c_4, c_5, c_6, c_7$ are given.