

# Differential Geometry. Home Assignments

## Assignment 1

### Problem 1

A circle with radius  $r$  that rolls without slipping along the outer circumference of the fixed circle with radius  $R$  is given. A point  $M$  is fixed on the circumference of the rolling circle.

Determine the equation describing the curve being traced by point  $M$ . Find singular points on this curve or justify that it has no singular points.

### Problem 2

Prove the assertion below.

Suppose

$$\begin{cases} x = x(t), \\ y = y(t), \\ z = z(t) \end{cases}$$

is an arbitrary parametrisation of an elementary curve. Then any other parametrisation has the form

$$\begin{cases} x = x(\sigma(\tau)), \\ y = y(\sigma(\tau)), \\ z = z(\sigma(\tau)) \end{cases}$$

where  $\sigma(\tau)$  is a continuous, strictly monotonic function.

### Problem 3

Find the conditions for the existence of an asymptote to the space curve

$$\begin{cases} x = x(t), \\ y = y(t), \\ z = z(t), \end{cases} \quad t \in (a, b),$$


which tends to infinity as  $t \rightarrow a$ .

# HW 1 Week 11th.

## Problem 1

A circle with radius  $r$  that rolls without slipping along the outer circumference of the fixed circle with radius  $R$  is given. A point  $M$  is fixed on the circumference of the rolling circle.

Determine the equation describing the curve being traced by point  $M$ . Find singular points on this curve or justify that it has no singular points.



Consider  $M_0$  at  $(R+r, 0)$  with counter-clockwise rolling

during the rolling process, denote the change of central angle of fixed circle by  $\varphi$ .

No slipping  $\Rightarrow R\varphi = r\psi$  the change of central angle of rolling circle by  $\psi$ .

denote  $P: (R+r)\cos\varphi, (R+r)\sin\varphi$ .

thus  $M: (R+r)\cos\varphi + r\cos(\psi+\varphi), (R+r)\sin\varphi + r\sin(\psi+\varphi)$ .

then parametrize  $\gamma_M: \begin{cases} x(\varphi) = (R+r)\cos\varphi + r\cos(\frac{R+r}{r}\varphi) \\ y(\varphi) = (R+r)\sin\varphi + r\sin(\frac{R+r}{r}\varphi) \end{cases}$

find singular point:  $x'(\varphi) = -(R+r)\sin\varphi - (R+r)\sin(\frac{R+r}{r}\varphi)$

$y'(\varphi) = (R+r)\cos\varphi + (R+r)\cos(\frac{R+r}{r}\varphi)$ .

$$\begin{aligned} [x'(\varphi)]^2 + [y'(\varphi)]^2 &= 2(R+r) + 2(R+r) [\sin\varphi \sin(\frac{R+r}{r}\varphi) + \cos\varphi \cos(\frac{R+r}{r}\varphi)] \\ &= 2(R+r) + 2(R+r) [\cos(\frac{R+r}{r}\varphi - \varphi)] = 2(R+r) (1 - \cos\frac{R}{r}\varphi) \end{aligned}$$

which = 0 iff  $\frac{R}{r}\varphi = \frac{\pi}{2} + 2k\pi \Rightarrow \varphi = \frac{\pi r}{2R} + \frac{2k\pi r}{R}$ .

thus, when  $\varphi = \frac{\pi r}{2R} + \frac{2k\pi r}{R}$ ,  $\gamma_M$  has singular points.

$$\left( (R+r)\cos\left(\frac{\pi r}{2R} + \frac{2k\pi r}{R}\right) - r\sin\left(\frac{\pi r}{2R} + \frac{2k\pi r}{R}\right), (R+r)\sin\left(\frac{\pi r}{2R} + \frac{2k\pi r}{R}\right) + r\cos\left(\frac{\pi r}{2R} + \frac{2k\pi r}{R}\right) \right)$$

Prove the assertion below.

Suppose

$$\gamma: \begin{cases} x = x(t), \\ y = y(t), \\ z = z(t) \end{cases}$$

is an arbitrary parametrisation of an elementary curve. Then any other parametrisation has the form

$$\begin{cases} x = x(\sigma(\tau)), \\ y = y(\sigma(\tau)), \\ z = z(\sigma(\tau)) \end{cases}$$

where  $\sigma(\tau)$  is a continuous, strictly monotonic function.

Pf: Assume there exists another parametrization of  $\gamma$ .  $\begin{cases} x = x(\sigma) \\ y = y(\sigma) \\ z = z(\sigma) \end{cases}$ . Since  $\gamma$  is elementary,  $\sigma \in (\alpha, \beta)$ .

also since  $\gamma$  is elementary,  $t \in (a, b)$ , denote map.  $\sigma: (a, b) \rightarrow (\alpha, \beta)$

(by elementary),  $x, y, z$  and  $x^{-1}, y^{-1}, z^{-1}$  are top. map.  $\sigma = x^{-1} \circ x = y^{-1} \circ y = z^{-1} \circ z$  is topological map.

A topological map in  $\mathbb{R}$  is continuous and monotonic.

by the elementarity of  $\gamma$ ,  $\exists \tau_1, \tau_2 \in (a, b)$  s.t.  $\sigma(\tau_1) = \sigma(\tau_2)$ , thus,  $\sigma$  is strictly monotonic.

### Problem 3

Find the conditions for the existence of an asymptote to the space curve

$$\begin{cases} x = x(t), \\ y = y(t), \\ z = z(t), \end{cases} \quad t \in (a, b),$$

which tends to infinity as  $t \rightarrow a$ .

Sol: the asymptote (straight line in space) has expression :  $\frac{x - x_0}{dx} = \frac{y - y_0}{dy} = \frac{z - z_0}{dz} = k$ .

parametrize:  $\begin{cases} x = dx t + x_0 \\ y = dy t + y_0 \\ z = dz t + z_0 \end{cases}$  we need both direction vector  $(dx, dy, dz)$  and initial point exists.

that is,  $\begin{cases} \lim_{t \rightarrow a} \frac{x(t)}{t} = dx < \infty \\ \lim_{t \rightarrow a} \frac{y(t)}{t} = dy < \infty \\ \lim_{t \rightarrow a} \frac{z(t)}{t} = dz < \infty \end{cases}$  and  $\begin{cases} \lim_{t \rightarrow a} (x(t) - dx t) = x_0 < \infty \\ \lim_{t \rightarrow a} (y(t) - dy t) = y_0 < \infty \\ \lim_{t \rightarrow a} (z(t) - dz t) = z_0 < \infty \end{cases}$

Prove the assertion below.

Suppose

$$\gamma: \begin{cases} x = x(t), \\ y = y(t), \\ z = z(t) \end{cases}$$

is an arbitrary parametrisation of an elementary curve. Then any other parametrisation has the form

$$\begin{cases} x = x(\sigma(\tau)), \\ y = y(\sigma(\tau)), \\ z = z(\sigma(\tau)) \end{cases}$$

where  $\sigma(\tau)$  is a continuous, strictly monotonic function.

Pf: denote  $t = \sigma(\tau)$ . by the definition of elementary function.  $\sigma$  should be a topological mapping.

That is,  $\sigma$  is bijection. and  $\sigma, \sigma^{-1}$  is continuous

It suffices to check  $\sigma$  is strictly monotonic.

assume the converse.  $\exists \tau_1 < \tau_2 < \tau_3$ . s.t.  $\sigma(\tau_2) \geq \max(\sigma(\tau_1), \sigma(\tau_3))$  . or  $\leq \min(\sigma(\tau_1), \sigma(\tau_3))$ .

w.l.g.  $\sigma(\tau_2) \geq \max(\sigma(\tau_1), \sigma(\tau_3))$

by the continuity and mean value theorem,  $\exists \tau_1' \in (\tau_1, \tau_2)$ .  $\tau_2' \in (\tau_2, \tau_3)$  s.t.  $\max(\sigma(\tau_1), \sigma(\tau_3)) \leq \sigma(\tau_1') = \sigma(\tau_2') \leq \sigma(\tau_2)$ .

which contradicts with the injective of  $\sigma$ .

Thus,  $\sigma$  is cont. and strictly monotonic.

II.

### Problem 3

Find the conditions for the existence of an asymptote to the space curve

$$\begin{cases} x = x(t), \\ y = y(t), \\ z = z(t), \end{cases} \quad t \in (a, b),$$

which tends to infinity as  $t \rightarrow a$ .

Sol: denote  $\vec{r}(t) = (x(t), y(t), z(t))$ .

Assume there exist a asymptote with direction vector  $\vec{a}$  and initial point  $\vec{b}$ .

then the asymptote could be parametrised:  $\vec{r} = \vec{a}u + \vec{b} \Rightarrow \begin{cases} x(u) = x_0 + a_x u \\ y(u) = y_0 + a_y u \\ z(u) = z_0 + a_z u \end{cases}$

$$(*) \begin{cases} \lim_{t \rightarrow a} \frac{\vec{r}(t)}{\|\vec{r}(t)\|} = \vec{a} = (a_x, a_y, a_z). & \text{- finite} \\ \lim_{t \rightarrow a} [\vec{r}(t) - (\vec{r}(t) \cdot \vec{a}) \cdot \vec{a}] = \vec{b} = (x_0, y_0, z_0) & \text{- finite point} \end{cases}$$

That is if the condition (\*) holds. ( i.e. the direction limit exists and orthogonal projection converges to finite point ).

the curve exists asymptote.