

Core Concepts and Theorems in Classical Optimization Theory

(Complete Compilation of Part I: Optimization Theory)

Compiled by Gemini AI Assistant

November 2025

Contents

1	Unconstrained Optimization Concepts and Theorems	3
1.1	Unconstrained Optimization Concepts (Definitions)	3
1.2	Relevant Theorems	3
2	Line-Search Concepts and Theorems	4
2.1	Line-Search Concepts (Definitions)	4
2.2	Relevant Results	4
3	Newton and Quasi-Newton Methods (Statements)	5
3.1	Core Concepts	5
3.2	Relevant Theorems/Lemmas	5
4	Trust-Region Methods	6
4.1	Core Concepts	6
4.2	Relevant Results	6
5	Conjugate Gradient and Conjugate Directions	6
5.1	Core Concepts	6
5.2	Relevant Theorems	7
6	Constrained Optimization	8
6.1	Core Definitions	8
7	Lagrangian, KKT, and First-Order Optimality	8
7.1	Core Concepts	8
7.2	Relevant Items in the Notes	8
8	Second-Order Optimality for Constrained Problems	9
8.1	Core Concepts	9
8.2	Relevant Theorems	9
9	Duality Theory (Nonlinear and LP)	10
9.1	Core Concepts	10
9.2	Relevant Theorems	10
10	Linear Programming and the Simplex Method (Concepts)	11
10.1	Core Concepts	11
10.2	Relevant Results	11
11	Quadratic Programming (QP)	11
11.1	Core Concepts	11
11.2	Relevant Theorems	11

12 Descriptive Regression and Least Squares Estimation (LSE)	12
12.1 Descriptive Regression General Setting	12
12.2 Parametric Regression and Least Squares Formulation	12
12.3 Least Squares Estimation (LSE)	12
13 Classical Linear Regression Model and BLUE	13
13.1 Classical Linear Regression Model	13
13.2 Best Linear Unbiased Estimator (BLUE)	13
14 OLS Estimation in the Singular Case	14
14.1 OLS Estimation in the Singular Case	14
15 WLS/GLS and Design of Experiments	15
15.1 Weighted and Generalized Least Squares (WLS / GLS)	15
15.2 Design of Experiments: Basic Concepts	15
15.3 Information Matrix and Variance of Estimates	15
16 Optimality Criteria and Equivalence Theorem	16
16.1 Optimality Criteria in Design	16

1 Unconstrained Optimization Concepts and Theorems

1.1 Unconstrained Optimization Concepts (Definitions)

1. **Unconstrained Optimization Problem** The problem of finding $\min_{x \in \mathbb{R}^n} f(x)$ or $f(x) \rightarrow \min_{x \in \mathbb{R}^n}$, where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a smooth (continuously differentiable) scalar function.
2. **Global Minimum** A point $x^* \in \mathbb{R}^n$ is a global minimum of f if and only if $f(x^*) \leq f(x)$ holds for all $x \in \mathbb{R}^n$.
3. **Local Minimum** A point $x^* \in \mathbb{R}^n$ is a local minimum of f if and only if there exists $\epsilon > 0$ such that $f(x^*) \leq f(x)$ holds for all x satisfying $\|x - x^*\| < \epsilon$.
4. **Strict Local Minimum** A vector $x^* \in \mathbb{R}^n$ is a strict local solution if there exists a neighborhood \mathcal{N} such that $f(x) > f(x^*)$ holds for all x satisfying $x \in \mathcal{N} \cap \Omega, x \neq x^*$.
5. **Stationary Point** A point $x^* \in \mathbb{R}^n$ is a stationary point if and only if $\nabla f(x^*) = 0$.
6. **Descent Direction** For a function f and a point x , a vector $p \in \mathbb{R}^n$ is called a descent direction if it satisfies $\nabla f(x)^T p < 0$.
7. **Gradient** (∇f) ∇f is the gradient vector of the function f .
8. **Hessian** ($\nabla^2 f$) $\nabla^2 f$ is the Hessian matrix of the function f .
9. **Positive Definiteness** A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is positive definite if and only if $p^T A p > 0$ for all $p \neq 0$.
10. **Positive Semidefiniteness** A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is positive semidefinite if and only if $p^T A p \geq 0$ for all $p \in \mathbb{R}^n$.

1.2 Relevant Theorems

Theorem 1.1 (Taylor's Theorem). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable, and $p \in \mathbb{R}^n$. Then there exists $t \in (0, 1)$ such that:*

$$f(x + p) = f(x) + \nabla f(x + tp)^T p$$

If f is twice continuously differentiable, then there exists $t \in (0, 1)$ such that:

$$f(x + p) = f(x) + \nabla f(x)^T p + \frac{1}{2} p^T \nabla^2 f(x + tp) p$$

Theorem 1.2 (Necessary Condition for a Local Minimum). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable in an open neighborhood of x^* . If x^* is a local minimum of f , then:*

$$\nabla f(x^*) = 0$$

Theorem 1.3 (Second-Order Necessary Condition). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable in an open neighborhood of x^* . If x^* is a local minimum of f , then:*

$$\nabla f(x^*) = 0 \quad \text{and} \quad \nabla^2 f(x^*) \text{ is positive semidefinite.}$$

Theorem 1.4 (Second-Order Sufficient Condition). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable in an open neighborhood of x^* . Suppose:*

$$\nabla f(x^*) = 0 \quad \text{and} \quad \nabla^2 f(x^*) \text{ is positive definite.}$$

Then x^ is a strict local minimum of f .*

Theorem 1.5 (Global Minimum for Convex Functions). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex. Then any local minimizer x^* is also a global minimizer. If in addition f is differentiable, then any stationary point x^* with $\nabla f(x^*) = 0$ is a global minimizer.*

2 Line-Search Concepts and Theorems

2.1 Line-Search Concepts (Definitions)

1. **Descent Direction** For a function f and a point x , a vector $p \in \mathbb{R}^n$ is called a descent direction if it satisfies $\nabla f(x)^T p < 0$.
2. **Step Length (α)** In the iteration $x_{k+1} = x_k + \alpha_k p_k$, $\alpha_k > 0$ is the distance moved along the search direction p_k .
3. **Sufficient Decrease Condition (Armijo Condition)** A step length $\alpha > 0$ satisfies the sufficient decrease condition if and only if:

$$f(x + \alpha p) \leq f(x) + c_1 \alpha \nabla f(x)^T p$$

where $c_1 \in (0, 1)$.

4. **Curvature Condition** A step length $\alpha > 0$ satisfies the curvature condition if and only if:

$$\nabla f(x + \alpha p)^T p \geq c_2 \nabla f(x)^T p$$

where $c_2 \in (c_1, 1)$.

5. **Wolfe Conditions** A step length $\alpha > 0$ satisfies the Wolfe conditions if and only if it satisfies both:

$$(a) \text{ Sufficient decrease: } f(x_k + \alpha p_k) \leq f(x_k) + c_1 \alpha \nabla f(x_k)^T p_k$$

$$(b) \text{ Curvature condition: } \nabla f(x_k + \alpha p_k)^T p_k \geq c_2 \nabla f(x_k)^T p_k$$

where $0 < c_1 < c_2 < 1$.

6. **Strong Wolfe Conditions** A step length $\alpha > 0$ satisfies the strong Wolfe conditions if and only if it satisfies the sufficient decrease condition (1) and replaces the curvature condition (2) with:

$$|\nabla f(x_k + \alpha p_k)^T p_k| \leq c_2 |\nabla f(x_k)^T p_k|$$

where $0 < c_1 < c_2 < 1$.

7. **Goldstein Conditions** A step length $\alpha > 0$ satisfies the Goldstein conditions if and only if:

$$f(x_k) + (1 - c) \alpha \nabla f(x_k)^T p_k \leq f(x_k + \alpha p_k) \leq f(x_k) + c \alpha \nabla f(x_k)^T p_k$$

where $c \in (0, 0.5)$.

2.2 Relevant Results

Lemma 2.1 (Existence of Step Lengths). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable. Suppose p_k is a descent direction at x_k , and f is bounded below along the ray $\{x_k + \alpha p_k | \alpha > 0\}$. Then, for any $0 < c_1 < c_2 < 1$, there exists an interval of step lengths α such that the Wolfe and strong Wolfe conditions are satisfied.*

Theorem 2.2 (Global Convergence of Line Search Methods). *Let $\{x_k\}$ be a sequence generated by the iteration $x_{k+1} = x_k + \alpha_k p_k$ where p_k is a descent direction, and the step length α_k satisfies Wolfe conditions. Assume:*

1. *The function f is bounded below on \mathbb{R}^n .*
2. *f is continuously differentiable in an open set \mathcal{N} containing the level set $\mathcal{L} = \{x \in \mathbb{R}^n | f(x) \leq f(x_0)\}$, where x_0 is the starting point of the iteration.*
3. *The gradient ∇f is Lipschitz continuous on \mathcal{N} , i.e., $\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\|$ for all $x, y \in \mathcal{N}$.*

Then the Zoutendijk condition holds:

$$\sum_{k \geq 0} (\cos \theta_k)^2 \|\nabla f(x_k)\|^2 < \infty$$

where $\cos \theta_k = \frac{-\nabla f(x_k)^T p_k}{\|\nabla f(x_k)\| \|p_k\|}$.

3 Newton and Quasi-Newton Methods (Statements)

3.1 Core Concepts

1. **Newton Step Definition** The search direction for Newton's method is defined as $p_k^N = -(\nabla^2 f(x_k))^{-1} \nabla f(x_k)$.
2. **Local Quadratic Convergence Statement** If the initial point x_0 is sufficiently close to x^* (where $\nabla f(x^*) = 0$ and $\nabla^2 f(x^*) \succ 0$), the sequence of iterates $\{x_k\}$ generated by the pure Newton method converges to x^* at a Q-quadratic rate, i.e., $\|x_{k+1} - x^*\| \leq M \|x_k - x^*\|^2$ for some constant M .
3. **The Hessian Modification Idea** If the Hessian $\nabla^2 f(x_k)$ is not positive definite, it is replaced by a modified matrix $B_k = \nabla^2 f(x_k) + E_k$, where E_k is chosen such that B_k is positive definite, ensuring that $p_k = -B_k^{-1} \nabla f(x_k)$ is a descent direction.
4. **Quasi-Newton Secant Condition** The updated Quasi-Newton matrix B_{k+1} must satisfy the secant equation: $B_{k+1} s_k = y_k$, where $s_k = x_{k+1} - x_k$ and $y_k = \nabla f_{k+1} - \nabla f_k$.

3.2 Relevant Theorems/Lemmas

Theorem 3.1 (Local Convergence of Newton's Method). *Suppose f is twice differentiable and that $\nabla^2 f(x)$ is Lipschitz continuous near a solution x^* satisfying the second-order sufficient conditions. Consider the iteration $x_{k+1} = x_k + p_k$ where p_k is the Newton step. Then:*

1. *If the starting point x_0 is sufficiently close to x^* , the sequence of iterates converges to x^* .*
2. *The convergence rate of $\{x_k\}$ is quadratic.*
3. *The sequence $\{\|\nabla f_k\|\}$ converges quadratically to zero.*

Theorem 3.2 (Quasi-Newton / Superlinear Convergence Statements). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable. Let the iteration $x_{k+1} = x_k + \alpha_k p_k$ be generated by a descent direction p_k and a step length α_k satisfying Wolfe conditions (with $c_1 \leq \frac{1}{2}$). If the sequence $\{x_k\}$ converges to x^* such that $\nabla f(x^*) = 0$ and $\nabla^2 f(x^*)$ is positive definite, and if the search direction satisfies:*

$$\lim_{k \rightarrow \infty} \frac{\|\nabla f(x_k) + \nabla^2 f(x_k) p_k\|}{\|p_k\|} = 0$$

then:

1. *The step length $\alpha_k = 1$ is admissible for all k greater than a certain index k_0 .*
2. *If $\alpha_k = 1$ for all $k > k_0$, then $\{x_k\}$ converges to x^* superlinearly.*

4 Trust-Region Methods

4.1 Core Concepts

1. **The Quadratic Trust-Region Model** The model is $m_k(p) = f_k + g_k^T p + \frac{1}{2} p^T B_k p$ minimized within the constraint $\|p\| \leq \Delta_k$.
2. **Trust-Region Radius Strategy** The strategy chooses the trust-region radius Δ_k based on the agreement between the model m_k and the objective f , using the ratio $\rho_k = \frac{f(x_k) - f(x_k + p_k)}{m_k(0) - m_k(p_k)}$.
3. **The Cauchy Point (Closed-Form)** The Cauchy point p_k^c is defined as $-\tau_k \frac{\Delta_k}{\|g_k\|} g_k$, where:

$$\tau_k = \begin{cases} 1, & \text{if } g_k^T B_k g_k \leq 0; \\ \min\left(\frac{\|g_k\|^3}{\Delta_k g_k^T B_k g_k}, 1\right), & \text{otherwise.} \end{cases}$$

4. **Dogleg Path Idea** The Dogleg Method constructs a piecewise linear path from the origin to the unconstrained minimum along the steepest descent direction (p^U), and then toward the full Newton step (p^B), to approximate the trust-region solution.

4.2 Relevant Results

Lemma 4.1 (Existence of Global Minimum of Unconstrained Quadratic Model). *Let $m(p) = g^T p + \frac{1}{2} p^T B p$, where B is symmetric.*

1. m attains a minimum if and only if B is positive semidefinite and g is in the range of B . If B is positive semidefinite, then every p satisfying $Bp = -g$ is a global minimizer of m .
2. m has a unique minimizer if and only if B is positive definite.

Lemma 4.2 (Dogleg Properties). *Let B be positive definite. Then along the dogleg path $\tilde{p}(\tau)$:*

1. $\|\tilde{p}(\tau)\|$ is an increasing function of τ .
2. $m(\tilde{p}(\tau))$ is a decreasing function of τ .

Lemma 4.3 (Cauchy Decrease). *The Cauchy point p_k^c satisfies the bound:*

$$m_k(0) - m_k(p_k^c) \geq \frac{1}{2} \|g_k\| \min\left(\Delta_k, \frac{\|g_k\|}{\|B_k\|}\right)$$

Theorem 4.4 (Characterization of Trust-Region Solution / Moré-Sorensen Type Result). *The vector p^* is a global solution of the trust-region subproblem $\min_{p \in \mathbb{R}^n} m(p)$ s.t. $\|p\| \leq \Delta$, if and only if p^* is feasible and there exists a scalar $\lambda \geq 0$ such that:*

1. $(B + \lambda I)p^* = -g$
2. $\lambda(\Delta - \|p^*\|) = 0$
3. $B + \lambda I$ is positive semidefinite

5 Conjugate Gradient and Conjugate Directions

5.1 Core Concepts

1. **Definition of A-conjugacy** A set of nonzero vectors $\{p_0, p_1, \dots, p_l\}$ is conjugate with respect to an $n \times n$ symmetric positive definite matrix A if it satisfies $p_i^T A p_j = 0$ for all $i \neq j$.
2. **Krylov Subspaces** A Krylov subspace $\mathcal{K}(r_0; k)$ is defined as $\text{span}\{r_0, A r_0, \dots, A^{k-1} r_0\}$.
3. **Finite Termination Property** If the matrix A has only r distinct eigenvalues, the Conjugate Gradient (CG) iteration will terminate at the solution in at most r iterations.

4. **Orthogonality of Residuals** The sequence of residuals $\{r_k\}$ generated by the CG method satisfies $r_k^T r_i = 0$ for $i \neq k$, i.e., the residuals are mutually orthogonal.
5. **Spectral Convergence Bounds** The convergence rate bounds depend on the distribution of eigenvalues of the matrix A .
6. **Preconditioning Concept** Preconditioning involves transforming the original system $Ax = b$ into an equivalent system with a better-conditioned matrix (e.g., $M^{-1}A$ or $C^{-T}AC^{-1}$) to accelerate the convergence rate of CG.

5.2 Relevant Theorems

Theorem 5.1 (Convergence of Conjugate Direction Method). *For any $x_0 \in \mathbb{R}^n$, the sequence $\{x_k\}$ generated by the conjugate direction algorithm converges to the solution x^* of the linear system $Ax = b$ in at most n steps.*

Theorem 5.2 (Finite-Termination Property). *If A has only r distinct eigenvalues, then the Conjugate Gradient (CG) iteration will terminate at the solution in at most r iterations.*

6 Constrained Optimization

6.1 Core Definitions

1. **Problem Statement** Minimize $f(x)$ over $x \in \mathbb{R}^n$ subject to $c_i(x) = 0, i \in E$ (equality constraints) and $c_i(x) \geq 0, i \in I$ (inequality constraints).
2. **Feasible Set (Ω)** The feasible set Ω is the set of all vectors $x \in \mathbb{R}^n$ satisfying all constraints.
3. **Feasible Point** A feasible point is any vector $x \in \mathbb{R}^n$ belonging to the feasible set Ω .
4. **Active Set ($A(x)$)** The active set $A(x)$ at a feasible point x consists of the equality constraint indices E together with the indices of the inequality constraints $i \in I$ for which $c_i(x) = 0$.
5. **Active Constraint** An inequality constraint $i \in I$ is active at x if $c_i(x) = 0$. All equality constraints $i \in E$ are always active.
6. **Inactive Constraint** An inequality constraint $i \in I$ is inactive at x if the strict inequality $c_i(x) > 0$ is satisfied.
7. **Feasible Sequence** A sequence $\{z_k\}$ approaching x is called a feasible sequence if $z_k \in \Omega$ for all k sufficiently large and $z_k \rightarrow x$.
8. **Tangent Cone ($T_\Omega(x)$)** The set of all limiting directions d of feasible sequences approaching x , where $d = \lim_{k \rightarrow \infty} \frac{z_k - x}{t_k}$ for some feasible sequence $\{z_k\}$ and positive scalars $\{t_k\}$ with $t_k \rightarrow 0$.
9. **Set of Linearized Feasible Directions ($F(x)$)** The set of vectors d satisfying:

$$F(x) = \{d \mid d^T \nabla c_i(x) = 0, \text{ for all } i \in E; \quad d^T \nabla c_i(x) \geq 0, \text{ for all } i \in A(x) \cap I\}$$

10. **LICQ (Linear Independence Constraint Qualification)** LICQ holds at point x with active set $A(x)$ if the set of active constraint gradients $\{\nabla c_i(x), i \in A(x)\}$ is linearly independent.
11. **MFCQ (Mangasarian-Fromovitz Constraint Qualification)** MFCQ holds, if there exists $w \in \mathbb{R}^n$ such that $\nabla c_i(x^*)^T w > 0$ for all $i \in A(x^*) \cap I$, $\nabla c_i(x^*)^T w = 0$ for all $i \in E$, and the set $\{\nabla c_i(x^*), i \in E\}$ is linearly independent.

7 Lagrangian, KKT, and First-Order Optimality

7.1 Core Concepts

1. **The Lagrangian Function** The Lagrangian is defined as $\mathcal{L}(x, \lambda) = f(x) - \sum_{i \in E \cup I} \lambda_i c_i(x)$.
2. **KKT Conditions** The necessary conditions for a local solution x^* (assuming LICQ) require the existence of λ^* satisfying:
 - (a) **Stationarity:** $\nabla_x \mathcal{L}(x^*, \lambda^*) = 0$.
 - (b) **Primal Feasibility:** $c_i(x^*) = 0, i \in E$ and $c_i(x^*) \geq 0, i \in I$.
 - (c) **Dual Feasibility:** $\lambda_i^* \geq 0$ for all $i \in I$.
 - (d) **Complementary Slackness:** $\lambda_i^* c_i(x^*) = 0$ for all $i \in E \cup I$.
3. **Role of Constraint Qualifications for Necessity/Sufficiency** Constraint Qualifications guarantee that the KKT conditions are **necessary** for a local solution.

7.2 Relevant Items in the Notes

Lemma 7.1 (Tangent Cone and First-Order Feasible Directions). $T_\Omega(x^*) \subset F(x^*)$, and if LICQ holds at x^* , then $T_\Omega(x^*) = F(x^*)$.

Theorem 7.2 (First-Order Necessary Conditions). Suppose x^* is a local solution, f and c_i are continuously differentiable, and that LICQ holds at x^* . Then there is a Lagrange multiplier vector λ^* satisfying the KKT conditions.

8 Second-Order Optimality for Constrained Problems

8.1 Core Concepts

1. **Critical Cone Definition** The critical cone $C(x^*, \lambda^*)$ contains directions $w \in F(x^*)$ for which $w^T \nabla f(x^*) = 0$, defined by the conditions in Section 7.2.1.
2. **Second-Order Necessary Conditions** They require $w^T \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) w \geq 0$ for all $w \in C(x^*, \lambda^*)$.
3. **Second-Order Sufficient Conditions** They require $w^T \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) w > 0$ for all $w \in C(x^*, \lambda^*)$, $w \neq 0$.
4. **Normal Cone Definition** The normal cone $N_\Omega(x)$ is defined as $N_\Omega(x) = \{v \mid v^T w \leq 0 \text{ for all } w \in T_\Omega(x)\}$.

8.2 Relevant Theorems

Theorem 8.1 (Second-Order Necessary Conditions). *Suppose x^* is a local solution and that LICQ holds. Let λ^* be a Lagrange multiplier satisfying the KKT conditions. Then $w^T \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) w \geq 0$ for all $w \in C(x^*, \lambda^*)$.*

Theorem 8.2 (Second-Order Sufficient Conditions). *Suppose that for some feasible point x^* , there exists a multiplier λ^* such that the KKT conditions are satisfied. Further suppose that $w^T \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) w > 0$ for all $w \in C(x^*, \lambda^*)$, $w \neq 0$. Then x^* is a strict local solution of the problem.*

9 Duality Theory (Nonlinear and LP)

9.1 Core Concepts

1. **Dual Function** $q(\lambda) = \inf_x [f(x) - \lambda^T c(x)]$.
2. **Weak Duality** Weak duality states that $q(\bar{\lambda}) \leq f(\bar{x})$ for any primal feasible \bar{x} and dual feasible $\bar{\lambda} \geq 0$.
3. **Duality for Convex Quadratic Programming** The dual problem maximizes a concave quadratic objective subject to nonnegativity constraints.
4. **Wolfe Dual (Construction and Interpretation)** The Wolfe dual is $\max_{x, \lambda} \mathcal{L}(x, \lambda)$ s.t. $\nabla_x \mathcal{L}(x, \lambda) = 0$ and $\lambda \geq 0$.

9.2 Relevant Theorems

Theorem 9.1 (Concavity of the Dual Objective). *The dual objective $q(\lambda) = \inf_x \mathcal{L}(x, \lambda)$ is concave, and its domain $D = \{\lambda \mid q(\lambda) > -\infty\}$ is convex.*

Theorem 9.2 (Weak Duality). *For any feasible \bar{x} in $\min_x f(x)$ s.t. $c(x) \geq 0$ and $\bar{\lambda} \geq 0$, $q(\bar{\lambda}) \leq f(\bar{x})$.*

Theorem 9.3 (Solutions of the Dual Problem). *If \bar{x} solves $\min_x f(x)$ s.t. $c(x) \geq 0$, and $f, -c_i$ are convex and differentiable at \bar{x} , then any $\bar{\lambda}$ satisfying the KKT conditions with \bar{x} is a solution of the dual problem.*

Theorem 9.4 (Wolfe Dual Formulation). *If f and $-c_i$ are convex and continuously differentiable, and $(\bar{x}, \bar{\lambda})$ satisfies the KKT conditions, then $(\bar{x}, \bar{\lambda})$ solves the Wolfe Dual $\max_{x, \lambda} \mathcal{L}(x, \lambda)$ s.t. $\nabla_x \mathcal{L}(x, \lambda) = 0, \lambda \geq 0$.* </theorem

10 Linear Programming and the Simplex Method (Concepts)

10.1 Core Concepts

1. **Standard Form of LP**

$$\min c^T x, \quad \text{s.t.} \quad Ax = b, \quad x \geq 0$$

2. **Vertices and Basic Feasible Points** All basic feasible points for the LP problem are vertices of the feasible polytope $\{x \mid Ax = b, x \geq 0\}$, and vice versa.

3. **Statement of the Fundamental Theorem of Linear Programming** See Theorem 10.1.

10.2 Relevant Results

Theorem 10.1 (Fundamental Theorem of Linear Programming). 1. If the primal problem has a nonempty feasible region, then there is at least one basic feasible point.

2. If the primal problem has solutions, then at least one solution is a basic optimal point.

3. If the nonempty feasible region is bounded, then the primal problem has an optimal solution.

11 Quadratic Programming (QP)

11.1 Core Concepts

1. **Definition of a Quadratic Program** A QP minimizes a quadratic objective function $q(x) = \frac{1}{2}x^T Gx + x^T c$ subject to linear constraints.

2. **Equality-Constrained QP** A QP where all constraints are linear equalities $Ax = b$.

3. **KKT system for Equality-Constrained QP** The KKT system is:

$$\begin{pmatrix} G & A^T \\ A & 0 \end{pmatrix} \begin{pmatrix} x^* \\ \lambda^* \end{pmatrix} = \begin{pmatrix} -c \\ b \end{pmatrix}$$

4. **Sufficiency Conditions for Convex QP** For a convex QP (G is positive semidefinite), KKT conditions are sufficient for global optimality.

11.2 Relevant Theorems

Theorem 11.1 (Global Solution for the Equality-Constrained QP). If A has full row rank and $Z^T GZ$ is positive definite (where Z is the null-space basis for A), then x^* satisfying the KKT system is the unique global solution of the equality-constrained QP.

Theorem 11.2 (Sufficiency for Convex QP). If x^* satisfies KKT conditions with λ_i^* , and G is positive semidefinite (convex QP), then x^* is a global solution of the QP.

12 Descriptive Regression and Least Squares Estimation (LSE)

12.1 Descriptive Regression General Setting

1. **Regression Purpose** To represent and understand the relationship between a response (output) and several influencing factors (inputs). Regression analysis is a statistical method for estimating relationships among variables, describing how a dependent variable changes as one or more independent variables change.
2. **Dependent Variable (y)** The outcome or response variable we are trying to predict or explain.
3. **Independent Variables (x_1, \dots, x_m)** The predictor variables or factors that influence the dependent variable. Also called regressors.
4. **Model Function ($\eta(x_1, \dots, x_m)$)** A mathematical representation of the relationship: $y = \eta(x_1, \dots, x_m)$.
5. **Role of Regression Analysis as a Descriptive Modeling Tool** The focus of descriptive regression is to build a model based on observed data without making strong assumptions about the underlying statistical distribution.

12.2 Parametric Regression and Least Squares Formulation

1. **Idea of Model Fitting** The core challenge is that exact functional relationships are often too complex to determine, so we use simplified parametric models $\eta(x, \theta)$ that approximate the statistical dependence with satisfactory accuracy.
2. **Residuals and Error Minimization** The deviation (error) for the j -th observation is $\epsilon_j = y_j - \tilde{y}_j$, where \tilde{y}_j are the predicted values. We seek to minimize the magnitude of the error vector $\epsilon = Y - X\theta$.
3. **Least Squares Criterion and its Justification** The criterion is to minimize the sum of squared errors: $\sum \epsilon_j^2 = \epsilon^T \epsilon = (Y - X\theta)^T (Y - X\theta)$. This is preferred due to its computational simplicity and strong optimality properties (e.g., resulting in a convex optimization problem).

12.3 Least Squares Estimation (LSE)

1. **Definition of the Least Squares Estimator (LSE)** The vector $\hat{\theta} = \arg \min_{\theta \in \mathbb{R}^m} (Y - X\theta)^T (Y - X\theta) = \arg \min_{\theta \in \mathbb{R}^m} \epsilon^T \epsilon$ is called the empirical least squares estimator.
2. **System of Normal Equations** The system of equations $X^T X \theta = X^T Y$ provides a closed-form condition for the optimality of θ .
3. **Convexity of the Least Squares Criterion** The least squares criterion $f(\theta) = (Y - X\theta)^T (Y - X\theta)$ is quadratic in θ , leading to a convex optimization problem.

Lemma 12.1 (Normal Equations Existence and Optimality Condition). For any matrix X and vector Y of compatible dimensions, the system of normal equations $X^T X \theta = X^T Y$ always has at least one solution. Any vector $\hat{\theta}$ satisfying this system is a least squares estimator.

13 Classical Linear Regression Model and BLUE

13.1 Classical Linear Regression Model

1. **Definition of the Classical Linear Regression Model** The model is written as $y_j = \sum_{i=1}^m x_{ji}\theta_i + \epsilon_j$ for $j = 1, \dots, N$, or in matrix form: $Y = X\theta + \epsilon$.
2. **Assumptions of Zero Mean, Uncorrelated and Homoscedastic Errors** The error vector ϵ satisfies:
 - Zero Mean (Unbiasedness): $E[\epsilon_i] = 0$.
 - Homoscedasticity: $E[\epsilon_i^2] = \sigma^2$ (common variance).
 - Uncorrelated Errors: $E[\epsilon_i\epsilon_j] = 0$ for $i \neq j$.

The covariance matrix is $\text{Cov}(\epsilon) = \Sigma = E[\epsilon\epsilon^T] = \sigma^2 I_N$.

3. **Definition of OLS (Ordinary Least Squares) Estimator** For a classical linear regression model with a non-singular matrix $X^T X$, the vector $\hat{\theta} = (X^T X)^{-1} X^T Y$ is called the Ordinary Least Squares estimator.

13.2 Best Linear Unbiased Estimator (BLUE)

1. **Definition and Conditions for Linear Unbiased Estimators** A linear estimator $\tilde{\theta} = AY$ is unbiased if and only if $AX = I$.
2. **Covariance Dominance (Minimum Variance)** For any vector z of appropriate dimension and any unbiased estimator $\tilde{\theta}$, the inequality $D(z^T(\hat{\theta} - \theta)) \leq D(z^T(\tilde{\theta} - \theta))$ holds, where D denotes the covariance matrix, and $\hat{\theta}$ is the BLUE. This implies that the matrix $D_{\tilde{\theta}} - D_{\hat{\theta}}$ is positive semi-definite.
3. **Conditions (a)-(c) for BLUE** An estimator $\hat{\theta}$ is the Best Linear Unbiased Estimator (BLUE) if it satisfies:
 - (a) Estimator Unbiasedness: $E[\hat{\theta}] = \theta$.
 - (b) Minimum Variance: $\hat{\theta}$ minimizes the variance of any linear combination $z^T \hat{\theta}$ among all linear unbiased estimators.
 - (c) Linearity: $\hat{\theta} = SY$, where S is a matrix independent of Y .
4. **Variance Minimization Principle** The BLUE minimizes the variance of every linear combination $z^T \hat{\theta}$ across all possible linear unbiased estimators.

Theorem 13.1 (Gauss-Markov Theorem). Consider the classical linear regression model $(Y, X\theta, \sigma^2 I_N)$, where the error vector ϵ satisfies: (1) $E[\epsilon] = 0$, (2) $\text{Cov}(\epsilon) = \sigma^2 I_N$, and (3) components are uncorrelated. If the matrix $X^T X$ is nonsingular, then the vector

$$\hat{\theta} = (X^T X)^{-1} X^T Y$$

is the Best Linear Unbiased Estimator (BLUE) of θ . Its covariance matrix is given by $D_{\hat{\theta}} = \sigma^2 (X^T X)^{-1}$.

Lemma 13.2 (Properties of the OLS Estimator). The OLS estimator $\hat{\theta} = (X^T X)^{-1} X^T Y$ is a linear and unbiased estimator, i.e., it satisfies conditions (a) and (c) for BLUE.

Lemma 13.3 (Unbiasedness Condition). A linear estimator $\tilde{\theta} = AY$ is unbiased if and only if $AX = I$.

Lemma 13.4 (Covariance Form). Under the assumptions of the Gauss-Markov Theorem, the covariance matrix of any linear unbiased estimator $\tilde{\theta} = AY$ is $D_{\tilde{\theta}} = \sigma^2 AA^T$.

14 OLS Estimation in the Singular Case

14.1 OLS Estimation in the Singular Case

1. **OLS solutions for singular $X^T X$** If $X^T X$ is singular ($\text{rank}(X^T X) < m$), the normal equation $X^T X \theta = X^T Y$ still has solutions, but not a unique one. Any vector $\hat{\theta}$ satisfying this equation minimizes the residual sum of squares $\|Y - X\theta\|^2$ and is called an OLS estimator.
2. **Concept of the Generalized Inverse** A matrix $A^- \in \mathbb{R}^{m \times n}$ is called a generalized inverse of A if, for every vector $y \in \mathbb{R}^n$ such that the system $Ax = y$ is consistent, the vector $x = A^- y$ is a solution.
3. **Moore-Penrose Pseudoinverse** A matrix $A^+ \in \mathbb{R}^{m \times n}$ is called the Moore-Penrose pseudoinverse of A if it satisfies the four Penrose conditions: (1) $AA^+A = A$, (2) $A^+AA^+ = A^+$, (3) $(AA^+)^T = AA^+$, and (4) $(A^+A)^T = A^+A$.
4. **Penrose Conditions** The four defining algebraic conditions for the Moore-Penrose pseudoinverse (listed above).
5. **Representation of all OLS estimators via the Generalized Inverse** The general solution to the consistent system $X^T X \theta = X^T Y$ can be represented using a generalized inverse $A^- = (X^T X)^-$ as $\theta = A^- X^T Y + (H - I)z$, where $H = A^- A$ and z is an arbitrary vector.

Theorem 14.1 (Generalized Inverse Solution). A linear parametric function $\tau = T\theta$ is estimable if and only if $T(X^T X)^- X^T X = T$. If this condition is satisfied, the OLS-estimator $\hat{\tau} = T(X^T X)^- X^T Y$ is uniquely defined and represents the best linear unbiased estimator.

Lemma 14.2 (Condition for Generalized Inverse). For a matrix B to be a generalized inverse of matrix A , it is necessary and sufficient that $ABA = A$.

Lemma 14.3 (Existence of Generalized Inverse). For any matrix A , there exists a generalized inverse A^- .

15 WLS/GLS and Design of Experiments

15.1 Weighted and Generalized Least Squares (WLS / GLS)

1. **Generalized Linear Regression Model** The model is written as $(Y, X\theta, \sigma^2 W)$, where $W \in \mathbb{R}^{N \times N}$ is a known positive definite matrix, and $\sigma^2 > 0$ is an unknown scalar parameter.
2. **Definition of the GLS Estimator** The Generalized Least Squares (GLS) estimator is defined as:

$$\hat{\theta} = (X^T W^{-1} X)^{-1} X^T W^{-1} Y$$

3. **Properties of the GLS Estimator** The covariance matrix of $\hat{\theta}$ is $D_{\hat{\theta}} = \sigma^2 (X^T W^{-1} X)^{-1}$.
4. **Optimality (BLUE in the Generalized Model)** According to the Gauss-Markov theorem, the GLS estimator is the Best Linear Unbiased Estimator (BLUE) under the generalized linear model assumptions.

15.2 Design of Experiments: Basic Concepts

1. **Model Structure** $y_j = \eta(t_j, \theta) + \epsilon_j$ for $j = 1, \dots, N$, where $\eta(t, \theta) = \theta^T f(t)$ (linearity in θ), $t_j \in \chi$ (design points), and ϵ_j are observation errors.
2. **Standard Assumptions (a)-(f)** The model assumes (a) Unbiasedness ($E[\epsilon_j] = 0$), (b) Uncorrelated errors ($E[\epsilon_i \epsilon_j] = 0$ for $i \neq j$), (c) Homoscedasticity ($E[\epsilon_j^2] = \sigma^2 > 0$), (d) Linearity in θ , (e) Basis functions $f_i(t)$ are continuous and linearly independent on χ , and (f) Design space χ is compact.
3. **Definition of Discrete Design (ξ_N)** A design represented by $\xi_N = (\frac{t_1}{1/N} \dots \frac{t_N}{1/N})$, where t_i may repeat, and N is the total number of observations.
4. **Definition of Approximate Design (ξ)** A design represented by $\xi = (\frac{t_1}{\omega_1} \dots \frac{t_n}{\omega_n})$, where t_i are n distinct support points, $\omega_i \geq 0$ are weights (relative frequencies), and $\sum_{i=1}^n \omega_i = 1$.
5. **Full Design Space (Ξ)** The full design space Ξ is the union of all approximate designs with exactly n support points: $\Xi = \bigcup_{n=1}^{\infty} \Xi_n$.

15.3 Information Matrix and Variance of Estimates

1. **Definition of the Information Matrix ($M(\xi)$)** The information matrix of a design ξ is defined as the integral of the outer product of the regressor vector $f(t)$ with respect to the design measure $\xi(dt)$:

$$M(\xi) = \int_{\chi} f(t) f^T(t) \xi(dt) \in \mathbb{R}^{m \times m}$$

2. **Relation to Covariance of Parameter Estimates** For a nonsingular design ξ , the information matrix is proportional to the inverse of the covariance matrix of the OLS estimator: $D_{\hat{\theta}} = \sigma^2 N^{-1} (M(\xi))^{-1}$.
3. **Criteria of Optimality based on $M(\xi)$** Optimality criteria are real-valued functions defined on $M(\xi)$ (or its inverse $D(\xi) = M(\xi)^{-1}$) that quantify the precision of the design.

Theorem 15.1 (Properties of Information Matrices). The following statements hold for the set $\mathcal{M} = \{M : M = M(\xi), \xi \in \Xi\}$:

1. Every information matrix $M(\xi)$ is positive semidefinite.
2. If $n < m$ (fewer support points than parameters), then $\det M(\xi) = 0$.
3. The set \mathcal{M} of all information matrices is convex.
4. Under regularity conditions, \mathcal{M} is compact.
5. For any $\xi \in \Xi$, there exists $\tilde{\xi}$ with $n \leq \frac{m(m+1)}{2} + 1$ support points such that $M(\tilde{\xi}) = M(\xi)$.

16 Optimality Criteria and Equivalence Theorem

16.1 Optimality Criteria in Design

1. **D-optimality Criterion** Minimizes the volume of the confidence ellipsoid by maximizing the determinant of the information matrix: $\log \det M(\xi) \rightarrow \sup_{\xi} \text{ or } \log \det D(\xi) \rightarrow \inf_{\xi \in \Xi}$.
2. **L-optimality Criterion** Minimizes generalized quadratic loss by minimizing the trace of L times the inverse information matrix: $\text{tr}[LD(\xi)] \rightarrow \inf_{\xi \in \Xi_{NS}}$, where L is a fixed nonnegative definite matrix.
3. **E-optimality Criterion** Maximizes the smallest eigenvalue of the information matrix, which minimizes the longest axis of the confidence ellipsoid: $\lambda_{\min}(M(\xi)) \rightarrow \sup_{\xi \in \Xi_{NS}} \text{ or } \lambda_{\max}(D(\xi)) \rightarrow \inf_{\xi \in \Xi_{NS}}$.
4. **e_k -optimality Criterion** Minimizes the variance of the estimator for a single parameter coordinate θ_k : $e_k^T M^{-1}(\xi) e_k \rightarrow \inf_{\xi \in \Xi_{e_k}}$.
5. **G-optimality Criterion** Minimizes the maximum prediction variance over the design space: $\max_{t \in \chi} d(t, \xi) \rightarrow \inf_{\xi \in \Xi_{NS}}$, where $d(t, \xi) = f^T(t) M^{-1}(\xi) f(t)$.

Theorem 16.1 (Kiefer-Wolfowitz Equivalence Theorem). *Under standard regularity assumptions (including compactness of the set of information matrices), the following conditions are equivalent for a design ξ^* :*

1. ξ^* is D-optimal.
2. ξ^* is G-optimal.
3. $\max_{x \in \chi} d(x, \xi^*) = m$, where m is the number of parameters.

Theorem 16.2 (D-Optimal Designs for the Polynomial Model). *For polynomial regression of degree $m - 1$ on $\chi = [-1, 1]$, a unique approximate D-optimal design exists. It is supported with equal weights ($1/m$) on m points, which are the roots of the polynomial $(x^2 - 1)P'_{m-1}(x)$, where $P_{m-1}(x)$ is the Legendre polynomial of degree $m - 1$.*

Theorem 16.3 (D-Optimal Designs for the Trigonometric Model). *An approximate D-optimal design for the trigonometric regression model of order m on $\chi = [-\pi, \pi]$ is any design supported with equal weights ($1/N$) on $N \geq 2m + 1$ equally spaced points $t_i^* = \frac{i-1}{N} 2\pi - \pi$ for $i = 1, \dots, N$. The continuous uniform design $\xi^* = \frac{1}{2\pi} dx$ is also D-optimal.*