

Fourier series

TASKS

1. Decompose the function into a Fourier series $f(x) = \text{sign } x, -\pi < x < \pi$, and using the resulting decomposition, find the sum of the Leibniz series

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1}$$

Answer: $\text{sign } x = \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{\sin(2n-1)x}{2n-1}$, Leibniz series: $\sum_{n=1}^{\infty} \frac{(-1)^n}{2n+1} = \frac{\pi}{4}$

Decompose the Fourier series of the function $f(x)$ on the specified interval, the length of the interval is the period (2-11).

$$2. f(x) = \begin{cases} A, & 0 < x < l, \\ A/2, & x = l, \\ 0, & l < x < 2l, \end{cases} \quad \text{on the interval } (0, 2l).$$

Answer: $\frac{A}{2} + \frac{2A}{\pi} \sum_{n=1}^{\infty} \frac{1}{2n-1} \sin \frac{2n-1}{l} \pi x$

$$3. f(x) = |x| \text{ on the segment } [-1; 1].$$

Answer: $\frac{1}{2} - \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{\cos \pi(2n-1)x}{(2n-1)^2}$

$$4. f(x) = \begin{cases} ax, & -\pi < x < 0, \\ bx, & 0 \leq x < \pi, \end{cases} \quad \text{in the interval } (-\pi, \pi).$$

Answer: $\frac{(b-a)\pi}{4} - \sum_{n=1}^{\infty} ((b-a)(1 - (-1)^n) \frac{\cos nx}{\pi n^2} + (-1)^n(a+b) \frac{\sin nx}{n})$

$$5. f(x) = \begin{cases} a, & -\pi/2 < x < \pi/2, \\ b, & \pi/2 \leq x < 3\pi/2, \end{cases} \quad \text{in the range } (-\pi/2, 3\pi/2).$$

Answer: $\frac{a+b}{2} + \frac{2(a-b)}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1} \cos(2n-1)x$

$$6. f(x) = x + \text{sign } x \text{ on the interval } (-\pi; \pi).$$

Answer: $\frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1+(-1)^{n+1}(1+\pi)}{n} \sin nx$

$$7. f(x) = \pi^2 - x^2 \text{ on the interval } (-\pi; \pi).$$

Answer: $\frac{2}{3}\pi^2 + 4 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} \cos nx$

$$8. f(x) = x^3 \text{ on the interval } (-\pi; \pi).$$

Answer: $\sum_{n=1}^{\infty} (-1)^n (\frac{12}{n^3} - \frac{2\pi^2}{n}) \sin nx$

$$9. f(x) = e^{ax}, a \neq 0, \text{ in the interval } (-\pi; \pi).$$

Answer: $\frac{2}{\pi} \text{sh } a\pi (\frac{1}{2a} + \sum_{n=1}^{\infty} \frac{(-1)^n}{a^2+n^2} (a \cos(nx) - n \sin nx))$

$$10. f(x) = e^{2|x|} \text{ in the interval } (-\pi; \pi).$$

Answer: $\frac{e^{2\pi}-1}{2\pi} + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n e^{2\pi}-1}{n^2+4} \cos nx$

11. $f(x) = \sin ax, a \in Z$ in the interval $(-\pi; \pi)$.
 Answer: $\frac{2\sin\pi a}{\pi} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{n\sin nx}{n^2 - a^2}, \quad -\pi < x < \pi$