

# HW1. Version 5.

$$1. \int \left( \sin \frac{x}{2} - \cos \frac{x}{2} \right)^2 dx$$

$$= \int (1 - 2 \sin \frac{x}{2} \cos \frac{x}{2}) dx = \int (1 - \sin x) dx$$

$$= x + \cos x + C$$

$$2. \int \frac{x^4}{x^5+7} dx$$

$$= \frac{1}{5} \int \frac{d(x^5+7)}{x^5+7} = \frac{1}{5} \ln |x^5+7| + C.$$

$$3. \int e^x \sin x dx$$

$$\int e^x \sin x dx = -e^x \cos x + \int e^x \cos x dx$$

$$\int e^x \cos x dx = e^x \sin x - \int e^x \sin x dx$$

$$\Rightarrow \int e^x \sin x dx = \frac{e^x \sin x - e^x \cos x}{2} + C$$

$$4. \int \frac{5x+2}{x^2+2x+10} dx$$

$$= \int \frac{\frac{5}{2} d(x^2+2x+10)}{x^2+2x+10} - \frac{1}{9} \int \frac{d(\frac{x+1}{3})}{(\frac{x+1}{3})^2+1}$$

$$= \frac{5}{2} \ln(x^2+2x+10) - \frac{1}{9} \arctan \frac{x+1}{3} + C$$

$$5. \int \frac{dx}{3 \sin x + 4 \cos x}$$

$$= \int \frac{dx}{5 \sin(x+\varphi)} \quad \varphi = \arctan \frac{4}{3}$$

$$= \frac{1}{5} \ln \left| \tan \frac{x+\varphi}{2} \right| + C \quad \varphi = \arctan \frac{4}{3}.$$

## HW2 Version 5.

1.  $\int_0^{\pi/3} \cos^3 x \sin 2x \, dx$
2.  $\int_{-3}^3 \frac{x^3 \sin 2x}{x^2+1} \, dx$ .
3. Find the area of a flat figure bounded by a single cycloid arch and the Ox-axis:  
 $x=2(t-\sin t)$ ,  $y=2(1-\cos t)$
4. Find the volumes of bodies formed by rotation around the Ox-axes of a figure bounded by lines:  
 $y=\sqrt{x}e^x$ ,  $x=1$ ,  $y=0$ .
5. Calculate improper integrals (or establish their divergence):  
 $\int_1^{+\infty} \frac{e^{-x^2}}{x^2} \, dx$ .

$$1. \int_0^{\frac{\pi}{3}} \cos^3 x \sin 2x \, dx = -2 \int_0^{\frac{\pi}{3}} \cos^4 x \, d(\cos x) = -2 \left. \frac{\cos^5 x}{5} \right|_0^{\frac{\pi}{3}} = \frac{31}{80}$$


$$2. \int_{-3}^3 \frac{x^3 \sin 2x}{x^2+1} \, dx = 0 \quad \text{odd function with symmetric domain.}$$

$$3. \text{ single cycloid : } 0 \leq t \leq 2\pi$$

$$S = \int y(x) \, dx = \int_0^{2\pi} y(t) \cdot x'(t) \, dt = 4 \int_0^{2\pi} (1-\cos t)^2 \, dt = 4 \int_0^{2\pi} (1-2\cos t + \cos^2 t) \, dt$$

$$= 4 \left( 2\pi + 2\pi \cdot \frac{1}{2} \right) = 12\pi.$$

4.



$$V = \pi \int_a^b y^2(x) \, dx = \pi \int_0^1 (\sqrt{x} e^x)^2 \, dx = \pi \int_0^1 x e^{2x} \, dx$$

$$= \pi \left[ \frac{1}{2} x \cdot e^{2x} \Big|_0^1 - \frac{1}{2} \int_0^1 e^{2x} \, dx \right] = \pi \left[ \frac{e^2}{2} - \frac{e^2-1}{4} \right] = \pi \frac{e^2+1}{4}.$$

$$5. \int_1^{+\infty} \frac{e^{-x^2}}{x^2} \, dx$$

$$e^{-x^2} < 1 \quad \text{in } [1, +\infty)$$

$$\int_1^{+\infty} \frac{1}{x^2} \, dx = -\frac{1}{x} \Big|_1^{+\infty} = 1 \quad \text{conv.} \quad \text{thus the integral } \int_1^{+\infty} \frac{e^{-x^2}}{x^2} \, dx \quad \text{conv.}$$