

COMENIUS UNIVERSITY IN BRATISLAVA
FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS

TRIANGULATION OF IMPLICIT SURFACE WITH
SINGULARITIES
MASTER'S THESIS

2021

BC. KRISTÍNA KORECOVÁ

COMENIUS UNIVERSITY IN BRATISLAVA
FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS

TRIANGULATION OF IMPLICIT SURFACE WITH SINGULARITIES

MASTER'S THESIS

Study Programme: Computer Graphics and Geometry
Field of Study: Mathematics
Department: Department of Algebra and Geometry
Supervisor: doc. RNDr. Pavel Chalmovianský, PhD.

Bratislava, 2021

Bc. Kristína Korecová



Univerzita Komenského v Bratislave
Fakulta matematiky, fyziky a informatiky

ZADANIE ZÁVEREČNEJ PRÁCE

Meno a priezvisko študenta: Bc. Kristína Korecová
Študijný program: počítačová grafika a geometria (Jednoodborové štúdium, magisterský II. st., denná forma)
Študijný odbor: matematika
Typ záverečnej práce: diplomová
Jazyk záverečnej práce: anglický
Sekundárny jazyk: slovenský

Názov: Triangulation of implicitly defined surfaces
Triangulácia implicitne definovaných plôch

Anotácia:

Cieľ:

Literatúra:

**Kľúčové
slová:**

Vedúci: doc. RNDr. Pavel Chalmovianský, PhD.
Katedra: FMFI.KAG - Katedra algebry a geometrie
Vedúci katedry: doc. RNDr. Pavel Chalmovianský, PhD.

Spôsob sprístupnenia elektronickej verzie práce:
prípustná pre vlastnú VŠ

Dátum zadania: 19.10.2021

Dátum schválenia: 20.10.2021

prof. RNDr. Július Korbaš, CSc.
garant študijného programu

.....
študent

.....
vedúci práce

Acknowledgements: TODO Acknowledgements

Abstrakt

TODO Abstrakt po Slovensky

Klíčové slová: TODO Klíčové slová

Abstract

TODO Abstract in English

Keywords: TODO Keywords

Contents

Introduction	1
1 Surface triangulation and its application	3
2 Theoretical background	5
2.1 Implicit surfaces	5
2.2 ADE singularities	9
2.3 Non-isolated translation surface singularities	11
2.4 Tringulation of regular implicit srufaces	11
2.5 Data structures for triangulation algorithm	11
2.6 CSG modelling for implcit surfaces	11
3 Our contributions	15
3.1 Triangulation adaptive to the local curvature	15
3.2 Triangulation of ADE singularities	16
3.3 Triangulation of non-isolated singularities of translation surfaces	23
4 Results	25
4.1 Quality criteria	25
4.2 Comparison with TODO	25
5 Future work	27
Conclusion	29
Appendix A	33
Appendix B	35

List of Figures

2.1	Implicit surfaces with corresponding equations	5
2.2	Isolated and non-isolated singularity	6
2.3	Normal cut	7
2.4	Visualisation of the curvature of the double-torus	8
2.5	Example of CSG tree	12
2.6	Boolean operations on implicit curves	13
2.7	Intersection of a sphere and a plane	14
3.1	Adaptive height of the new triangle	15
3.2	A_{n--} singularities	17
3.3	A_{n+-} singularities	17
3.4	D_{n+-} singularities	18
3.5	D_{n--} singularities	19
3.6	Intersection of D_{n--} singularities with plane $z = 0$	19
3.7	Triangulation vectors for two branches of D_{n--} singularities.	19
3.8	E_n singularities.	20
3.9	Intersection of E_{7++} and E_{8++} singularities with plane $z = 0$	20
3.10	Triangulation of A_{n--} singularity.	21
3.11	Equidistant points on ellipse.	22
3.12	TODO.	22

List of Tables

Introduction

TODO Introduction

Chapter 1

Surface triangulation and its application

Chapter 2

Theoretical background

2.1 Implicit surfaces

Implicit functions are a tool for surface representation and manipulation. In computer graphics, they can be used for modelling of complex surfaces using boolean operations, realistic animations, rendering and other.

Implicit functions do not define the boundary explicitly, instead the surface is defined as a zero set of a function.


Definition 1 *Given a function $F : \mathbb{R}^3 \rightarrow \mathbb{R}$, one can define an implicit surface as a set of points that fulfil $F(x, y, z) = 0$.*

Some examples of implicit surfaces and their equations can be seen on image 2.1.

Unit normal vector of the implicit surface in point (x_0, y_0, z_0) is normalized gradient of the implicit function in that point.


Definition 2 *Gradient vector of an implicit function $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ is defined as*

$$\nabla F(x, y, z) = \left(\frac{\partial F(x, y, z)}{\partial x}, \frac{\partial F(x, y, z)}{\partial y}, \frac{\partial F(x, y, z)}{\partial z} \right).$$



$$x^2 + y^2 + z^2 - 1 = 0$$

a) sphere



$$x^2 + y^2 - z^2 = 0$$

b) cone



$$z = 0$$

c) plane

Figure 2.1: Implicit surfaces with corresponding equations.




Figure 2.2: Isolated and non-isolated singularity.

If $\nabla F(x, y, z) \neq 0$, we can define the unit normal vector of F as a normalized gradient vector

$$N(F(x, y, z)) = \frac{\nabla F(x, y, z)}{\|\nabla F(x, y, z)\|} \quad \text{for } \nabla F(x, y, z) \neq 0.$$

Points lying on the implicit surface can be classified as regular or singular based on the value of the gradient vector in that point.

Definition 3 Point $P = (x, y, z)$ lying on the implicit surface is said to be regular, if $\nabla F(x, y, z) \neq 0$. On the contrary, point P is said to be singular, if $\nabla F(x, y, z) = 0$.

Singular points can be further classified as isolated or non-isolated based on their surroundings.


Definition 4 Singular point P is said to be isolated, if there exists an open ball $B_\epsilon(P)$, which does not contain any other singular point. Singular point P is said to be non-isolated if it is not isolated.

On the image 2.2 we can see example of isolated and non-isolated singularities.

Curvature of a surface

Curvature is a fundamental concept in differential geometry of curves and surfaces. In case of curves, curvature is a measure of how much does the curve differ from a straight line. It is defined as the inverse of the radius of the osculating circle, which is the second order approximation of the curve.

For surfaces, curvature is a measure of how much does the surface differ from a plane. The definition of the curvature of a surface is not as straightforward as in the case of curves, as the curvature depends on the choice of the direction in which we measure the curvature.

Figure 2.3: Normal cut of the parametric surface $S(u, v)$.

The idea of measuring the curvature of a surface has a long history in mathematics. One of the first contributors was a mathematician Carl Friedrich Gauss, who developed the idea of the Gaussian curvature of surfaces. In this subsection, we are drawing from the summary presented by Tiago Novello et al.[7].

Normal curvature of the surface

Let S be a parametric surface

$$S(u, v) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

$$S(u, v) = (S_x(u, v), S_y(u, v), S_z(u, v)).$$

Let us denote the unit normal vector of the surface S in the point $S(u, v)$ as $\overrightarrow{n(u, v)}$.

We define the normal curvature of the surface as a function of the location of the point on the surface given by parameters u and v and the unit tangent vector in that point \overrightarrow{u} .

Definition 5 *Normal cut of a surface S in the regular point P in the direction of the unit tangent vector \overrightarrow{u} is defined as an intersection of the surface S and a plane given by the vectors \overrightarrow{u} and $\overrightarrow{n(u, v)}$.*

The visualisation of the normal cut is shown on the image 2.3. It is clear, that the normal cut is a plane curve lying on the surface, we denote this normal cut as $\nu_S(u, v, \overrightarrow{u})$.

Definition 6 *Oriented normal curvature of the surface in the regular point P in the direction of the unit tangent vector \overrightarrow{u} is defined as the curvature of the normal cut $\nu_S(u, v, \overrightarrow{u})$. Non-oriented normal curvature is defined as an absolute value of the oriented normal curvature.*

Definition 7 *Minimal and maximal curvature in the point $P = S(u, v)$ are defined as*

$$\kappa_{min}(u, v) = \min_{\overrightarrow{u} \in T_P(u, v)} \nu_S(u, v, \overrightarrow{u}),$$




Figure 2.4: Visualisation of the curvature of the double-torus.

$$\kappa_{max}(u, v) = \max_{\vec{u} \in T_P(u, v)} \nu_S(u, v, \vec{u}),$$

where $T_S(u, v)$ is a tangent plane of the surface S in the point P .

Minimal and maximal curvature are called *principal curvatures*.

Definition 8 *Gaussian curvature is defined as a product of principal curvatures:*

$$\kappa_G(u, v) = \kappa_{min}(u, v)\kappa_{max}(u, v).$$

Gaussian curvature describes the shape of the surface in the local neighborhood of the point. The points where Gaussian curvature is positive are called elliptic points. The points where Gaussian curvature is negative are called hyperbolic points. The points where only one of κ_{min} , κ_{max} is zero are called parabolic and the points where both κ_{min} and κ_{max} are zero are called planar. The shape of the surface in the local neighborhoods of the points is as follows:

- elliptic points \longrightarrow surface is curved like a sphere,
- hyperbolic points \longrightarrow surface is curved like a saddle,
- parabolic points \longrightarrow surface is curved like a parabolic cylinder,
- planar points \longrightarrow surface is flat.

Gaussian curvature is an intrinsic property, which means that it is also independent of the placement of the surface in the space.

Definition 9 *Mean curvature is defined as an arithmetic mean of principal curvatures:*

$$\kappa_M(u, v) = \frac{\kappa_{min}(u, v) + \kappa_{max}(u, v)}{2}.$$

Minimal, maximal, Gaussian and mean curvature are visualized on the image 2.4.

Curvature formulas for implicit surface

A version of curvature formulas for implicit surfaces appeared in [10] and were reformulated, summarized and proved by Ron Goldman [4]. In this subsection we point out these formulas.

Let $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ be an implicit function which defines surface by the equation $F(x, y, z) = 0$. Let us denote $F_t = \frac{\partial F}{\partial t}$ and $F_{ts} = \frac{\partial^2 F}{\partial t \partial s}$. Hessian matrix - the matrix of second derivatives is defined as

$$H(F) = \begin{pmatrix} F_{xx} & F_{xy} & F_{xz} \\ F_{yx} & F_{yy} & F_{yz} \\ F_{zx} & F_{zy} & F_{zz} \end{pmatrix},$$

and the adjoint of the Hessian is defined as

$$H^*(F) = \begin{pmatrix} F_{yy}F_{zz} - F_{yz}F_{zy} & F_{yz}F_{zx} - F_{yx}F_{zz} & F_{yx}F_{zy} - F_{yy}F_{zx} \\ F_{xz}F_{zy} - F_{xy}F_{zz} & F_{xx}F_{zz} - F_{xz}F_{zx} & F_{xy}F_{zx} - F_{xx}F_{zy} \\ F_{xy}F_{yz} - F_{yx}F_{zy} & F_{yx}F_{xz} - F_{xx}F_{yz} & F_{xx}F_{yy} - F_{xy}F_{yx} \end{pmatrix}.$$

We can now formulate the formulas of Gaussian, mean, minimal and maximal curvature.

Gaussian curvature of the implicit surface defined by function F is given by

$$\kappa_G = \frac{\nabla F * H^*(F) * \nabla F^T}{|\nabla F|^4}.$$

Mean curvature of the implicit surface defined by function F is given by

$$\kappa_M = \frac{\nabla F * H^*(F) * \nabla F^T - |\nabla F|^2 \text{Trace}(H)}{2|\nabla F|^3}.$$

The principal curvatures κ_{min} and κ_{max} can be calculated from Gaussian curvature and mean curvature as

$$\kappa_{min}, \kappa_{max} = \kappa_M \pm \sqrt{\kappa_M^2 - \kappa_G}.$$

2.2 ADE singularities

ADE singularities, also referred to as du Val singularities are a specific class of simple, isolated surface singularities. They were classified by Arnold's [1] according to ADE classification [5] based on correspondence of these singularities to simply laced Dynkin diagrams [2]. We know infinitely many A singularities – A_1, A_2, \dots , infinitely many D singularities – D_4, D_5, \dots and three E singularities – E_6, E_7 and E_8 . ADE singularities are specified by their normal forms. When working in complex space, each singularity has a single normal form:

- A_n $F(x, y, z) = x^{n+1} + y^2 + z^2,$
- D_n $F(x, y, z) = yx^2 + y^{n-1} + z^2,$
- E_6 $F(x, y, z) = x^3 + y^4 + z^2,$
- E_7 $F(x, y, z) = x^3 + xy^3 + z^2,$
- E_8 $F(x, y, z) = x^3 + y^5 + z^2.$

Each ADE singularity on a surface can be locally expressed by their normal form.

In the real case, changing the signs in these equations produces different surfaces and therefore, ADE singularities can be further classified by their signature.

Definition 10 *Let's mark real surface singularities based on their signature as follows:*

- $A_{n\pm\pm}$ $F(x, y, z) = x^{n+1} \pm y^2 \pm z^2,$
- $D_{n\pm\pm}$ $F(x, y, z) = yx^2 \pm y^{n-1} \pm z^2,$
- $E_{6\pm\pm}$ $F(x, y, z) = x^3 \pm y^4 \pm z^2,$
- $E_{7\pm\pm}$ $F(x, y, z) = x^3 \pm xy^3 \pm z^2,$
- $E_{8\pm\pm}$ $F(x, y, z) = x^3 \pm y^5 \pm z^2.$

The most common example of a surface with ADE singularity is an ordinary cone. Given as the zero set of the function $F(x, y, z) = x^2 - y^2 - z^2$, cone has a singular point $P = (0, 0, 0)$. This singular point is an example of A_{1--} singularity.

Correspondence between $SO(3, \mathbb{R})$ group and ADE singularities

$SO(3, \mathbb{R})$ is special orthogonal group over the field of real numbers in three dimensions. It is also called 3D rotation group, as it is a group of all rotations about the origin in \mathbb{R}^3 .

Definition 11 *$SO(3, \mathbb{R})$ is a group of 3×3 orthogonal matrices of real numbers with determinant 1.*

$$SO(3, \mathbb{R}) = \left\{ A \in \mathbb{R}^{3 \times 3} \mid AA^T = I, \det(A) = 1 \right\}.$$

Simply laced Dynkin diagrams correspond to all finite subgroups of $SO(3, \mathbb{R})$. Finite subgroups of $SO(3, \mathbb{R})$ are the rotational symmetry groups of

- pyramid with n vertices (cyclic subgroup \overline{C}_n),
- double pyramid with n vertices (dihedral subgroup \overline{D}_n),

- platonic solids
 - tetrahedron (tetrahedral subgroup \overline{T})
 - octahedron (octahedral subgroup \overline{O})
 - icosahedron (icosahedral subgroup \overline{I})

The correspondence is as follows:

- $A_n \iff \overline{C}_{n+1}$,
- $D_n \iff \overline{D}_{n+2}$,
- $E_6 \iff \overline{T}$,
- $E_7 \iff \overline{O}$,
- $E_8 \iff \overline{I}$.

The conclusion is that ADE singularities correspond to finite subgroups of $SO(3, \mathbb{R})$, which represent certain types of symmetries in \mathbb{R}^3 .

2.3 Non-isolated translation surface singularities

2.4 Tringulation of regular implicit srufaces

2.5 Data structures for triangulation algorithm

2.6 CSG modelling for implcit surfaces

In this section we will discuss how constructive solid geometry can be used for modelling complex implicit surfaces. First major publication regarding constructive solid geometry was published in 1977 by Requicha and Voelcker [9]. More detailed mathematical foundations were published a year later, in 1978 by Requicha and Tilove [8].

Constructive solid geometry (CSG)

Constructive solid geometry is a technique used for modelling complex geometric objects using boolean operations on sets of points – union, intersection and difference. It gained populatity in 1980s as a powerful tool to create complex shapes from sets of geometrical primitives, such as cylinders, spheres and cones. The resulting model can be represented as a CSG tree, which contains geometrical primitives in its leaves and boolean operations in its internal nodes [3]. An example of such CSG tree can be seen on figure 2.5.




Figure 2.5: Example of CSG tree [3].

CSG for implicit surfaces

As the implicit surface divides space into inside ($F(x, y, z) < 0$) and outside ($F(x, y, z) > 0$), one can easily combine the idea of implicit surfaces and CSG modelling. Given two implicit surfaces represented by implicit functions F and G , the surface of the intersection of the interiors can be defined by implicit function $H = \min(F, G)$. Similarly, the surface of the union of the interiors can be defined by implicit function $H = \max(F, G)$ and lastly, the surface of the difference of the interiors can be defined by implicit function $H = \max(F, -G)$. Two dimensional example of this idea can be seen on figure 2.6.

Theorem 1 *The minimum function $\min(F, G)$ of two functions $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ is defined as*

$$\min(F, G) = F + G - \sqrt{F^2 + G^2}$$

. *The maximum function $\max(F, G)$ of two functions $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ is defined as*

$$\max(F, G) = F + G + \sqrt{F^2 + G^2}$$

.

These formulas allow us to model implicit surfaces which are the result of performing finite number of operations union, intersection and difference on arbitrary implicit functions.




Figure 2.6: Boolean operations on implicit curves.

An easy example is creating an implicit equation representing a half of a sphere. Let us have an implicit function $F(x, y, z) = x^2 + y^2 + z^2 - 1$, which represents a unit sphere and another implicit function $G(x, y, z) = z$ which represents the xy plane. The minimum function of these two functions

$$\min(F, G) = x^2 + y^2 + y^2 - 1 + z - \sqrt{(x^2 + y^2 + y^2 - 1)^2 + z^2}$$

represents the surface of a half sphere. The visualisation of these surfaces can be seen on figure 2.7.




Figure 2.7: Intersection of a sphere and a plane.

Chapter 3

Our contributions

3.1 Triangulation adaptive to the local curvature

As we explained in the beginning of chapter 2.1, curvature of the surface is a measure of how much the surface bends.

The triangulation of the surface should be accurate enough, but also memory efficient. This can be achieved by creating a triangulation which is locally adaptive to the curvature of the surface. Therefore having smaller triangles in the places where the surface is curved and having bigger triangles where surface is flatter.

In this section we present our implementation of the triangulation adaptive to the local curvature.

In the original algorithm, the height of the triangle which is projected to the surface is set to the constant value $\frac{\sqrt{3}}{2}e$, where e is the required length of the side of the triangle. To achieve the adaptivity of the triangles size, we set the height of the triangle to depend on the curvature in the given point, as shown in the image 3.1.

To identify the curved areas, we decided to use the maximal curvature. As the minimal and maximal curvature are both signed, we want to identify the areas depending on the absolute value of these curvatures.

We do not allow arbitrary height of the triangle to avoid edge-cases. We decided to restrict the allowed height to $\frac{1}{4}\frac{\sqrt{3}}{2}e > h > 4\frac{\sqrt{3}}{2}e$. We create a variable m -multiplier, depending on κ_T and set the height of the new triangle to $h = m\frac{\sqrt{3}}{2}e$.

As κ_T has values in range $\langle 0, \infty \rangle$.




Figure 3.1: Adaptive height of the new triangle.

Definition 12 *Let us define triangulation curvature of the surface S in the point $P = S(u, v)$ as $\kappa_T(u, v) = \max(|\kappa_{\min}|, |\kappa_{\max}|)$.*

NOT FINISHED YET

3.2 Triangulation of ADE singularities

Analysis of the geometry of ADE singularities

ADE singularities are simple, isolated surface singularities, which can be expressed by corresponding implicit equations.

We already know, that A_{1--} singularity is locally represented as a cone. In this section we discuss geometric structure of other ADE surface singularities.

Definition 13 *(TODO rewrite) Let us define branch of ADE singularity as the part of the surface, which is connected to the rest only by the singular point.*



For our needs, we pick one triangulation vector for each branch of each ADE singularity. This triangulation vector is normalized vector either in the direction each ADE singularity. This triangulation vector is normalized vector either in the direction of rotation symmetry axis or an intersection of reflection symmetry planes of the corresponding branch. If the branch has only one reflection symmetry plane, the triangulation vector is picked to lie in the reflection symmetry plane. of the corresponding branch. If the branch has only one reflection symmetry plane, the triangulation vector is picked to lie in the reflection symmetry plane.

In the general case, triangulation vectors serve us as a partial information about the orientation of a singularity with respect to its normal form. In the general case, triangulation vectors serve us as a partial information about the orientation of a singularity with respect to its normal form.

A_n singularities

As we can see from the equations $F(x, y, z) = x^{n+1} \pm y^2 \pm z^2$, A_{n-+} singularities are just rotated A_{n+-} singularities and A_{n++} singularities are a single point if n is odd and reflected A_{n--} singularities if n is even. are a single point if n is odd and reflected A_{n--} singularities if n is even. We therefore only discuss geometry of A_{n--} and A_{n+-} singularities.

A_{n--} singularities are topologically equivalent to a cone if n is odd, therefore they have two branches. If n is even, they are topologically equivalent to a half cone or a plane, therefore they have a single branch. As n gets bigger, the tip of the cone gets sharper. As A_{n--} singularities are rotationally symmetrical, we pick the direction of

Figure 3.2: A_{n--} singularities. [6]Figure 3.3: A_{n+-} singularities. [6]

axis of symmetry as triangulation vector. For a normal form, the triangulation vectors are $(1, 0, 0)$ (and $(-1, 0, 0)$ if n is odd). First four A_{n--} singularities can be seen on image 3.2.

A_{n+-} singularities are topologically equivalent to a cone if n is odd, therefore they have two branches. In the contrary with the previous singularities, as n gets bigger, the tip of the cone gets less sharp and flatter. Branches of these singularities have reflection symmetry planes $x = 0$ and $y = 0$, therefore we pick the vectors $(0, 0, 1)$ and $(0, 0, -1)$ as the triangulation vectors.


If n is even, A_{n+-} singularities are topologically equivalent to a plane with shape similar to hyperbolic paraboloid, therefore they have a single branch. First four A_{n+-} singularities can be seen on image 3.3. For this case, we pick the vector $(1, 0, 0)$ as a triangulation vector as these singularities have reflection symmetry planes $y = 0$ and $z = 0$.

D_n singularities

Given by equations $F(x, y, z) = yx^2 \pm y^{n-1} \pm z^2$, we consider 8 categories. For given sign combination and parity of n , the singularities are topologically equivalent, with sharper(or flatter) features around the singularities for increasing value of n similar to A_n singularities.

We can therefore say that D_n singularities can be classified into 8 categories locally represented by the following equations:

- $D_{4++} \quad yx^2 + y^3 + z^2$

Figure 3.4: D_{n+-} singularities. [6]

- $D_{5++} \quad yx^2 + y^4 + z^2$
- $D_{4+-} \quad yx^2 + y^3 - z^2$
- $D_{5+-} \quad yx^2 + y^4 - z^2$
- $D_{4-+} \quad yx^2 - y^3 + z^2$
- $D_{5-+} \quad yx^2 - y^4 + z^2$
- $D_{4--} \quad yx^2 - y^3 - z^2$
- $D_{5--} \quad yx^2 - y^4 - z^2$.

Now we look at some equivalences between these 8 categories. D_{4++} singularity is reflected D_{4+-} singularity. D_{5++} singularity is reflected D_{5--} singularity. D_{5-+} singularity is reflected D_{5+-} singularity. D_{4-+} singularity is reflected D_{4--} singularity.




We therefore only analyze geometry of D_{n+-} singularities and D_{n--} singularities.



D_{n+-} singularities are topologically equivalent to a plane when n is even and to a cone when n is odd. Again, as n gets bigger, the features around singularities get sharper. Symmetry planes of these singularities are $x = 0$ and $z = 0$, therefore we pick $(0, 1, 0)$ (and $(0, -1, 0)$ when n is odd) as triangulation vectors. First four D_{n+-} singularities can be seen on image 3.4.

D_{n--} singularities are topologically equivalent to a cone when n is odd and to a 3 halfcones connected in the singular point when n is even. First four D_{n--} singularities can be seen on image 3.5.

Symmetry plane for all branches of these singularities is $z = 0$. the intersection of the surface and plane $z = 0$ is displayed on image 3.6.

For D_{n--} singularity, the intersections of the two branches where $y \geq 0$ are bounded by curves $y = 0$ and $x^2 = y^{n-2}$. For given r , we pick the triangulation vectors as $(r, \frac{1}{2}r^{\frac{2}{n-2}}, 0)$ and $(-r, \frac{1}{2}r^{\frac{2}{n-2}}, 0)$. The resulting vectors are displayed on image 3.7 by blue arrow. Parameter r is changed based on the length of the edge of triangulation triangle. displayed on image 3.7 by blue arrow. Parameter r is changed based on the length of the edge of triangulation triangle.


 Figure 3.5: D_{n--} singularities. [6]

 Figure 3.6: Intersection of D_{n--} singularities with plane $z = 0$.

 Figure 3.7: Triangulation vectors for two branches of D_{n--} singularities.

Figure 3.8: E_n singularities. [6]Figure 3.9: Intersection of E_{7++} and E_{8++} singularities with plane $z = 0$.

The third branch where $y \leq 0$ has another plane of symmetry $x = 0$, therefore triangulation vector for this branch is chosen as $(0, -1, 0)$.

E_6, E_7 and E_8 singularities

Given by equations $F(x, y, z) = x^3 \pm y^4 \pm z^2$, $F(x, y, z) = x^3 \pm xy^3 \pm z^2$ and $F(x, y, z) = x^3 \pm y^5 \pm z^2$, we can see the following equivalences: E_{6++} singularity is reflected E_{6--} singularity. E_{6+-} singularity is reflected E_{6-+} singularity. E_{7+-}, E_{7-+} and E_{7--} are all reflected E_{7++} singularity. E_{8+-}, E_{8-+} and E_{8--} are all reflected E_{8++} singularity.

We only analyze geometry of E_{6++} , E_{6+-} , E_{7++} and E_{8++} singularities. These singularities are displayed on the image 3.8.


Both E_{6++} and E_{6+-} are topologically equivalent to a plane, thus they each have only one branch. The planes of symmetry of both of these branches are $y = 0$ and $z = 0$, therefore we pick $(-1, 0, 0)$ as the triangulation vector.

E_{7++} singularity is topologically equivalent to a cone, therefore it has E_{7++} singularity is topologically equivalent to a cone, therefore it has two branches. The plane of symmetry of this singularity is $z = 0$.

E_{8++} singularity is also topologically equivalent to a plane, therefore it has only one branch. This branch has only one plane of symmetry $z = 0$.

We again look at the intersection of the surfaces with the plane of symmetry, this is displayed on image 3.9.

For E_{7++} singularity, we pick $(-1, 0, 0)$ and $(\frac{1}{2}r^{\frac{3}{2}}, -r, 0)$ as triangulation vectors.

Figure 3.10: Triangulation of A_{n--} singularity.

For E_{8++} singularity, we pick $(-1, -1, 0)$ as a triangulation vector. These vectors are displayed on the image 3.9 as blue arrows.

Analytical calculation of local triangulation of some ADE singularities

For given edge size e , we want to calculate the local triangulation of ADE singularities, such that edges on the border of the local triangulation have length e .

A_{n--} singularities

For A_{n--} singularities, we create a disc of 6 isosceles triangles with vertex in the singular point. The bases of these triangles create regular hexagon in the plane P parallel to the plane $x = 0$, as showed on the image 3.10. 3.10. Given by equation $x^{n+1} - y^2 - z^2 = 0$, we find the distance of the plane P from the plane $x = 0$ for the given length e of the sides of the hexagon.

Let e be the length of the side of the hexagon, then the circumscribed circle has radius e . This circle is identical with the intersection of the surface and the plane $x = h$. The equation of the intersecting circle is $y^2 + z^2 = h^{n+1}$ therefore, the radius can be also expressed as $r = h^{\frac{n+1}{2}}$, which emerges $h = e^{\frac{2}{n+1}}$. Knowing the distance of the plane, one can easily calculate the length of the arms of the triangles using Pythagorean theorem:

$$a^2 = h^2 + e^2 \implies a = \sqrt{e^{\frac{4}{n+1}} + e^2}$$

D_n singularities

Some D_n singularities have branches with elliptical intersection with a plane parallel to the plane $y = 0$. As ellipses have 2 axes of symmetry, we create 8 triangles for these branches.

Let us have an ellipse E with semi-major axis a and semi-minor axis b . We are going to create eight triangles with apex in the singular point. The other points of the triangles lie on the ellipse and they have the same length of the base.




Figure 3.11: Equidistant points on ellipse.




Figure 3.12: TODO.

As displayed on image 3.11, we pick the leftmost, the rightmost, the top and the bottom points. Then we can calculate the point P on ellipse equidistant from points P_1 and P_2 .

$$\begin{aligned} \frac{1}{2}(a, b) + \frac{t}{2}(b, a) \in E &\implies \frac{(a + tb)^2}{4a^2} + \frac{(b + ta)^2}{4b^2} = 1 \\ 4b^2(a^2 + 2atb + t^2b^2) + 4a^2(b^2 + 2atb + t^2a^2) - a^2b^2 &= 0 \\ 4(b^4 + a^4)t^2 + 8ab(b^2 + a^2)t + 7a^2b^2 &= 0 \\ t = \frac{ab(\sqrt{3a^4 + 2a^2b^2 + 3b^4} - a^2 - b^2)}{a^4 + b^4} \\ P = \frac{1}{2}(a, b) + \frac{ab(\sqrt{3a^4 + 2a^2b^2 + 3b^4} - a^2 - b^2)}{2(a^4 + b^4)}(b, a) \end{aligned}$$

TODO binary search for height

Given edge length e , we are not able to calculate the height in which the distance between points P_1 and P is e . The visualization showing this is on the image 3.12.

We use binary search to find such height. Given the height and the singularity class, we can calculate the semi-major axis and semi-minor axis as

$$D_{n+-} : -hx^2 + h^{n-1} - z^2 = 0 \quad h > 0$$

$$x^2 + \frac{z^2}{h} = h^{n-2}$$

$$\frac{x^2}{h^{n-2}} + \frac{z^2}{h^{n-1}} = 1 \implies a_h = \max(h^{\frac{n-2}{2}}, h^{\frac{n-1}{2}}) \wedge b_h = \min(h^{\frac{n-2}{2}}, h^{\frac{n-1}{2}}).$$

As we can see, we get the same ellipse for D_{n--} singularities:

$$D_{n--} : -hx^2 - h^{n-1} - z^2 = 0 \quad h > 0$$

$$2|n \wedge x^2 + \frac{z^2}{h} = -h^{n-2} \implies x^2 + \frac{z^2}{h} = h^{n-2}.$$

Then

$$P_h = \frac{1}{2}(h^{\frac{n-2}{2}}, h^{\frac{n-1}{2}}) + \frac{h^{\frac{2n-3}{2}}(\sqrt{3h^{2n-4} + 2h^{2n-3} + 3h^{2n-2}} - h^{n-2} - h^{n-1})}{2(h^{2n-4} + h^{2n-2})}(h^{\frac{n-1}{2}}, h^{\frac{n-2}{2}})$$

$$P_h = \frac{1}{2}(h^{\frac{n-2}{2}}, h^{\frac{n-1}{2}}) + \frac{h^{\frac{1}{2}}(\sqrt{3 + 2h + 3h^2} - 1 - h)}{2(1 + h^2)}(h^{\frac{n-1}{2}}, h^{\frac{n-2}{2}})$$

and we can calculate $e_h = ||P_h - P_1||$.

As $e \leq a_h$, we can start the binary search on the interval $\langle 0, a^{\frac{2}{n-2}} \rangle$ or $\langle 0, a^{\frac{2}{n-1}} \rangle$ and finish, when required precision is reached.

TODO want to try to prove that $||P_h - P_1||$ is monotone in h .

E_6, E_7 and E_8 singularities

Numerical calculation of local triangulation of ADE singularities

3.3 Triangulation of non-isolated singularities of translation surfaces

Chapter 4

Results

4.1 Quality criteria

4.2 Comparison with TODO

Chapter 5

Future work

Conclusion

TODO Conclusion

Bibliography

- [1] Normal forms for functions near degenerate critical points, the weyl groups of A_n , D_n , E_n and lagrangian singularities. *Functional Analysis and its applications*, 6:254–272, 1972.
- [2] Evgenii Borisovich Dynkin. The structure of semi-simple algebras. *Uspekhi Matematicheskikh Nauk*, 2(4):59–127, 1947.
- [3] James D Foley, Foley Dan Van, Andries Van Dam, Steven K Feiner, and John F Hughes. *Computer graphics: principles and practice*, volume 12110. Addison-Wesley Professional, 1996.
- [4] Ron Goldman. Curvature formulas for implicit curves and surfaces. *Computer Aided Geometric Design*, 22(7):632–658, 2005.
- [5] M Hazewinkel, W Hesselink, D Siersma, and FD Veldkamp. The ubiquity of coxeter-dynkin diagrams. *Nieuw Arch. Wisk*, 25:257–307, 1977.
- [6] Dr. Richard Morris. 3d viewer-generator.
- [7] TIAGO NOVELLO, VINÍCIUS DA SILVA, GUILHERME SCHARDONG, LUIZ SCHIRMER, HÉLIO LOPES, and LUIZ VELHO. Differential geometry of implicit surfaces. 2021.
- [8] Aristides Requicha and Robert Tilove. Mathematical foundations of constructive solid geometry: General topology of closed regular sets. 1978.
- [9] Aristides AG Requicha and Herbert B Voelcker. Constructive solid geometry. 1977.
- [10] Michael Spivak. *A comprehensive introduction to differential geometry*, volume 3. Publish or Perish, Incorporated, 1975.

Appendix A

Appendix B