#### 1

# Solution of question 9.3.8

## Gagan Singla - EE22BTECH11021

Question: Five cards are drawn successively with replacement from a well-shuffled deck of 52 cards. What is the probability that

- (a) all the five cards are spades?
- (b) only 3 cards are spades?
- (c) none is a spade?

**Solution:** Let us define:

Parameter	Value	Description
n	5	number of cards drawn
p	$\frac{1}{4}$	drawing a spade card
q	$\frac{3}{4}$	drawing any other card
$\mu = np$	<u>5</u>	mean of the distribution
$\sigma^2 = npq$	15 16	variance of the distribution
Y	{0,1,2,3,4,5}	Random Variable

### (i) Gaussian Distribution

The gaussian distribution function is defined as:

$$p_Y(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \qquad (x \in Y) \quad (1)$$

(a) If we consider all cards to be spades,

$$Y = 5 \tag{2}$$

Substituting values in (1),

$$p_Y(5) = \frac{1}{\sqrt{2\pi \left(\frac{15}{16}\right)}} e^{-\frac{\left(5 - \frac{5}{4}\right)^2}{2\left(\frac{15}{16}\right)}}$$
(3)

$$=\frac{1}{\sqrt{2\pi}}e^{-\frac{15}{2}}\tag{4}$$

= 0.0001245 (5)

(b) If we consider 3 cards to be spades,

$$Y = 3 \tag{6}$$

Substituting values in (1),

$$p_Y(3) = \frac{1}{\sqrt{2\pi \left(\frac{15}{16}\right)}} e^{-\frac{\left(3-\frac{5}{4}\right)^2}{2\left(\frac{15}{16}\right)}} \tag{7}$$

$$=\frac{1}{\sqrt{2\pi}}e^{-\frac{49}{30}}\tag{8}$$

$$= 0.044$$
 (9)

(c) If we consider 0 cards to be spades,

$$Y = 0 \tag{10}$$

Substituting values in (1),

$$p_Y(0) = \frac{1}{\sqrt{2\pi\left(\frac{15}{16}\right)}} e^{-\frac{\left(0-\frac{5}{4}\right)^2}{2\left(\frac{15}{16}\right)}}$$
(11)

$$= \frac{1}{\sqrt{2\pi}} e^{-\frac{5}{6}} \tag{12}$$

$$= 0.0978$$
 (13)

## (ii) Gaussian vs Binomial Comparison

Y	Gaussian	Binomial
0	0.0978	0.2373
3	0.044	0.08789
5	0.0001245	0.00098

## (iii) Binomial vs Gaussian Graph

