

SOCCOM IN2015_V01 SOTS HPLC

Avuntaura Gullledge¹, Nils Haëntjens¹, Emmanuel Boss¹, and Lynne Talley²

¹University of Maine, ²Scripps Institution of Oceanography

March 1, 2018

Sample collection

Near-surface samples from SOCCOM CTD stations were taken for HPLC analysis. Volume filtered varied from 1 to 2 L to achieve maximum loading without blockage. Filters used were GF/F 25 mm diameter.

More information on the cruise are available at:

<https://socom.princeton.edu/content/shipboard-data-reports>

Analysis method

Analysis was performed by Lesley Clementson (lesley.clementson@csiro.au) at CSIRO following the protocol of Hooker et al. 2012.

References

Hooker, Stanford B., Lesley Clementson, Crystal S. Thomas, Louise Schlüter, Merete Allerup, Josephine Ras, Herve Claustre, Claire Normandeau, John Cullen, Markus Kienast, Wendy Kozlowski, Maria Vernet, Sumit Chakraborty, Steven Lohrenz, Merritt Tuel, Donald Redalje, Paulo Cartaxana, Carlos R. Mendes, Vanda Brotas, S.G. Prabhu Matondkar, Sushma G. Parab, Aimee Neeley, and Einar Skarstad Egeland. The Fifth SeaWiFS HPLC Analysis Round-Robin Experiment (SeaHARRE-5), 2012, NASA/TM-2012-217503.