
SAMPLE_R1_KM original R1 subtype composition
0
1
2
3
4
5
0.75
0.50
0.25
Arterial EC
C1QC TAM
CD4 Helper/Conventional
CD4 Naive/Memory-like
CD8 Exhausted
CD8 GZMK/Effector-memory-like
Capillary EC
Classical Mono
Epithelial
GC B
LAMP3 DC
Lymphatic EC
Mast
Matrix_CAF
NK
Naive/Memory B
Neutrophil
Pericyte
Plasma
SMC
SPP1 TAM
Tfh/CXCL13-like
Treg
Venous EC
cDC1
cDC2
iCAF
pDC
rCAF
vCAF
SAMPLE_R1_KM original R2 subtype composition
0
1
2
3
4
5
0.75
0.50
0.25
Arterial EC
C1QC TAM
CD4 Helper/Conventional
CD4 Naive/Memory-like
CD8 Exhausted
CD8 GZMK/Effector-memory-like
Capillary EC
Classical Mono
Epithelial
GC B
LAMP3 DC
Lymphatic EC
Mast
Matrix_CAF
NK
Naive/Memory B
Neutrophil
Pericyte
Plasma
SMC
SPP1 TAM
Tfh/CXCL13-like
Treg
Venous EC
cDC1
cDC2
iCAF
pDC
rCAF
vCAF