




GSE171351_A1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12















Tissue 768 | Epi 488 | internal 229 | P95 12
Exact front in Epi 177 | R12 labelled 409


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE171351_A1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE171351_A1 | exploratory scatter display selected by coverage and |delta r|, not P












Treg / Budden S1P RESPONSE 18: delta r=0.254 [nan, nan]; common 300–721 µm; spatial P=nan
Matrix_CAF / Budden TNF EMT RESPONSE 15: delta r=-0.172 [nan, nan]; common 0–500 µm; spatial P=nan
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.


P95 nonP95 distance 0


GSE171351_B1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12















Tissue 1017 | Epi 959 | internal 651 | P95 33
Exact front in Epi 429 | R12 labelled 853


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE171351_B1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE171351_B1 | exploratory scatter display selected by coverage and |delta r|, not P










Neutrophil / Flores pro-invasion 10: delta r=-0.571 [-0.730, -0.348]; common 100–557 µm; spatial P=1.1e-05
Naive/Memory B / Flores pro-invasion 10: delta r=0.514 [-0.017, 0.853]; common 0–1229 µm; spatial P=0.021
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.


P95 nonP95 distance 0

GSE171351_C1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 1469 | Epi 1465 | internal 493 | P95 25
Exact front in Epi 894 | R12 labelled 1290


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE171351_C1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE171351_C1 | exploratory scatter display selected by coverage and |delta r|, not P










Treg / Budden S1P RESPONSE 18: delta r=-0.094 [-0.593, 0.433]; common 100–557 µm; spatial P=0.71
Matrix_CAF / Budden TNF EMT RESPONSE 15: delta r=0.029 [-0.345, 0.553]; common 100–1308 µm; spatial P=0.88
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.


P95 nonP95 distance 0

GSE171351_D1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12

MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled


UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 832 | Epi 3 | internal 1 | P95 1
Exact front in Epi 0 | R12 labelled 0


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE171351_D1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE171351_D1 | exploratory scatter display selected by coverage and |delta r|, not P













Treg / Budden S1P RESPONSE 18: delta r=nan [nan, nan]; common nan–-inf μm; spatial P=nan
Matrix_CAF / Budden TNF EMT RESPONSE 15: delta r=nan [nan, nan]; common nan–-inf μm; spatial P=nan
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 μm blocks.

P95 nonP95 distance 0


GSE171351 | Neutrophil / Flores pro-invasion 10
Section heterogeneity; missing patient IDs remain sample units
P95 | section
nonP95 | n shown on left
delta r | common support ∩ [0,1000] µm
GSE171351_A1 (n=12)
GSE171351_B1 (n=33)
GSE171351_C1 (n=25)
GSE171351_D1 (n=0)
207
527
464
0
−1.0
−0.5
0.0
0.5
1.0
−2
−1
0
1
2
Pearson r / joint spatial 95% CI
Pearson r / joint spatial 95% CI
P95 r minus nonP95 r


GSE171351 / Neutrophil / Flores pro-invasion 10
Gouin2021::1204-1A
Gouin2021::72-1A
Gouin2021::8-1A
−1.00
−0.75
−0.50
−0.25
0.00
0.25
0.50
0.75
1.00
Equal-section Fisher-z aggregate / conditional spatial 95% CI
P95
nonP95


GSE171351 | iCAF / Budden CHOLESTEROL SYNTHESIS 27
Section heterogeneity; missing patient IDs remain sample units
P95 | section
GSE171351_A1 (n=11)
GSE171351_B1 (n=29)
GSE171351_C1 (n=24)
GSE171351_D1 (n=0)
−1.0
−0.5
0.0
0.5
1.0
Pearson r / joint spatial 95% CI
nonP95 | n shown on left
124
472
468
0
−1.0
−0.5
0.0
0.5
1.0
Pearson r / joint spatial 95% CI
delta r | common support ∩ [0,1000] µm
−2
−1
0
1
2
P95 r minus nonP95 r


GSE171351 / iCAF / Budden CHOLESTEROL SYNTHESIS 27
Gouin2021::1204-1A
Gouin2021::72-1A
Gouin2021::8-1A
−1.00
−0.75
−0.50
−0.25
0.00
0.25
0.50
0.75
1.00
Equal-section Fisher-z aggregate / conditional spatial 95% CI
P95
nonP95