

GSE285715_N1 | adjacent_normal_source
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 1417 | Epi 213 | internal 0 | P95 0
Exact front in Epi 0 | R12 labelled 38


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_N1 | all 29 candidates; fixed core programs


normal_Epi raw Pearson r | full observed range




P95 minus nonP95 r | common support | grey = not estimable



GSE285715_N1 | exploratory scatter display selected by coverage and |delta r|, not P



Adjacent normal reference
Tumor MP10-P95 is not defined.





Adjacent normal reference
Tumor MP10-P95 is not defined.


No tumor common-support contrast; no range is fabricated.
No tumor common-support contrast; no range is fabricated.
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 μm blocks.


P95 nonP95 distance 0


GSE285715_N2 | adjacent_normal_source
Frozen mixed-spot MP10 / front / R12
















Tissue 1722 | Epi 72 | internal 0 | P95 0
Exact front in Epi 0 | R12 labelled 29

GSE285715_N2 | all 29 candidates; fixed core programs


normal_Epi raw Pearson r | full observed range




P95 minus nonP95 r | common support | grey = not estimable



GSE285715_N2 | exploratory scatter display selected by coverage and |delta r|, not P



Adjacent normal reference
Tumor MP10-P95 is not defined.




Adjacent normal reference
Tumor MP10-P95 is not defined.

No tumor common-support contrast; no range is fabricated.
No tumor common-support contrast; no range is fabricated.
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 μm blocks.


P95 nonP95 distance 0

GSE285715_N3 | adjacent_normal_source
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 2489 | Epi 2024 | internal 0 | P95 0
Exact front in Epi 0 | R12 labelled 1490


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_N3 | all 29 candidates; fixed core programs


normal_Epi raw Pearson r | full observed range




P95 minus nonP95 r | common support | grey = not estimable



GSE285715_N3 | exploratory scatter display selected by coverage and |delta r|, not P



Adjacent normal reference
Tumor MP10-P95 is not defined.




Adjacent normal reference
Tumor MP10-P95 is not defined.

No tumor common-support contrast; no range is fabricated.
No tumor common-support contrast; no range is fabricated.
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 μm blocks.


P95 nonP95 distance 0

GSE285715_T1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 4285 | Epi 4167 | internal 3504 | P95 177
Exact front in Epi 1869 | R12 labelled 3885


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_T1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE285715_T1 | exploratory scatter display selected by coverage and |delta r|, not P










Capillary EC / GLYCOLYSIS: delta r=-0.487 [-0.624, -0.070]; common 0–1136 μm; spatial P=0.00032
Plasma / OXIDATIVE PHOSPHORYLATION: delta r=0.360 [0.143, 0.557]; common 0–1400 μm; spatial P=0.00041
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 μm blocks.


P95 nonP95 distance 0

GSE285715_T2 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 3987 | Epi 3821 | internal 2110 | P95 106
Exact front in Epi 2277 | R12 labelled 3443


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_T2 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE285715_T2 | exploratory scatter display selected by coverage and |delta r|, not P










SMC / Budden CHOLESTEROL SYNTHESIS noMP10 18: delta r=-0.474 [-0.655, -0.274]; common 200–1500 µm; spatial P=8.4e-06
GC B / Puram p-EMT Top30: delta r=0.337 [0.150, 0.507]; common 0–2066 µm; spatial P=0.00053
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.


P95 nonP95 distance 0

GSE285715_T3 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 3341 | Epi 3317 | internal 1881 | P95 97
Exact front in Epi 875 | R12 labelled 3000


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_T3 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE285715_T3 | exploratory scatter display selected by coverage and |delta r|, not P













NK / Budden CHOLESTEROL SYNTHESIS 27: delta r=-0.608 [-0.755, -0.256]; common 0–854 µm; spatial P=1.5e-05
GC B / G2M CHECKPOINT: delta r=0.470 [0.122, 0.714]; common 0–889 µm; spatial P=0.0012
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0


GSE285715 | SMC / Budden CHOLESTEROL SYNTHESIS 27
Section heterogeneity; missing patient IDs remain sample units
P95 | section
nonP95 | n shown on left
delta r | common support ∩ [0,1000] µm
GSE285715_N1 (NA)
GSE285715_N2 (NA)
GSE285715_N3 (NA)
GSE285715_T1 (n=177)
GSE285715_T2 (n=77)
GSE285715_T3 (n=96)
NA
NA
NA
3287
1453
1741
−1.0
−0.5
0.0
0.5
1.0
−2
−1
0
1
2
Pearson r / joint spatial 95% CI
Pearson r / joint spatial 95% CI
P95 r minus nonP95 r


GSE285715 / SMC / Budden CHOLESTEROL SYNTHESIS 27
−1.00
−0.75
−0.50
−0.25
0.00
0.25
0.50
0.75
1.00
Equal-section Fisher-z aggregate / conditional spatial 95% CI
P95
nonP95


GSE285715 | Pericyte / MTORC1 SIGNALING (no MP10)
Section heterogeneity; missing patient IDs remain sample units
P95 | section
GSE285715_N1 (NA)
GSE285715_N2 (NA)
GSE285715_N3 (NA)
GSE285715_T1 (n=177)
GSE285715_T2 (n=103)
GSE285715_T3 (n=97)
−1.0
−0.5
0.0
0.5
1.0
Pearson r / joint spatial 95% CI
nonP95 | n shown on left
NA
NA
NA
3321
1910
1765
−1.0
−0.5
0.0
0.5
1.0
Pearson r / joint spatial 95% CI
delta r | common support ∩ [0,1000] µm
−2
−1
0
1
2
P95 r minus nonP95 r


GSE285715 / Pericyte / MTORC1 SIGNALING (no MP10)
−1.00
−0.75
−0.50
−0.25
0.00
0.25
0.50
0.75
1.00
Equal-section Fisher-z aggregate / conditional spatial 95% CI
P95
nonP95