

CanDis_2026_Sample2 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 10062 | Epi 4928 | internal 4363 | P95 219
Exact front in Epi 2460 | R12 labelled 4580




CanDis_2026_Sample2 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




CanDis_2026_Sample2 | exploratory scatter display selected by coverage and |delta r|, not P









Pericyte / Budden S1P RESPONSE 18: delta r=0.497 [0.353, 0.624]; common 100–2960 µm; spatial P=6.7e-09
Pericyte / Budden CHOLESTEROL SYNTHESIS 27: delta r=-0.431 [-0.573, -0.280]; common 100–2960 µm; spatial P=1.4e-07
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0