

CanDis_2026_Sample7 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 2295 | Epi 1858 | internal 1366 | P95 69
Exact front in Epi 1033 | R12 labelled 1576


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

CanDis_2026_Sample7 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




CanDis_2026_Sample7 | exploratory scatter display selected by coverage and |delta r|, not P









Classical Mono / GLYCOLYSIS: delta r=0.371 [0.182, 0.547]; common 0–1905 µm; spatial P=0.00016
Pericyte / Budden S1P RESPONSE 18: delta r=-0.329 [-0.574, -0.057]; common 0–2052 µm; spatial P=0.02
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0