

GSE285715_T2 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 3987 | Epi 3821 | internal 2110 | P95 106
Exact front in Epi 2277 | R12 labelled 3443


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_T2 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE285715_T2 | exploratory scatter display selected by coverage and |delta r|, not P









SMC / Budden CHOLESTEROL SYNTHESIS noMP10 18: delta r=-0.474 [-0.655, -0.274]; common 200–1500 µm; spatial P=8.4e-06
GC B / Puram p-EMT Top30: delta r=0.337 [0.150, 0.507]; common 0–2066 µm; spatial P=0.00053
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0