

GSE285715_T3 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 3341 | Epi 3317 | internal 1881 | P95 97
Exact front in Epi 875 | R12 labelled 3000


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE285715_T3 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE285715_T3 | exploratory scatter display selected by coverage and |delta r|, not P









NK / Budden CHOLESTEROL SYNTHESIS 27: delta r=-0.608 [-0.755, -0.256]; common 0–854 µm; spatial P=1.5e-05
GC B / G2M CHECKPOINT: delta r=0.470 [0.122, 0.714]; common 0–889 µm; spatial P=0.0012
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0