

GSE319536_BC08 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 3444 | Epi 1297 | internal 1121 | P95 57
Exact front in Epi 482 | R12 labelled 1088


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE319536_BC08 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE319536_BC08 | exploratory scatter display selected by coverage and |delta r|, not P









cDC1 / Puram p-EMT Top30: delta r=-0.614 [-0.943, -0.386]; common 100–1513 µm; spatial P=1.4e-05
Mast / Budden CHOLESTEROL SYNTHESIS noMP10 18: delta r=0.334 [0.055, 0.599]; common 436–2234 µm; spatial P=0.013
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0