

GSE319536_BC12 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 1654 | Epi 884 | internal 764 | P95 39
Exact front in Epi 142 | R12 labelled 698


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE319536_BC12 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE319536_BC12 | exploratory scatter display selected by coverage and |delta r|, not P









Mast / E2F TARGETS: delta r=-0.600 [-0.840, -0.318]; common 173–1752 µm; spatial P=1.2e-05
vCAF / Budden CHOLESTEROL SYNTHESIS 27: delta r=0.532 [0.232, 0.761]; common 0–917 µm; spatial P=0.0013
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0