

GSE319536_BC18 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 2847 | Epi 2656 | internal 2204 | P95 111
Exact front in Epi 649 | R12 labelled 2297


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

GSE319536_BC18 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




GSE319536_BC18 | exploratory scatter display selected by coverage and |delta r|, not P









Mast / Budden CHOLESTEROL SYNTHESIS 27: delta r=-0.367 [-0.557, -0.099]; common 0–1127 µm; spatial P=0.0018
Mast / GLYCOLYSIS: delta r=0.268 [0.078, 0.439]; common 0–1127 µm; spatial P=0.0054
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0