

NPJ_sample1 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95




Strict v3.4 signed distance (µm)




Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient



Observed coverage only; no extrapolation



Frozen P95 vs nonP95 within [-200, 0] µm




Tissue 2076 | Epi 996 | internal 589 | P95 30
Exact front in Epi 508 | R12 labelled 766


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

NPJ_sample1 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range





P95 minus nonP95 r | common support | grey = not estimable




NPJ_sample1 | exploratory scatter display selected by coverage and |delta r|, not P









Treg / MTORC1 SIGNALING (no MP10): delta r=-0.903 [-1.232, -0.066]; common 100–1044 µm; spatial P=0.001
cDC1 / Budden S1P RESPONSE 18: delta r=0.637 [0.213, 1.066]; common 100–889 µm; spatial P=0.0023
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0