

NPJ_sample2 | tumor_study_specimen_not_region_diagnosis
Frozen mixed-spot MP10 / front / R12


MP10 AUCell and frozen P95





Strict v3.4 signed distance (µm)





Formal R12 0–8; grey = unlabelled




UCell neighborhood: adjusted MP10 coefficient




Observed coverage only; no extrapolation




Frozen P95 vs nonP95 within [-200, 0] µm





Tissue 4506 | Epi 3217 | internal 3006 | P95 151
Exact front in Epi 1026 | R12 labelled 2832


0 1 2 3 4 5 6 7 8 P95 nonP95 / all Epi

NPJ_sample2 | all 29 candidates; fixed core programs


P95 raw Pearson r | full observed range






P95 minus nonP95 r | common support | grey = not estimable




NPJ_sample2 | exploratory scatter display selected by coverage and |delta r|, not P




rCAF / Budden TNF EMT RESPONSE 15: delta r=0.638 [0.411, 0.822]; common 100–3372 µm; spatial P=4.7e-07
NK / Budden CHOLESTEROL SYNTHESIS noMP10 18: delta r=-0.526 [-0.679, -0.347]; common 0–1868 µm; spatial P=5.5e-09
Orange points: distance = 0. Pearson P assumes independent spots; spatial intervals use 500 µm blocks.

P95 nonP95 distance 0