

The Patient-Specific Functional Scale

This useful questionnaire can be used to quantify activity limitation and measure functional outcome for patients with any orthopaedic condition.

Clinician to read and fill in below: Complete at the end of the history and prior to physical examination.

Initial Assessment: 2/06/2026 - pre trial

I am going to ask you to identify up to three important activities that you are unable to do or are having difficulty with as a result of your Multiple Sclerosis problem. Today, are there any activities that you are unable to do or having difficulty with because of your _____ problem? (Clinician: show scale to patient and have the patient rate each activity).

Follow-up Assessments:

When I assessed you on (state previous assessment date), you told me that you had difficulty with (read all activities from list at a time). Today, do you still have difficulty with: (read and have patient score each item in the list)?

Patient-specific activity scoring scheme (Point to one number):

0	1	2	3	4	5	6	7	8	9	10
Unable to perform activity									Able to perform activity at the same level as before injury or problem	

(Date and Score)

Activity	Initial	Mid Trial (1/5/26)	4 weeks post end of trial		
1. Walk to the corner shops from my house	0	6	1		
2. Walk on uneven surfaces	2	7	2		
3. To build confidence walking to and navigating medical appointments	1	8	1		
4. To build confidence to be able to walk at my local Westfield shopping centre	0	3	0		
To be able to do stairs	2	5	2		
Additional					
Additional					

Total score = sum of the activity scores/number of activities

Minimum detectable change (90%CI) for average score = 2 points

Minimum detectable change (90%CI) for single activity score = 3 points

PSFS developed by: Stratford, P., Gill, C., Westaway, M., & Binkley, J. (1995). Assessing disability and change on individual patients: a report of a patient specific measure. Physiotherapy Canada, 47, 258-263.

Reproduced with the permission of the authors.