

List of Figures

2.1	An example potential for a driven particle with periodic boundary conditions.	17
2.2	Two slowest timescales for the 1D driven system depending on lagtime.	21
2.3	Chapman-Kolmogorov test for the metastable states for the 1D driven system.	21
2.4	(a) Probability distribution for the equilibrium system with metastable states identified by PCCA+. (b) First and second eigenvector Ψ_i representing the probability flow during relaxation process.	23
2.5	Mean first-passage time depending on the lagtime τ for all six observed processes for the 1D system.	25
2.6	First-passage time distribution of process from metastable state B to A at different lagtimes.	26
3.1	Equilibrium-reweighted energy distribution of the interacting self-avoiding walk.	34
3.2	Illustration of symmetric and asymmetric observables under time reversal and space inversion.	39
3.3	Stationary distribution and first-passage time distribution for the driven 1D system reweighted with respect to global entropy production.	42
3.4	Stationary distribution and first-passage time distribution for the driven 1D system reweighted with respect to local entropy production.	44
3.5	Transition probability matrix of the 1D driven system and the violation of detailed balance.	46
3.6	Stationary distribution and first-passage time distribution for the driven 1D system reweighted with respect to local entropy production and global balance.	47
3.7	Deviation of constraints when attempting to find a full solution to the Caliber maximisation by numerical minimisation.	51
4.1	Potential surface, stationary distribution and first-passage time distribution of a chosen process for the 1D equilibrium system	54

4.2	The first three moments and the population of metastable states for the equilibrium system under varying tilt of the potential.	56
4.3	Potential surface, stationary distribution and first-passage time distribution of a chosen process for the 1D driven system	57
4.4	First-passage time distribution of transition $C \rightarrow A$ simulated at 3 different driving forces for the 1D driven system.	58
4.5	The first three moments and the population of metastable states for the 1D driven system for varying external force.	60
4.6	Sketch of a 4-state laser model with potential surface and pumping force.	61
4.7	Stationary distribution and first-passage time distribution of the excitation process for the laser model.	62
4.8	The first three moments and the population of metastable states for the laser system for varying external force.	65
4.9	Potential surface, stationary distribution and first-passage time distribution of a chosen process for the 2D driven system	67
4.10	The first three moments and the population of metastable states for the 2D system for varying external force.	68
4.11	First-passage time distribution of transition $A \rightarrow B$ simulated at 3 different driving forces for the 2 system.	69
5.1	Free energy surface, stationary distribution and first-passage time distribution of a chosen process for the 1D reduced system	77
5.2	The first three moments and the population of metastable states for the reduced 1D system for varying driving forces.	79
5.3	First-passage time distribution of transition $A \rightarrow B$ simulated at three different driving forces for the 2D and the reduced model. . .	80
5.4	Atomistic and coarse-grained representation of the tetraalanine peptide.	82
5.5	Free energy surface and lagtime analysis for the two slowest processes of the reference tetraalanine peptide.	82
5.6	Detailed relaxation process analysis of the two slowest processes of the tetraalanine peptide reference model and modified by an end-to-end potential.	84
5.7	The first three moments, the population of metastable states and the first two timescales for the tetraalanine peptide for varying driving along the end-to-end distance.	86
5.8	Stationary distribution and first-passage time distribution of a chosen process for tetraalanine peptide driven along the dihedral angle. . .	88
5.9	The first three moments and the population of metastable states for the tetraalanine peptide for varying driving along the dihedral angle. . .	89

5.10	Transition matrices of chosen initial point for the tetraalanine peptide reference model and a model driven along the dihedral angle.	90
6.1	The dynamical invariant of the 1D driven system.	95
6.2	Two example trajectories from simulation of the 2D system between two microstates.	97
6.3	Distribution of entropy productions for the single particle in 1D potential and single particle in 2D potential for chosen transitions.	98
6.4	Deviation of local entropy production between analytic solution and averaging over simulated trajectories for (a) a single particle on a 1D potential surface and (b) a single particle on a 2D potential surface for a chosen starting point.	99

List of Technical Points

2.1	Stratonovich vs Itô Integration	10
3.1	Maxwell's Demon	28
3.2	Interacting Self-Avoiding Walk	33
4.1	The Kramers Model	72

