

The effects of prolonged term blood transfusion in Postpartum Hemorrhage

*Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²

¹Department of Medical Laboratory Science, Kampala International University, Uganda.

²School of Nursing Science, Kampala International University, Uganda.

*Corresponding authour: Emmanuel Ifeanyi Obeagu, Department of Medical Laboratory Science, Kampala International University, Uganda.

E-mail: emmanuelobeagu@yahoo.com, obeagu.emmanuel@kiu.ac.ug, 0000-0002-4538-0161

Abstract

Postpartum hemorrhage (PPH) remains a significant cause of maternal morbidity and mortality globally. In severe cases of PPH, prolonged or repeated blood transfusions are often necessary to restore hemodynamic stability and prevent adverse outcomes. This review aims to comprehensively examine the effects and implications of prolonged-term blood transfusion in postpartum hemorrhage scenarios. The paper begins by elucidating the physiological consequences of extended blood transfusion in the postpartum period, encompassing alterations in hemoglobin levels, changes in coagulation parameters, immune modulation, and the impact on erythropoiesis and iron metabolism. It also delves into the clinical outcomes associated with prolonged transfusion, encompassing maternal morbidity, transfusion-related complications, and long-term health implications. Additionally, this paper discusses critical considerations for healthcare providers managing prolonged blood transfusion in PPH cases. It explores strategies for optimizing transfusion practices, minimizing risks of complications, and deliberates on the ethical and counseling aspects of prolonged-term transfusion. Through this comprehensive analysis, the review aims to provide insights for healthcare professionals to make informed decisions, optimize management strategies, and improve maternal health outcomes in severe postpartum hemorrhage scenarios. Further research and concerted efforts are essential to advance transfusion practices and enhance maternal care in these critical situations.

Keywords: *postpartum hemorrhage; blood transfusion; maternal outcomes; prolonged transfusion effects; alloimmunization; maternal health; immunomodulation*

Introduction

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

Postpartum hemorrhage (PPH) remains a substantial contributor to maternal mortality and morbidity worldwide, posing significant challenges to obstetric care.¹ Defined as excessive bleeding following childbirth, severe cases of PPH often necessitate urgent interventions to stabilize the mother's condition. Among these interventions, blood transfusion stands as a vital therapeutic modality, particularly in cases of persistent or profound hemorrhage. While blood transfusions are life-saving, the implications of prolonged or repeated transfusions in the postpartum period warrant close examination.² The increasing frequency of severe obstetric hemorrhages and subsequent transfusions has sparked interest in understanding the multifaceted impact of prolonged-term blood transfusion in this specific clinical context.

Postpartum hemorrhage is considered a significant obstetric challenge due to its potential to rapidly escalate into a life-threatening condition for the mother.³ PPH involves excessive bleeding following childbirth, typically occurring within the first 24 hours after delivery.⁴ It poses a substantial risk to maternal health and is a leading cause of maternal mortality and morbidity globally. The challenges associated with PPH arise from its unpredictability, as it can occur suddenly and progress rapidly, leading to severe blood loss and hemodynamic instability.⁵ Common causes include uterine atony (failure of the uterus to contract), retained placental tissue, genital tract trauma, coagulation disorders, or complications arising from cesarean sections or instrumental deliveries. Addressing PPH effectively requires immediate and coordinated interventions by healthcare providers. Failure to manage PPH promptly can result in severe consequences such as hypovolemic shock, organ dysfunction, disseminated intravascular coagulation, and ultimately maternal death. Given its critical nature and potential for adverse outcomes, managing postpartum hemorrhage, especially in severe cases, requires a multidisciplinary approach, rapid assessment, and swift implementation of evidence-based interventions. PPH represents a significant obstetric challenge that necessitates vigilance, expertise, and timely intervention to prevent severe morbidity or mortality associated with excessive postpartum bleeding.⁶ Postpartum hemorrhage, characterized by excessive bleeding following childbirth, represents a scenario where there is an urgent need for intervention due to the potential rapid progression to a life-threatening situation for the mother.⁷ When PPH occurs, immediate and decisive actions are vital to control bleeding, stabilize the mother's condition, and prevent severe consequences associated with substantial blood loss. Given the unpredictable nature of PPH and its capacity to escalate quickly, healthcare providers must swiftly identify and manage the underlying causes to avoid complications such as hypovolemic shock or organ failure.

The physiological consequences of prolonged blood transfusion in the context of postpartum hemorrhage encompass a range of effects on the recipient's body due to extended exposure to transfused blood components.⁸ Prolonged blood transfusion can lead to changes in hematological parameters. This may include alterations in hemoglobin levels, hematocrit, and red blood cell indices, potentially impacting the body's natural erythropoiesis. Extended-term transfusion may affect the coagulation system. It can lead to changes in clotting factors, platelet function, and other components of the coagulation cascade, potentially predisposing individuals to bleeding or thrombotic complications. The immune system may be influenced by transfused blood components, potentially resulting in immune modulation. Prolonged exposure to foreign antigens

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

might lead to immune sensitization or the development of alloantibodies, affecting future transfusion compatibility and increasing the risk of transfusion reactions. Repeated transfusions can result in iron overload due to the accumulation of excess iron from transfused blood.⁹ This excess iron may deposit in organs such as the liver, heart, and pancreas, leading to complications like organ dysfunction, endocrine disturbances, and increased oxidative stress. Continuous exposure to transfused blood may suppress endogenous erythropoietin production and bone marrow function, potentially leading to decreased production of new blood cells and prolonged dependency on transfusions.¹⁰ Prolonged-term transfusion increases the risk of various transfusion-related complications, including allergic reactions, febrile non-hemolytic reactions, transfusion-associated circulatory overload (TACO), and transfusion-transmitted infections (e.g., HIV, hepatitis B and C). Large volumes of transfused blood can lead to volume overload, impacting cardiac function and predisposing individuals to complications like heart failure or pulmonary edema, particularly in patients with compromised cardiac status.

Clinical outcomes and considerations regarding prolonged blood transfusion in the context of postpartum hemorrhage encompass a spectrum of potential effects and strategic approaches for managing patients requiring extended transfusion support.¹¹ Prolonged blood transfusion in postpartum hemorrhage may prevent immediate mortality by restoring blood volume and stability. However, it is associated with increased risks of various morbidities, such as transfusion reactions, infections, and complications related to iron overload. Adverse reactions to transfusions, including febrile non-hemolytic reactions, allergic reactions, acute hemolytic reactions, or transfusion-related circulatory overload (TACO), can contribute to increased maternal morbidity.¹² Despite stringent screening, there remains a small risk of transfusion-transmitted infections, which can pose significant risks to maternal health. Frequent transfusions may lead to iron overload, contributing to complications like liver dysfunction, cardiac issues, endocrine disturbances, and increased oxidative stress. Repeated transfusions can trigger immune responses, resulting in alloimmunization. This may complicate future transfusions, leading to delayed hemolytic reactions or challenges in finding compatible blood products.

This review seeks to comprehensively explore the effects of prolonged blood transfusion in postpartum hemorrhage scenarios. It aims to dissect the physiological alterations induced by extended-term transfusions, delve into associated clinical outcomes, and address critical considerations for healthcare providers involved in managing severe PPH cases requiring prolonged transfusion support.

Postpartum Hemorrhage

Postpartum hemorrhage (PPH) represents a significant obstetric complication and remains a leading cause of maternal mortality and morbidity globally.¹³ The incidence of PPH varies across regions but is notably prevalent in both developed and developing countries.¹⁴ Certain risk factors predispose women to PPH, including uterine atony, retained placental tissue, genital tract trauma, coagulation disorders, and complications arising from instrumental or cesarean deliveries.¹⁵ Identifying these risk factors and promptly managing PPH are critical in preventing adverse

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

maternal outcomes. PPH can manifest within minutes to hours after delivery. Clinical signs include excessive vaginal bleeding, hypotension, tachycardia, and signs of shock. Timely recognition and prompt intervention are pivotal in managing PPH. Initial measures include uterine massage, administration of uterotonic agents (such as oxytocin or misoprostol), and ensuring adequate fluid resuscitation. In severe cases of PPH, where bleeding cannot be controlled by conservative measures alone, blood transfusion becomes a cornerstone in stabilizing the patient.¹⁶ Transfusion of red blood cells, fresh frozen plasma, platelets, and clotting factors may be required to restore hemodynamic stability and correct coagulopathy. While blood transfusion is a life-saving intervention in severe PPH, prolonged or repeated transfusions may introduce complexities and potential risks.¹⁷ Extended-term transfusion in the postpartum period can lead to alterations in hemoglobin levels, disturbances in coagulation, immunomodulation, and complications associated with iron overload or transfusion reactions. These effects necessitate a careful balance between the benefits of transfusion and the potential risks.

Efforts aimed at preventing PPH involve proper antenatal care, skilled birth attendance, active management of the third stage of labor, and timely recognition and management of obstetric complications.¹⁸ Additionally, ongoing research into alternative therapies, transfusion protocols, and strategies to minimize the need for prolonged-term blood transfusions in severe PPH cases remains an area of interest for improving maternal outcomes. Postpartum hemorrhage represents a critical obstetric emergency requiring prompt recognition, early intervention, and appropriate management.¹⁹ While blood transfusion plays a crucial role in severe cases, understanding the implications of prolonged-term transfusion in PPH is essential for optimizing maternal care and improving outcomes in these challenging scenarios.

Blood Transfusion in Postpartum Hemorrhage

Postpartum hemorrhage (PPH) is a serious obstetric complication characterized by excessive bleeding following childbirth, often necessitating urgent interventions to prevent adverse maternal outcomes.²⁰ Blood transfusion stands as a critical therapeutic intervention in cases of severe PPH, aiming to restore blood volume, hemoglobin levels, and coagulation parameters.²¹ However, the administration of blood products in this context poses specific challenges and considerations. In cases of severe PPH, unresponsive to initial management strategies, blood transfusion becomes necessary to correct hypovolemia, maintain tissue oxygenation, and manage coagulopathy. Indications for transfusion include persistent hemorrhage leading to hemodynamic instability, profound anemia, or coagulopathy refractory to other interventions. Choosing the appropriate blood components (packed red blood cells, fresh frozen plasma, platelets, clotting factors) and their ratios is crucial in addressing specific deficits and restoring coagulation function. Tailoring transfusion therapy to individual patient needs while minimizing unnecessary transfusions is essential in preventing complications associated with excessive transfusion. Developing evidence-based transfusion protocols or guidelines specific to postpartum hemorrhage helps standardize transfusion practices, ensuring timely and appropriate administration of blood products. These

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

protocols should consider hemoglobin thresholds, coagulation parameters, and individual patient factors to guide transfusion decisions. Prolonged or repeated blood transfusions in PPH carry inherent risks, including transfusion reactions, transfusion-related infections, volume overload, and immune sensitization.²² Close monitoring of vital signs, laboratory parameters, and clinical assessment for signs of complications is crucial during and after transfusion. In certain scenarios, alternative therapies or adjuncts to blood transfusion might be considered, such as tranexamic acid to aid in coagulation, cell salvage techniques to reinfuse autologous blood, or the use of erythropoietin to stimulate red blood cell production. Effective communication and shared decision-making between healthcare providers and patients are imperative. Informed consent, discussing potential risks and benefits of blood transfusion, respecting patient preferences, and considering cultural or religious beliefs are essential aspects of managing transfusion in PPH. Continued research into transfusion strategies, development of targeted therapies, and innovations in obstetric care are pivotal to optimize transfusion practices and improve outcomes in postpartum hemorrhage scenarios. A multidisciplinary approach involving obstetricians, hematologists, anesthesiologists, and transfusion medicine specialists is essential for comprehensive and effective management. While blood transfusion remains a vital intervention in severe PPH, a tailored and judicious approach considering patient-specific factors, transfusion protocols, alternatives, and ethical considerations is crucial to mitigate risks and improve maternal outcomes.

Physiological Effects of Prolonged Blood Transfusion

Prolonged blood transfusion, especially in the context of postpartum hemorrhage or other conditions requiring extensive transfusion support, can exert various physiological effects on the recipient's body.²³ These effects encompass alterations in hematological parameters, immune modulation, iron metabolism, and potential complications associated with prolonged exposure to transfused blood components. Prolonged blood transfusion can lead to alterations in hemoglobin levels, with subsequent fluctuations in hematocrit and red blood cell indices.²⁴ Repeated transfusions might affect the body's ability to regulate its own erythropoiesis, potentially leading to erythropoietin suppression and a state of relative bone marrow suppression. Blood transfusion may impact the coagulation system, particularly in cases where multiple components such as red blood cells, platelets, and plasma products are transfused. Altered coagulation parameters, including changes in clotting factor levels and platelet function, might occur, potentially affecting hemostasis and contributing to bleeding or thrombotic complications. Transfused blood components can influence the recipient's immune system.²⁵ Prolonged exposure to foreign antigens from transfused blood may result in immune modulation, including immune sensitization or the development of alloantibodies. This can lead to increased risks of transfusion reactions and, in some cases, limit the compatibility of future transfusions. Frequent blood transfusions can contribute to iron overload in recipients, especially in patients who receive transfusions over an extended period.²⁶ This excess iron can deposit in various organs, leading to complications such as organ damage, endocrine dysfunction, and increased oxidative stress. Prolonged-term transfusion increases the risk of transfusion-related adverse events. These complications may include transfusion reactions (such as febrile reactions, allergic reactions, or hemolytic reactions), transfusion-transmitted infections, and transfusion-associated circulatory overload (TACO) due to volume overload. Continuous exposure to transfused blood might suppress endogenous

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

erythropoietin production and lead to bone marrow suppression.²⁷ This could result in decreased production of new blood cells and prolonged dependency on transfusions. Prolonged transfusion may contribute to a state of immune suppression, potentially increasing the risk of infections, particularly in recipients with compromised immune systems or those exposed to multiple transfusions.

Clinical Outcomes and Complications

Prolonged-term blood transfusion in postpartum hemorrhage may mitigate the immediate risk of mortality by stabilizing hemodynamics and restoring blood volume. However, it is associated with an increased risk of various morbidities, including infections, allergic reactions, and transfusion-related acute lung injury (TRALI).²⁸ Adverse reactions to blood transfusions, such as febrile non-hemolytic reactions, allergic reactions, acute hemolytic reactions, or transfusion-related circulatory overload (TACO), can occur and contribute to increased maternal morbidity.²⁹ Despite stringent screening measures, the risk of transfusion-transmitted infections (e.g., HIV, hepatitis B and C, bacterial infections) remains, albeit low. Infections pose a significant concern, especially in settings with inadequate blood screening protocols. Prolonged-term transfusion can lead to iron overload in recipients. This excess iron can accumulate in organs (e.g., liver, heart, pancreas), leading to complications such as liver dysfunction, cardiac issues, endocrine disturbances, and increased oxidative stress. Repeated exposure to transfused blood components can trigger alloimmunization, leading to the development of antibodies against certain blood antigens. This can pose challenges in finding compatible blood for future transfusions and may result in delayed hemolytic transfusion reactions.

Large volumes of transfused blood can result in volume overload, leading to cardiac strain, pulmonary edema, and exacerbation of pre-existing cardiac conditions, particularly in patients with compromised cardiac function.³⁰ Alterations in coagulation parameters due to prolonged-term transfusion might predispose individuals to thrombotic complications such as deep vein thrombosis (DVT) or pulmonary embolism (PE). Prolonged-term transfusion can suppress the immune system, increasing susceptibility to infections, which can be particularly problematic in postpartum women who are already at an increased risk of infections. Prolonged dependency on blood transfusions can lead to delayed recovery, affecting a woman's overall health and potentially impacting future pregnancies due to complications arising from iron overload, alloimmunization, or other transfusion-related issues.³¹

Considerations and Strategies

Certainly, managing prolonged blood transfusion in postpartum hemorrhage requires careful considerations and specific strategies to optimize patient care and outcomes. Establishing evidence-based transfusion protocols and hemoglobin thresholds specific to postpartum hemorrhage aids in guiding transfusion decisions.³² Tailoring transfusion to individual patient needs while minimizing unnecessary transfusions helps prevent complications associated with excessive transfusion. Close monitoring of vital signs, hematological parameters, and clinical status before, during, and after transfusion is essential.³³ This monitoring assists in detecting and

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58

managing potential complications promptly. Considering patient-specific factors, such as baseline hemoglobin levels, comorbidities, and the severity of hemorrhage, is crucial in determining the appropriate transfusion strategy. Customizing care based on individual needs helps mitigate risks and optimize outcomes. Implementing strategies to minimize transfusion reactions involves proper blood component selection, ensuring compatibility, and employing leukoreduction techniques. Adequate pre-transfusion screening and compatibility testing reduce the risk of adverse reactions. Given the risk of iron overload from prolonged transfusions, iron management strategies (e.g., iron chelation therapy) might be considered to mitigate iron-related complications, particularly in patients receiving frequent transfusions. Exploring alternative therapies or adjuncts to blood transfusion, such as tranexamic acid to aid in coagulation, cell salvage techniques, or erythropoietin to stimulate erythropoiesis, can reduce dependency on blood products and minimize associated risks. Engaging patients in informed discussions about the risks and benefits of blood transfusion, discussing potential complications, and addressing concerns or preferences is crucial. Informed consent and clear communication are essential components of patient-centered care. Collaborating with a multidisciplinary team involving obstetricians, hematologists, anesthesiologists, and transfusion medicine specialists ensures comprehensive care, optimal decision-making, and access to specialized expertise in managing complex cases of postpartum hemorrhage requiring prolonged transfusion. Encouraging research efforts to explore innovative transfusion strategies, identify alternative treatments, and enhance transfusion practices in postpartum hemorrhage. Advancements in technology and understanding may lead to improved management approaches. Respecting patient autonomy, cultural or religious beliefs, and ensuring informed consent are critical ethical considerations. Upholding ethical standards in decision-making and care delivery is integral in managing transfusions in postpartum hemorrhage.³⁴⁻⁶⁰

Conclusion

Postpartum hemorrhage (PPH) remains a critical obstetric emergency requiring prompt intervention to prevent maternal morbidity and mortality. In severe cases of PPH, prolonged-term blood transfusion plays a pivotal role in stabilizing hemodynamics and correcting deficits in blood volume and coagulation factors. However, the administration of prolonged blood transfusions in this context necessitates a nuanced approach considering the physiological effects, potential complications, and individualized patient care. The management of prolonged blood transfusion in postpartum hemorrhage necessitates a comprehensive, patient-centered approach, integrating current evidence, multidisciplinary expertise, and a commitment to continually improve care delivery in these critical situations.

References

1. Lancaster L, Barnes RF, Correia M, Luis E, Boaventura I, Silva P, von Drygalski A, Angchaisuksiri P. Maternal death and postpartum hemorrhage in sub-Saharan Africa—A pilot study in metropolitan Mozambique. *Research and practice in thrombosis and haemostasis*. 2020;4(3): e12311.

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. *Elite Journal of Laboratory Medicine*, 2024; 2(2): 47-58

2. Hedrick HL, Flake AW, Crombleholme TM, Howell LJ, Johnson MP, Wilson RD, Adzick NS. Sacrococcygeal teratoma: prenatal assessment, fetal intervention, and outcome. *Journal of pediatric surgery*. 2004;39(3):430-8.
3. Owen MD, Cassidy AL, Weeks AD. Why are women still dying from obstetric hemorrhage? A narrative review of perspectives from high and low resource settings. *International journal of obstetric anesthesia*. 2021; 46:102982.
4. Oyelese Y, Ananth CV. Postpartum hemorrhage: epidemiology, risk factors, and causes. *Clinical obstetrics and gynecology*. 2010;53(1):147-56.
5. Shakur H, Elbourne D, Gülmezoglu M, Alfirevic Z, Ronsmans C, Allen E, Roberts I. The WOMAN Trial (World Maternal Antifibrinolytic Trial): tranexamic acid for the treatment of postpartum haemorrhage: an international randomised, double-blind placebo-controlled trial. *Trials*. 2010;11(1):1-4.
6. Espinoza IS. Obstetric Hemorrhage, its role in maternal morbidity and mortality and the importance of its diagnosis, prevention and timely management. *Mexican Journal of Medical Research ICSA*. 2020;8(15):37-44.
7. Muñoz M, Stensballe J, Ducloy-Bouthors AS, Bonnet MP, De Robertis E, Fornet I, Goffinet F, Hofer S, Holzgreve W, Manrique S, Nizard J. Patient blood management in obstetrics: prevention and treatment of postpartum haemorrhage. A NATA consensus statement. *Blood Transfusion*. 2019;17(2):112.
8. Matkovic E, Lindholm PF. Role of viscoelastic and conventional coagulation tests for management of blood product replacement in the bleeding patient. In *Seminars in thrombosis and hemostasis* 2022; 48(07):785-795). 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA: Thieme Medical Publishers, Inc.
9. Shander A, Cappellini MD, Goodnough LT. Iron overload and toxicity: the hidden risk of multiple blood transfusions. *Vox sanguinis*. 2009;97(3):185-97.
10. Kaphan E, Laurin D, Lafeuillade B, Drillat P, Park S. Impact of transfusion on survival in patients with myelodysplastic syndromes: current knowledge, new insights and transfusion clinical practice. *Blood Reviews*. 2020; 41:100649.
11. Benson G, Auerswald G, Dolan G, Duffy A, Hermans C, Ljung R, Morfini M, Šalek SZ. Diagnosis and care of patients with mild haemophilia: practical recommendations for clinical management. *Blood Transfusion*. 2018;16(6):535.
12. Rahim R, Varghese S, Thambi R, Suma MS. Study on acute transfusion related adverse reactions in surgery department. *Int J Res Med Sci*. 2021;9(5):1381-7.
13. Maswime S, Buchmann E. A systematic review of maternal near miss and mortality due to postpartum hemorrhage. *Int J Gynaecol Obstet*. 2017;137(1):1-7.
14. Prick BW, auf Altenstadt JF, Hukkelhoven CW, Bonsel GJ, Steegers EA, Mol BW, Schutte JM, Bloemenkamp KW, Duvekot JJ. Regional differences in severe postpartum hemorrhage: a nationwide comparative study of 1.6 million deliveries. *BMC pregnancy and childbirth*. 2015; 15:1-0.
15. Nyfløt LT, Sandven I, Stray-Pedersen B, Pettersen S, Al-Zirqi I, Rosenberg M, Jacobsen AF, Vangen S. Risk factors for severe postpartum hemorrhage: a case-control study. *BMC pregnancy and childbirth*. 2017; 17:1-9.

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. *Elite Journal of Laboratory Medicine*, 2024; 2(2): 47-58

16. Sebghati M, Chandraharan E. An update on the risk factors for and management of obstetric haemorrhage. *Women's Health*. 2017;13(2):34-40.
17. Shaikh AG, Shaikh VF, Nasim MT, Farid E, Ali SA, Azim D, Afzal H, Iqbal K, Iqbal A. Risk Factors for Blood Transfusion during Caesarean Section. *Pakistan Journal of Medical & Health Sciences*. 2022;16(04):339-.
18. Bishanga DR, Charles J, Tibaijuka G, Mutayoba R, Drake M, Kim YM, Plotkin M, Rusibamayila N, Rawlins B. Improvement in the active management of the third stage of labor for the prevention of postpartum hemorrhage in Tanzania: a cross-sectional study. *BMC pregnancy and childbirth*. 2018;18(1):1-0.
19. Traoré Y, Tégoué I, Bocoum A, Traoré M, Dao S, Bomini MK, Mounkoro N, Dolo A. Management and Prognosis of Early Postpartum Hemorrhage in African Low Setting Health. *Open Journal of Obstetrics and Gynecology*. 2018;8(1):1-9.
20. Ghosh M, Chandraharan E. Management of post-partum haemorrhage. *Obstetrics, Gynaecology & Reproductive Medicine*. 2017;27(8):239-44.
21. Stanworth SJ, Dowling K, Curry N, Doughty H, Hunt BJ, Fraser L, Narayan S, Smith J, Sullivan I, Green L, Transfusion Task Force of the British Society for Haematology. Haematological management of major haemorrhage: a British Society for Haematology Guideline. *British journal of haematology*. 2022;198(4):654-67.
22. Kozek-Langenecker SA, Ahmed AB, Afshari A, Albaladejo P, Aldecoa C, Barauskas G, De Robertis E, Faraoni D, Filipescu DC, Fries D, Haas T. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology: first update 2016. *European Journal of Anaesthesiology| EJA*. 2017;34(6):332-95.
23. Rossaint R, Bouillon B, Cerny V, Coats TJ, Duranteau J, Fernández-Mondéjar E, Hunt BJ, Komadina R, Nardi G, Neugebauer E, Ozier Y. Management of bleeding following major trauma: an updated European guideline. *Critical care*. 2010; 14:1-29.
24. García-Roa M, del Carmen Vicente-Ayuso M, Bobes AM, Pedraza AC, González-Fernández A, Martín MP, Sáez I, Seghatchian J, Gutiérrez L. Red blood cell storage time and transfusion: current practice, concerns and future perspectives. *Blood Transfusion*. 2017;15(3):222.
25. Brand A. Immunological aspects of blood transfusions. *Transplant immunology*. 2002;10(2-3):183-90.
26. Shander A, Cappellini MD, Goodnough LT. Iron overload and toxicity: the hidden risk of multiple blood transfusions. *Vox sanguinis*. 2009;97(3):185-97.
27. Jelkmann W. Erythropoietin after a century of research: younger than ever. *European journal of haematology*. 2007;78(3):183-205.
28. Kuldanek SA, Kelher M, Silliman CC. Risk factors, management and prevention of transfusion-related acute lung injury: a comprehensive update. *Expert review of hematology*. 2019;12(9):773-85.
29. Rahim R, Varghese S, Thambi R, Suma MS. Study on acute transfusion related adverse reactions in surgery department. *Int J Res Med Sci*. 2021;9(5):1381-7.
30. Bulle EB, Klanderman RB, Pendergrast J, Cserti-Gazdewich C, Callum J, Vlaar AP. The recipe for TACO: A narrative review on the pathophysiology and potential mitigation

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. *Elite Journal of Laboratory Medicine*, 2024; 2(2): 47-58

- strategies of transfusion-associated circulatory overload. *Blood Reviews*. 2022; 52:100891.
31. Chauhan S, Rishi B, Tanwar P, Mehdi G, Arif SH, Rabbani T, Rai S, Siraj F, Misra A. Therapeutic lessons from transfusion in pregnancy-effect on hematological parameters and coagulation profile. *American Journal of Blood Research*. 2021;11(3):303.
 32. Mo YD, Delaney M. Transfusion in pediatric patients: review of evidence-based guidelines. *Clinics in Laboratory Medicine*. 2021;41(1):1-4.
 33. Gehrie EA, Roubinian NH, Chowdhury D, Brambilla DJ, Murphy EL, Gottschall JL, Wu Y, Ness PM, Strauss RG, Hendrickson JE, NHLBI Recipient Epidemiology-Donor Evaluation Study (REDS-III). A multicentre study investigating vital sign changes occurring in complicated and uncomplicated transfusions. *Vox sanguinis*. 2018;113(2):160-169.
 34. Obeagu EI, Obeagu GU, Obiezu J, Ezeonwumelu C, Ogunnaya FU, Ngwoke AO, Emeka-Obi OR, Ugwu OP. Hematologic Support in HIV Patients: Blood Transfusion Strategies and Immunological Considerations. *Applied Sciences (NIJBAS)*. 2023;3(3).
 35. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. *Elite Journal of Laboratory Medicine*. 2024;2(1):33-45.
 36. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. *Elite Journal of Laboratory Medicine*. 2024;2(1):33-45.
 37. Obeagu EI, Obeagu GU. The Role of Blood Transfusion Strategies in HIV Management: Current Insights and Future Directions. *Elite Journal of Medicine*. 2024;2(1):10-22.
 38. Obeagu EI, Obeagu GU. Advances in Understanding the Impact of Blood Transfusion on Anemia Resolution in HIV-Positive Children with Severe Malaria: A Comprehensive Review. *Elite Journal of Haematology*. 2024;2(1):26-41.
 39. Obeagu EI, Babar Q, Obeagu GU. Allergic blood Transfusion reaction: A Review. *Int. J. Curr. Res. Med. Sci.* 2021;7(5):25-33.
 40. Obeagu EI, Ubosi NI, Uzoma G. Maternal Hemorrhage and Blood Transfusions: Safeguarding Pregnancy Health. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(11):26-35.
 41. Obeagu EI, Obeagu GU. Transfusion-Related Complications in Children Under 5 with Coexisting HIV and Severe Malaria: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2024;11(2):9-19.
 42. ObeaguEI AA, Obeagu GU. Synergistic Effects of Blood Transfusion and HIV in Children Under 5 Years with Severe Malaria: A Review. *Elite Journal of HIV*. 2024;2(1):31-50.
 43. Obeagu EI, Anyiam AF, Obeagu GU. Managing Anemia in HIV through Blood Transfusions: Clinical Considerations and Innovations. *Elite Journal of HIV*. 2024;2(1):16-30.
 44. Obeagu EI, Obeagu GU. Transfusion Therapy in HIV: Risk Mitigation and Benefits for Improved Patient Outcomes. *Sciences*. 2024;4(1):32-37.
 45. Obeagu EI, Obeagu GU, Obiezu J, Ezeonwumelu C, Ogunnaya FU, Ngwoke AO, Ugwu OP. Immunomodulatory Effects of Transfusions on Maternal Immunity in Pregnancy. *Applied Sciences (NIJBAS)*. 2023;3(3).

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. *Elite Journal of Laboratory Medicine*, 2024; 2(2): 47-58

46. Obeagu EI, Babar Q, Uduchi IO, Ibekwe AM, Chijioke UO, Okafor CJ, Vincent CC. An Update on Transfusion Related Immunomodulation (TRIM) in a Time of COVID-19 Pandemic. *Journal of Pharmaceutical Research International*. 2021 Aug 27;33(42A):135-146.
47. Okoroiwu IL, Obeagu EI, Elemchukwu Q, Ochei KC, Christian GS. Frequency of Transfusion Reactions Following Compatible Cross Matching of Blood: A Study in Owerri Metropolis. *International Journal of Current Research and Academic Review*. 2015;3(1):155-160.
48. Obeagu EI, Oshim IO, Ochei KC, Obeagu GU. Iron and blood donation: A Review. *Int. J. Curr. Res. Med. Sci*. 2016;2(10):16-48.
49. Ogar CO, Okoroiwu HU, Obeagu EI, Etura JE, Abunimye DA. Assessment of blood supply and usage pre-and during COVID-19 pandemic: a lesson from non-voluntary donation. *Transfusion Clinique et Biologique*. 2021;28(1):68-72.
50. Anyiam AF, Arinze-Anyiam OC, Ironi EA, Obeagu EI. Distribution of ABO and rhesus blood grouping with HIV infection among blood donors in Ekiti State Nigeria. *Medicine*. 2023;102(47): e36342.
51. Obeagu EI. Blood Transfusion: A Powerful Process of Saving Anaemic Patients. *EC Emergency Medicine and Critical Care*. 2020;4(7):33-40.
52. Obeagu EI, Buhari HA. Implications of Blood Transfusion in Renal Disease Patients. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(10):45-49.
53. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Ironi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal*. 2022;6(3):9-17.
54. Obeagu EI, Obeagu GU, Ukibe NR, Oyebadejo SA. Anemia, iron, and HIV: decoding the interconnected pathways: A review. *Medicine*. 2024;103(2): e36937.
55. Obeagu EI. An update on susceptibility of individuals to diseases based on ABO blood groups. *Int. J. Curr. Res. Med. Sci*. 2019;5(3):1-8.
56. Eze R, Obeagu EI, Nwakulite A, Vincent CC, Ogbodo SO, Ibekwe AM, Okafor CJ, Chukwurah EF. Frequency of Haemoglobin Genotype Variants, ABO and Rh 'D' Antigen among Madonna Undergraduates of South East Origin, Nigeria. *Journal of Pharmaceutical Research International*. 2021 May 26;33(29B):149-57.
57. Okoroiwu IL, Obeagu EI, Christian SG, Elemchukwu Q, Ochei KC. Determination of the haemoglobin, genotype and ABO blood group pattern of some students of Imo State University, Owerri, Nigeria. *International Journal of Current Research and Academic Review*. 2015;3(1):20-27.
58. Oloro OH, Obeagu EI, Puche RO, Lawal OA. Blood Products in Blood Banking: Preparation and Clinical Importance. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(3):102-109.
59. Asemota EA, Njar VE, Aguanah IT, Obeagu EI. Distribution of ABO, Rhesus Blood Group and Helicobacter Pylori Infection among Secondary School Students in Calabar South Local Government, Cross River State, Nigeria. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2023;3(1):32-45.

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. *Elite Journal of Laboratory Medicine*, 2024; 2(2): 47-58

60. Obeagu EI, Katya MC. A Systematic Review on Physiological Jaundice: Diagnosis and Management of the Affected Neonates. Madonna University journal of Medicine and Health Sciences. 2022;2(3):25-41.

Citation: Obeagu EI, Obeagu GU. The effects of prolonged term blood transfusion in Postpartum Hemorrhage. Elite Journal of Laboratory Medicine, 2024; 2(2): 47-58