

Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges

*Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²

¹Department of Medical Laboratory Science, Kampala International University, Ishaka, Uganda.

²School of Nursing Science, Kampala International University, Ishaka, Uganda.

*Corresponding author: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, \[emmanuelobeagu@yahoo.com\]\(mailto:emmanuelobeagu@yahoo.com\), ORCID: 0000-0002-4538-0161](#)

Abstract

The co-occurrence of malaria and Hepatitis B during pregnancy presents a complex interplay of challenges that impact maternal and neonatal health. Both infections individually pose significant risks: malaria can lead to adverse pregnancy outcomes such as preterm birth and low birth weight, while Hepatitis B increases the risk of chronic liver disease and potential transmission to the infant. This review explores the intersection of these two infections, examining their combined effects on pregnancy and the implications for maternal and child health. The review also summarizes epidemiological evidence on the prevalence of co-infection and its impact on pregnancy outcomes, including increased risks of severe disease and poor neonatal health. Public health interventions are crucial in addressing the dual burden of malaria and Hepatitis B. Integrated care approaches, including effective malaria prevention measures and Hepatitis B vaccination, are essential for mitigating the risks associated with co-infection. Enhancing antenatal care, community-based education, and ongoing research are key components for improving health outcomes for pregnant women and their infants.

Keywords: *Malaria, Pregnancy, Hepatitis B, Co-infection, Maternal Health, Neonatal Outcomes, Public Health Interventions, Integrated Care*

Introduction

Malaria and Hepatitis B are two major infectious diseases with significant global health impacts. Malaria, caused by Plasmodium parasites transmitted through Anopheles mosquitoes, affects millions of people annually, with a disproportionate burden in sub-Saharan Africa. Hepatitis B, caused by the Hepatitis B virus (HBV), is also prevalent in many regions, particularly in parts of Asia and Africa. Both infections pose considerable health risks, and their effects are magnified during pregnancy, making the management of these co-existing conditions a critical public health challenge. Malaria in pregnancy is associated with a range of complications, including anemia, **Citation:** Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. Elite Journal of Health Science, 2024; 2(7): 1-13

low birth weight, preterm birth, and maternal mortality. The parasites tend to sequester in the placenta, leading to inflammation and impaired nutrient transfer to the fetus. This disruption can result in adverse outcomes not only for the mother but also for the newborn. The impact of malaria on pregnancy outcomes emphasizes the need for effective prevention and treatment strategies to protect maternal and neonatal health. Hepatitis B infection during pregnancy presents its own set of challenges. Chronic Hepatitis B can lead to significant liver damage and increases the risk of liver-related complications. Additionally, the risk of vertical transmission from mother to infant is a major concern, as HBV can be transmitted during childbirth and potentially lead to chronic infection in the child. This vertical transmission can result in long-term health issues, including chronic liver disease and liver cancer.¹⁻¹⁰

The co-occurrence of malaria and Hepatitis B during pregnancy complicates the clinical management of both infections. Malaria-induced inflammation in the liver can exacerbate Hepatitis B complications, while the chronic effects of Hepatitis B can influence the severity and management of malaria. The pathophysiology of malaria and Hepatitis B during pregnancy involves complex interactions between the immune system, inflammatory responses, and the effects on liver function. Malaria can lead to liver inflammation and altered immune responses, which can complicate Hepatitis B management. Conversely, Hepatitis B-related liver damage can influence the course of malaria infection and its treatment. Exploring these mechanisms helps to elucidate the compounded health risks associated with co-infection. Epidemiological studies have provided valuable insights into the prevalence and impact of co-infection with malaria and Hepatitis B. Research indicates that these infections often overlap in endemic regions, with significant implications for maternal and child health. Evidence suggests that co-infection is associated with increased risks of severe disease outcomes and complications, highlighting the need for targeted public health interventions. Addressing the dual burden of malaria and Hepatitis B requires comprehensive public health interventions. Effective strategies include the use of insecticide-treated bed nets (ITNs) and intermittent preventive treatment in pregnancy (IPTp) for malaria, alongside Hepatitis B vaccination and antiviral therapy. Integrated care approaches that encompass both malaria and Hepatitis B management are essential for improving health outcomes for pregnant women and their infants. Community-based approaches and health education play a crucial role in managing the co-occurrence of malaria and Hepatitis B. Engaging communities in health promotion, providing information on prevention, and enhancing access to healthcare services are key components of effective public health strategies. Community health workers and local health programs are instrumental in delivering education and facilitating access to necessary interventions.¹¹⁻²⁰

Each condition affects the body in distinct ways, but their co-occurrence can exacerbate health complications. During pregnancy, Plasmodium parasites can sequester in the placenta, leading to localized inflammation and impaired placental function. This sequestration disrupts the transfer of nutrients and oxygen to the fetus, contributing to adverse outcomes such as low birth weight and preterm birth. The inflammatory response in the placenta also affects maternal immune responses and can lead to complications in malaria management. The inflammatory response triggered by malaria can extend to the liver, where the parasites can induce hepatitis-like symptoms. This inflammation can exacerbate existing liver conditions and complicate the management of Hepatitis

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. Elite Journal of Health Science, 2024; 2(7): 1-13

B. The liver's role in metabolizing drugs used to treat malaria may also be compromised, affecting treatment efficacy and safety. Hepatitis B infection leads to chronic inflammation of the liver, which can progress to cirrhosis and liver failure if untreated. During pregnancy, the increased liver burden from Hepatitis B can be compounded by malaria-induced liver inflammation. This chronic inflammation can affect the liver's ability to process medications and manage metabolic functions, complicating the treatment of both infections. One of the major concerns with Hepatitis B during pregnancy is the risk of vertical transmission. HBV can be transmitted from mother to child during delivery, leading to potential chronic infection in the infant. The presence of malaria can further complicate the delivery process and potentially increase the risk of transmission. Effective management of Hepatitis B is essential to minimize this risk and ensure the health of the infant.²¹⁻

30

The co-existence of malaria and Hepatitis B can lead to compounded inflammation and immune dysfunction. Malaria-induced hepatic inflammation can worsen Hepatitis B-related liver damage, while chronic Hepatitis B can alter immune responses, affecting the course and treatment of malaria. This interaction creates a complex clinical scenario where each condition can exacerbate the severity of the other. The presence of both malaria and Hepatitis B can influence the pharmacokinetics of treatments. Malaria medications, which are metabolized in the liver, may interact with antiviral drugs used for Hepatitis B. Liver dysfunction due to either infection can affect drug metabolism, leading to potential drug interactions and altered treatment efficacy. The combined burden of malaria and Hepatitis B can significantly impact maternal health. The increased liver inflammation from both infections can lead to severe complications, including liver failure and preterm labor. Managing these complications requires careful monitoring and integrated treatment strategies to address the multifaceted health challenges. For the fetus, the combined effects of malaria and Hepatitis B can lead to increased risks of adverse outcomes such as intrauterine growth restriction, preterm birth, and low birth weight. Additionally, the risk of vertical transmission of Hepatitis B and the potential for severe malaria-related complications highlight the need for effective prenatal care and interventions. Effective management of malaria and Hepatitis B during pregnancy requires coordinated care approaches. This includes integrating malaria prevention and treatment with Hepatitis B management strategies to address both conditions simultaneously. Regular monitoring, appropriate medication adjustments, and supportive care are essential for improving health outcomes for both mother and infant.³¹⁻⁴⁰

Epidemiological Evidence

Epidemiological evidence provides crucial insights into the prevalence, impact, and interactions of malaria and Hepatitis B during pregnancy. The co-occurrence of malaria and Hepatitis B is notably prevalent in regions where both infections are endemic. Sub-Saharan Africa and parts of Asia are significant hotspots for these diseases. Studies show that in malaria-endemic areas, the prevalence of Hepatitis B among pregnant women can range from 5% to 15%, with varying rates depending on specific geographic locations and population characteristics. The overlap in endemic areas underscores the need for integrated health strategies. Epidemiological surveys have revealed that the prevalence of co-infection varies within regions, influenced by factors such as local healthcare infrastructure, socio-economic conditions, and public health initiatives. For example,

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

in certain high-risk areas of West Africa, the co-infection rates among pregnant women can exceed 10%, indicating a significant public health challenge. Evidence suggests that the co-occurrence of malaria and Hepatitis B can lead to severe adverse outcomes during pregnancy. Studies have shown increased risks of preterm birth, low birth weight, and maternal anemia in women with both infections. The presence of both conditions exacerbates complications, potentially leading to poorer health outcomes for both mother and child. Research indicates that pregnant women co-infected with malaria and Hepatitis B face higher risks of severe disease and complications. Malaria can aggravate liver inflammation due to Hepatitis B, increasing the likelihood of severe maternal morbidity. Additionally, there is evidence of elevated maternal mortality rates in women with both infections, highlighting the need for targeted healthcare interventions. Infants born to mothers with co-infection are at increased risk of adverse outcomes, including intrauterine growth restriction and neonatal malaria. The risk of vertical transmission of Hepatitis B to the newborn is a major concern, with potential long-term implications for the child's health. Studies have shown that effective management of maternal Hepatitis B can significantly reduce the risk of transmission to the infant.⁴¹⁻⁵⁰

Epidemiological data reveal that the interaction between malaria and Hepatitis B can lead to compounded health risks. For example, the inflammatory effects of malaria can worsen liver damage in Hepatitis B patients, while chronic Hepatitis B can alter the severity and management of malaria. This interaction complicates clinical management and underscores the need for integrated treatment approaches. Studies evaluating the effectiveness of public health interventions for managing malaria and Hepatitis B co-infection provide valuable insights. Evidence shows that integrated care approaches, including combined prevention strategies and coordinated treatment, improve health outcomes. For example, the use of insecticide-treated bed nets (ITNs) alongside Hepatitis B vaccination programs has shown promise in reducing the burden of both infections. The integration of malaria and Hepatitis B control programs has been shown to enhance the effectiveness of interventions. Public health strategies that combine malaria prevention measures, such as ITNs and intermittent preventive treatment in pregnancy (IPTp), with Hepatitis B vaccination and antiviral therapy are crucial for addressing the dual burden of these infections. Regular screening for both malaria and Hepatitis B during pregnancy is essential for early detection and management. Evidence supports the implementation of routine antenatal care programs that include testing for both infections, providing appropriate treatment, and offering counseling to pregnant women. Community health education and engagement play a critical role in managing co-infections. Public health campaigns aimed at increasing awareness, promoting prevention measures, and improving access to healthcare services are important for reducing the prevalence and impact of malaria and Hepatitis B.⁵¹⁻⁶⁰

Public Health Interventions

Addressing the dual challenges of malaria and Hepatitis B during pregnancy requires multifaceted public health interventions. Effective strategies involve not only the prevention and treatment of each infection individually but also the integration of care approaches to manage the compounded effects of co-infection. Integrated control programs that address both malaria and Hepatitis B are essential for effective management. For malaria, strategies include the widespread use of

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

insecticide-treated bed nets (ITNs), intermittent preventive treatment in pregnancy (IPTp), and prompt diagnosis and treatment of malaria cases. For Hepatitis B, prevention focuses on vaccination, antiviral therapy, and screening pregnant women to identify and manage chronic infections. Implementing combined prevention strategies can enhance the effectiveness of public health efforts. For example, integrating Hepatitis B vaccination programs with malaria prevention initiatives, such as ITNs and IPTp, can simultaneously address both infections. Coordinated care ensures that pregnant women receive comprehensive health services, reducing the risk of complications associated with co-infection. Regular screening for malaria and Hepatitis B during antenatal care is crucial for early detection and management. Pregnant women should be routinely tested for both infections to identify those at risk and provide timely interventions. Screening helps in monitoring the health of the mother and fetus, enabling targeted treatment and reducing the risk of adverse outcomes. Antenatal care should include integrated management strategies that address both malaria and Hepatitis B. This involves providing appropriate treatment for malaria, managing Hepatitis B to prevent vertical transmission, and offering supportive care to address complications. Healthcare providers should be trained to recognize and manage co-infection effectively.⁶¹⁻⁷⁸

Hepatitis B vaccination is a key intervention to prevent vertical transmission and reduce the incidence of chronic Hepatitis B infection in infants. Vaccination programs should ensure that all pregnant women at risk of Hepatitis B receive the vaccine, and infants born to Hepatitis B-positive mothers should receive the hepatitis B immunoglobulin (HBIG) and vaccine shortly after birth. Although there is no widely available malaria vaccine for pregnant women, research into malaria vaccines is ongoing. Efforts are focused on developing vaccines that are safe and effective for pregnant women and their infants. Future availability of such vaccines could significantly impact malaria prevention strategies. Community-based health education plays a vital role in managing malaria and Hepatitis B. Public health campaigns should focus on raising awareness about the prevention and treatment of both infections. Educating communities about the importance of antenatal care, vaccination, and effective malaria prevention measures can improve health outcomes. Community health workers are instrumental in delivering education, providing screenings, and facilitating access to healthcare services. Training and supporting local health workers to address both malaria and Hepatitis B can enhance the reach and effectiveness of public health interventions. Effective public health interventions require supportive policies and resources. Governments and health organizations should develop and implement policies that prioritize the prevention and treatment of malaria and Hepatitis B during pregnancy. Funding and resources should be allocated to support integrated care programs and improve healthcare infrastructure. Ongoing monitoring and evaluation of public health interventions are essential for assessing their impact and effectiveness. Regular data collection and analysis help identify gaps, measure progress, and inform future strategies. Evaluating the outcomes of integrated care programs can guide improvements and ensure that interventions are meeting their goals.⁷⁹⁻⁸⁹

Conclusion

The co-occurrence of malaria and Hepatitis B during pregnancy presents significant challenges that require integrated and multifaceted public health interventions. Both infections independently pose considerable health risks, but their combined effects can exacerbate complications for both

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

the mother and the fetus. Addressing this dual burden effectively necessitates a comprehensive approach that incorporates prevention, diagnosis, treatment, and supportive care. The pathophysiological mechanisms underlying malaria and Hepatitis B co-infection highlight the complex interactions between these diseases. Malaria-induced inflammation can aggravate Hepatitis B-related liver damage, while chronic Hepatitis B can influence the severity and management of malaria. Recognizing these interactions is crucial for developing effective treatment strategies that address the compounded health risks.

Epidemiological evidence underscores the prevalence and impact of co-infection in endemic regions. Data reveal that the simultaneous presence of malaria and Hepatitis B increases the risk of severe maternal and neonatal outcomes, including preterm birth, low birth weight, and vertical transmission of Hepatitis B. Effective public health interventions must consider these increased risks and target both infections to improve health outcomes. Integrated public health interventions are essential for managing the dual burden of malaria and Hepatitis B. Coordinated prevention efforts, such as combined malaria control measures and Hepatitis B vaccination programs, enhance the effectiveness of interventions. Routine antenatal screening and care, community-based education, and supportive health policies play crucial roles in reducing the impact of these infections.

References

1. Uneke CJ. Impact of placental *Plasmodium falciparum* malaria on pregnancy and perinatal outcome in sub-Saharan Africa: part III: placental malaria, maternal health, and public health. *The Yale journal of biology and medicine*. 2008;81(1):1.
2. Gontie GB, Wolde HF, Baraki AG. Prevalence and associated factors of malaria among pregnant women in Sherkole district, Benishangul Gumuz regional state, West Ethiopia. *BMC Infectious Diseases*. 2020; 20:1-8.
3. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri*. 2023; 6 (1). 2023;138. links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf.
4. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci*. 2016;3(3):10-8. links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf.
5. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. *J Res Med Dent Sci*. 2021;9(4):145-8. links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf.
6. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
7. Jakheng SP, Obeagu EI. Seroprevalence of human immunodeficiency virus based on demographic and risk factors among pregnant women attending clinics in Zaria Metropolis, Nigeria. *J Pub Health Nutri*. 2022; 5 (8). 2022;137.

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

[links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf](https://epjournals.com/journals/EJHS/links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf).

8. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. Madonna University journal of Medicine and Health Sciences. 2022;2(2):1-9.
9. Bonilla FA, Oettgen HC. Adaptive immunity. Journal of Allergy and Clinical Immunology. 2010;125(2): S33-40.
10. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):1-9.
11. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfectd with HIV in Umuahia. Int. J. Curr. Res. Med. Sci. 2017;3(5):100-104.
12. Feeney ME. The immune response to malaria in utero. Immunological reviews. 2020 ;293(1):216-229.
13. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. Medicine. 2023;102(46): e36166.
14. Obeagu EI, Chijioke UO, Ekelozie IS. Malaria rapid diagnostic test (RDTs). Ann Clin Lab Res. 2018;6(4):275.
15. Ogomaka IA, Obeagu EI. Methods of Breast Feeding as Determinants of Malaria Infections among Babies in IMO State, Nigeria. International Journal of Medical Science and Dental Research. 2019;2(01):17-24.
16. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. Madonna University Journal of Medicine and Health Sciences. 2022;2(1):292-8.
<https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
17. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. J. Bio. Innov11 (3). 2022:936-941.
18. Obeagu EI, Obeagu GU, Egba SI, Emeka-Obi OR. Combatting Anemia in Pediatric Malaria: Effective Management Strategies. Int. J. Curr. Res. Med. Sci. 2023;9(11):1-7.
19. Hassan AO, Oso OV, Obeagu EI, Adeyemo AT. Malaria Vaccine: Prospects and Challenges. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):22-40.
20. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. Journal of Pharmaceutical Research International. 2021;33(4):10-9.
21. Moya-Alvarez V, Abellana R, Cot M. Pregnancy-associated malaria and malaria in infants: an old problem with present consequences. Malaria journal. 2014; 13:1-10.
22. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. Int. J. Curr. Res. Chem. Pharm. Sci. 2022;9(9): 21-6.DOI: 10.22192/ijcrps.2022.09.09.003

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. Elite Journal of Health Science, 2024; 2(7): 1-13

23. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
24. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State. Nigeria. *Journal of Research in Medical and Dental Science*. 2021;9(12):462-. [links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://doi.org/10.31838/JRMDS.2021.9.12.462).
25. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022 Jun 21;5(2):113-121.
26. Obeagu EI. Gestational Thrombocytopaenia. *J Gynecol Women's Health*. 2023;25(3):556163. [links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf](https://doi.org/10.31838/JGWH.2023.25.3.556163).
27. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
28. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
29. Obeagu EI, Busari AI, Uduchi IO, Ogomaka IA, Ibekwe AM, Vincent CC, Chijioke UO, Okafor CJ, Okoroiwu HU, Adike CN. Age-Related Haematological Variations in Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):218-24.
30. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
31. Ogbonna CO, Obeagu EI, Ufelle SA, Ogbonna LN. Evaluation of haematological alterations in children infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(1):38-45.
32. Appay V. The physiological role of cytotoxic CD4+ T-cells: the holy grail? *Clinical & Experimental Immunology*. 2004;138(1):10-13.
33. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(1):25-36.
34. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med*. 2018;3(9):1-4.
35. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

- Plasmodium falciparum malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2016;3(8):40-46.
36. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology.* 2019;41(3): e12573.
 37. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research.* 2017;5(4):206.
 38. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Ironi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal.* 2022;6(3):9-17.
 39. Madekwe CC, Madekwe CC, Obeagu EI. Inequality of monitoring in Human Immunodeficiency Virus, Tuberculosis and Malaria: A Review. *Madonna University journal of Medicine and Health Sciences.* 2022;2(3):6-15.
 40. Offie DC, Ibekwe AM, Agu CC, Esimai BN, Okpala PU, Obeagu EI, Ufelle SA, Ogbonna LN. Fibrinogen and C-Reactive Protein Significance in Children Infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(15):1-8.
 41. Obeagu EI, Ogunnaya FU. PREGNANCYINDUCED HAEMATOLOGICAL CHANGES: A KEY TO MATERNAL AND CHILD HEALTH. *European Journal of Biomedical.* 2023;10(8):42-43. [links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf](https://doi.org/10.22192/ijcrms.2023.09.02.003)
 42. Obeagu EI, Ofodile AC, Okwuanaso CB. A review of urinary tract infections in pregnant women: Risks factors. *J Pub Health Nutri.* 2023; 6 (1). 2023; 137:26-35. [links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf](https://doi.org/10.22192/ijcrms.2023.09.02.003).
 43. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. *Int. J. Curr. Res. Med. Sci.* 2023;9(2): 14-17.DOI: 10.22192/ijcrms.2023.09.02.003
 44. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 14-21.DOI: 10.22192/ijcreps.2023.10.08.002
 45. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology.* 2018; 9:1733.
 46. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9): 1-4.DOI: 10.22192/ijcrbm.2018.03.09.001
 47. Onyenweaku FC, Amah HC, Obeagu EI, Nwandikor UU, Onwuasoanya UF. Prevalence of asymptomatic bacteriuria and its antibiotic susceptibility pattern in pregnant women attending private ante natal clinics in Umuahia Metropolitan. *Int J Curr Res Biol Med.* 2017;2(2): 13-23.DOI: 10.22192/ijcrbm.2017.02.02.003

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

48. Okoroiwu IL, Chinedu-Madu JU, Obeagu EI, Vincent CC, Ochiabuto OM, Ibekwe AM, Amaechi CO, Agu CC, Anoh NV, Amadi NM. Evaluation of Iron Status, Haemoglobin and Protein Levels of Pregnant Women in Owerri Metropolis. *Journal of Pharmaceutical Research International*. 2021;33(27A):36-43.
49. Obeagu EI, Njar VE, Obeagu GU. Infertility: Prevalence and Consequences. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(7):43-50.
50. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
51. Broen K, Brustoski K, Engelmann I, Luty AJ. Placental Plasmodium falciparum infection: causes and consequences of in utero sensitization to parasite antigens. *Molecular and biochemical parasitology*. 2007;151(1):1-8.
52. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9):41-49.
53. Ogbonna LN, Ezeoru VC, Ofodile AC, Ochiabuto OM, Obi-Ezeani CN, Okpala PU, Okafor CJ, Obeagu GU, Busari AI, Obeagu EI. Gender Based Variations of Haematological Parameters of Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(8):75-80.
54. Eberendu IF, Ozims SJ, Agu GC, Amah HC, Obasi CC, Obioma-Elemba JE, Ihekaire DE, Ibanga IE, Amah CC, Obeagu EI, Nwosu DC. Impact of human activities on the breeding of mosquitoes of human disease in Owerri metropolis, Imo state. *Int J Adv Res Biol Sci IJARBS*. 2017;4(12):98-106.
55. Obeagu EI, Ofodile AC, Okwuanaso CB. A review on socio economic and behavioral aspects of malaria and its control among children under 5 years of age in Africa. *J Pub Health Nutri*. 2023; 6 (1): 136.
56. Djontu JC, Siewe Siewe S, Mpeke Edene YD, Nana BC, Chomga Foko EV, Bigoga JD, Leke RF, Megnekou R. Impact of placental Plasmodium falciparum malaria infection on the Cameroonian maternal and neonate's plasma levels of some cytokines known to regulate T cells differentiation and function. *Malaria journal*. 2016; 15:1-1.
57. Obeagu EI, Faduma MH, Uzoma G. Ectopic Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(4): 40-4.DOI: [10.22192/ijcrps.2023.10.04.004](https://doi.org/10.22192/ijcrps.2023.10.04.004)
58. Obeagu EI, Gamade SM, Obeagu GU. The roles of Neutrophils in pregnancy. *Int. J. Curr. Res. Med. Sci*. 2023;9(5): 31-35.DOI: [10.22192/ijcrms.2023.09.05.005](https://doi.org/10.22192/ijcrms.2023.09.05.005)
59. Obeagu EI, Obeagu GU. Molar Pregnancy: Update of prevalence and risk factors. *Int. J. Curr. Res. Med. Sci*. 2023;9(7): 25-28.DOI: [10.22192/ijcrms.2023.09.07.005](https://doi.org/10.22192/ijcrms.2023.09.07.005)
60. Kabyemela E, Gonçalves BP, Prevots DR, Morrison R, Harrington W, Gwamaka M, Kurtis JD, Fried M, Duffy PE. Cytokine profiles at birth predict malaria severity during infancy. *PloS one*. 2013;8(10):e77214.
61. Ibebuikie JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Barriers to utilization of maternal health services in southern senatorial district of Cross Rivers state, Nigeria. *International Journal of Advanced Multidisciplinary Research*. 2017;4(8): 1-9.DOI: [10.22192/ijamr.2017.04.08.001](https://doi.org/10.22192/ijamr.2017.04.08.001)

Citation: Obeagu EI, Obeagu GU. Malaria, Pregnancy, and Hepatitis B: Coexisting Challenges. *Elite Journal of Health Science*, 2024; 2(7): 1-13

62. Emmanuel G, Martin O, Peter OS, Obeagu EI, Daniel K. Factors Influencing Early Neonatal Adverse Outcomes among Women with HIV with Post Dated Pregnancies Delivering at Kampala International University Teaching Hospital, Uganda. *Asian Journal of Pregnancy and Childbirth*. 2023;6(1):203-211.
<http://research.sdpublishers.net/id/eprint/2819/>.
63. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9): 41-9.DOI: [10.22192/ijcrms.2018.04.09.006](https://doi.org/10.22192/ijcrms.2018.04.09.006)
64. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8): 22-6.DOI: [10.22192/ijcrmps.2023.10.08.003](https://doi.org/10.22192/ijcrmps.2023.10.08.003)
65. Obeagu E, Eze RI, Obeagu EI, Nnatuanya IN, Dara EC. ZINC LEVEL IN APPARENTLY PREGNANT WOMEN IN URBAN AREA. *Madonna University journal of Medicine and Health Sciences*. 2022;2(1):134-48.
<https://www.journal.madonnauniversity.edu.ng/index.php/medicine/article/view/40>.
66. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
67. Obeagu EI, Ogunnaya FU, Obeagu GU, Ndidi AC. SICKLE CELL ANAEMIA: A GESTATIONAL ENIGMA. migration. 2023; 17:18.
68. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology*. 2019;41(3):e12573.
69. Ifeanyi OE, Uzoma OG. A review on erythropietin in pregnancy. *J. Gynecol. Womens Health*. 2018;8(3):1-4.
https://www.academia.edu/download/56538560/A_Review_on_Erythropietin_in_Pregnancy.pdf.
70. Ifeanyi OE. A review on pregnancy and haematology. *Int. J. Curr. Res. Biol. Med*. 2018;3(5): 26-8.DOI: [10.22192/ijcrbm.2018.03.05.006](https://doi.org/10.22192/ijcrbm.2018.03.05.006)
71. Nwosu DC, Nwanjo HU, Obeagu EI, Ibebuike JE, Ezeama MC. Ihekireh. Changes in liver enzymes and lipid profile of pregnant women with malaria in Owerri, Nigeria. *International Journal of Current Research and Academic Review*. 2015;3(5):376-383.
72. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Factors that influence women's utilization of primary health care services in Calabar Cros river state, Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2017;4(7):28-33.
73. Elemchukwu Q, Obeagu EI, Ochei KC. Prevalence of Anaemia among Pregnant Women in Braithwaite Memorial Specialist Hospital (BMSH) Port Harcourt. *IOSR Journal of Pharmacy and Biological Sciences*. 2014;9(5):59-64.
74. Natama HM, Moncunill G, Rovira-Vallbona E, Sanz H, Sorgho H, Aguilar R, Coulibaly-Traoré M, Somé MA, Scott S, Valéa I, Mens PF. Modulation of innate immune responses at birth by prenatal malaria exposure and association with malaria risk during the first year of life. *BMC medicine*. 2018; 16:1-5.

75. Akandinda M, Obeagu EI, Katonera MT. Non Governmental Organizations and Women's Health Empowerment in Uganda: A Review. *Asian Research Journal of Gynaecology and Obstetrics*. 2022;8(3):12-26.
76. Gamde MS, Obeagu EI. IRON DEFICIENCY ANAEMIA: ENEMICAL TO PREGNANCY. *European Journal of Biomedical*. 2023;10(9):272-275. links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf.
77. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
78. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Studies of Some Haemostatic Variables in Preeclamptic Women in Owerri, Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):39-48.
79. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8):22-26.
80. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023;6(2):10-13.
81. Mutabingwa TK, Bolla MC, Li JL, Domingo GJ, Li X, Fried M, Duffy PE. Maternal malaria and gravidity interact to modify infant susceptibility to malaria. *PLoS medicine*. 2005;2(12):e407.
82. Gamble C, Ekwaru PJ, Garner P, Ter Kuile FO. Insecticide-treated nets for the prevention of malaria in pregnancy: a systematic review of randomised controlled trials. *PLoS medicine*. 2007;4(3):e107.
83. Okoko BJ, Enwere G, Ota MO. The epidemiology and consequences of maternal malaria: a review of immunological basis. *Acta tropica*. 2003;87(2):193-205.
84. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
85. Umbers AJ, Stanisic DI, Ome M, Wangnapi R, Hanieh S, Unger HW, Robinson LJ, Lufele E, Baiwog F, Siba PM, King CL. Does malaria affect placental development? Evidence from in vitro models. *PLoS One*. 2013;8(1):e55269.
86. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
87. Gbedande K, Carpio VH, Stephens R. Using two phases of the CD 4 T cell response to blood-stage murine malaria to understand regulation of systemic immunity and placental pathology in Plasmodium falciparum infection. *Immunological reviews*. 2020;293(1):88-114.
88. Lindsay SW, Thomas MB, Kleinschmidt I. Threats to the effectiveness of insecticide-treated bednets for malaria control: thinking beyond insecticide resistance. *The Lancet Global Health*. 2021;9(9):e1325-1331.

89. Akinleye SO, Falade CO, Ajayi IO. Knowledge and utilization of intermittent preventive treatment for malaria among pregnant women attending antenatal clinics in primary health care centers in rural southwest, Nigeria: a cross-sectional study. BMC pregnancy and childbirth. 2009; 9:1-9.