

## Anemia and Malaria: Twin Troubles in Pediatric Health

\*Emmanuel Ifeanyi Obeagu<sup>1</sup> and Getrude Uzoma Obeagu<sup>2</sup>

<sup>1</sup>Department of Medical Laboratory Science, Kampala International University, Uganda

<sup>3</sup>School of Nursing Science, Kampala International University, Uganda

\*Corresponding authour: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, emmanuelobeagu@yahoo.com, ORCID: 0000-0002-4538-0161](#)

### Abstract

Anemia and malaria represent twin troubles in pediatric health, particularly in regions where malaria transmission is endemic. These two conditions frequently coexist, exacerbating each other's impact and imposing a significant burden on children's health and well-being. This review explores the complex interplay between anemia and malaria in pediatric populations, focusing on the epidemiology, pathophysiology, diagnostic challenges, and management strategies. It examines the bidirectional relationship between these conditions, highlighting how malaria-induced hemolysis contributes to anemia while anemia predisposes children to severe malaria and its complications. The review also discusses the implications for public health interventions, emphasizing the importance of integrated approaches that address both malaria prevention/control and anemia management. By elucidating the intertwined nature of anemia and malaria in pediatric health, this review aims to inform healthcare providers, researchers, and policymakers about the challenges and opportunities for improving the health outcomes of children in malaria-endemic regions.

**Keywords:** *Anemia, Malaria, Pediatrics, Hemolytic Anemia, Plasmodium falciparum, Iron Deficiency, Children, Public Health, Co-infection, Management Strategies*

### Introduction

Anemia and malaria stand as twin troubles in pediatric health, particularly in regions where the disease burden is high. Anemia, characterized by a deficiency in red blood cells or hemoglobin, affects millions of children worldwide, leading to significant morbidity and mortality. Malaria, a mosquito-borne infectious disease caused by Plasmodium parasites, presents a pervasive threat to

**Citation:** Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. *Elite Journal of Health Science*, 2024; 2(3): 36-43

children, particularly in sub-Saharan Africa, Southeast Asia, and parts of South America. The co-occurrence of anemia and malaria exacerbates the health challenges faced by pediatric populations, creating a vicious cycle of illness, malnutrition, and impaired cognitive development. The relationship between anemia and malaria is complex and multifaceted. Malaria contributes significantly to the burden of anemia, as the infection directly destroys red blood cells and disrupts the body's ability to produce new ones. Moreover, the inflammatory response triggered by malaria infection can interfere with iron metabolism and erythropoiesis, further exacerbating anemia. Conversely, anemia increases the risk and severity of malaria infection, as reduced hemoglobin levels compromise the body's ability to deliver oxygen to tissues, weakening the immune response and increasing susceptibility to severe complications of malaria.<sup>1-14</sup>

Pediatric populations are particularly vulnerable to the dual burden of anemia and malaria due to their heightened nutritional needs, immature immune systems, and limited access to healthcare resources in many endemic regions. The consequences of these twin troubles extend beyond physical health, impacting children's cognitive development, school performance, and long-term productivity. Addressing anemia and malaria in pediatric populations requires a comprehensive approach that encompasses preventive measures, early diagnosis, prompt treatment, and efforts to address underlying socio-economic determinants of health. Despite significant progress in recent years, anemia and malaria remain formidable challenges in pediatric health, especially in resource-limited settings. Sustainable solutions necessitate interdisciplinary collaboration among healthcare providers, researchers, policymakers, and community stakeholders to implement evidence-based interventions, strengthen healthcare systems, and address the root causes of these twin troubles. By prioritizing pediatric health and investing in holistic strategies, we can mitigate the impact of anemia and malaria, improving the well-being and future prospects of millions of children worldwide.<sup>15-22</sup>

## **Epidemiology and Pathophysiology**

Anemia and malaria both pose significant public health challenges, particularly in regions with high disease prevalence and limited access to healthcare. In endemic areas such as sub-Saharan Africa, Southeast Asia, and parts of South America, these twin troubles disproportionately affect pediatric populations. Malaria, caused by *Plasmodium* parasites transmitted through the bites of infected mosquitoes, accounts for hundreds of thousands of deaths annually, with children under five years old bearing the greatest burden. Anemia, characterized by low levels of hemoglobin or red blood cells, affects approximately a third of the world's population, with children and pregnant women being the most vulnerable groups. The epidemiology of anemia and malaria is closely intertwined, with each condition exacerbating the other's impact. Malaria infection contributes significantly to the global burden of anemia, as the destruction of red blood cells by the parasite leads to hemolysis and subsequent decreases in hemoglobin levels. Additionally, chronic inflammation associated with malaria can disrupt iron metabolism and impair erythropoiesis, further worsening anemia. Conversely, anemia increases the risk and severity of malaria infection, as reduced hemoglobin levels compromise oxygen delivery to tissues, weakening the immune response and increasing susceptibility to severe malaria complications.<sup>23-32</sup>

**Citation:** Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. *Elite Journal of Health Science*, 2024; 2(3): 36-43

## Pathophysiology

The pathophysiology of anemia and malaria involves intricate interactions between host factors, parasite biology, and environmental influences. In malaria, the *Plasmodium* parasite undergoes a complex life cycle within the human host, invading red blood cells and replicating inside them before bursting out and infecting new cells. This process leads to the destruction of red blood cells, causing hemolysis and the release of malarial toxins and byproducts into the bloodstream. The resulting inflammatory response triggers the activation of immune cells and the release of cytokines, contributing to tissue damage and systemic symptoms. Anemia in the context of malaria can arise through multiple mechanisms, including direct hemolysis of infected red blood cells, dysregulated immune responses, and suppression of erythropoiesis. The parasite's ability to sequester within deep tissues and evade immune detection further complicates the host's ability to control infection and mitigate the associated anemia. Additionally, factors such as nutritional deficiencies, concurrent infections, and genetic predispositions can exacerbate both anemia and malaria, creating a vicious cycle of illness and malnutrition in pediatric populations.<sup>33-46</sup>

## Clinical Manifestations and Complications

The clinical manifestations of anemia and malaria in pediatric populations can vary widely depending on factors such as age, nutritional status, parasite species, and disease severity. In the context of anemia, children may present with symptoms such as fatigue, weakness, pallor, shortness of breath, and exercise intolerance. Severe cases of anemia can lead to tachycardia, palpitations, dizziness, fainting, and angina. In infants and young children, anemia may manifest as failure to thrive, developmental delays, and cognitive impairments. The severity of symptoms correlates with the degree of anemia, with profound anemia posing the greatest risk of morbidity and mortality. Malaria presents with a spectrum of clinical manifestations, ranging from asymptomatic infection to severe and life-threatening disease. The typical symptoms of uncomplicated malaria include fever, chills, headache, myalgia, and malaise. Children with malaria may also experience nausea, vomiting, abdominal pain, and diarrhea. In severe cases, malaria can progress rapidly, leading to complications such as cerebral malaria, severe anemia, respiratory distress, renal failure, and circulatory collapse. Neurological complications of cerebral malaria include seizures, altered consciousness, coma, and focal neurological deficits.<sup>42-45</sup>

## Complications

Anemia and malaria can give rise to a range of complications, particularly when they coexist or occur in vulnerable populations such as young children. Severe anemia resulting from malaria can exacerbate tissue hypoxia, leading to multi-organ dysfunction and metabolic acidosis. Children with severe anemia may require blood transfusions to restore oxygen-carrying capacity and prevent end-organ damage. Chronic anemia can impair growth and development, compromise immune function, and increase susceptibility to other infections. Malaria complications can be life-threatening, especially in cases of severe and untreated disease. Cerebral malaria, characterized by altered mental status, seizures, and coma, is a major cause of mortality and neurological morbidity in pediatric malaria cases. Other complications of severe malaria include acute respiratory distress

**Citation:** Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. *Elite Journal of Health Science*, 2024; 2(3): 36-43

syndrome (ARDS), acute kidney injury, liver dysfunction, disseminated intravascular coagulation (DIC), and shock. Prompt recognition and management of severe malaria are essential to prevent irreversible organ damage and improve outcomes. In endemic regions, the burden of anemia and malaria is compounded by factors such as limited access to healthcare, inadequate diagnostic facilities, and drug resistance. Integrated approaches that address both conditions simultaneously, such as intermittent preventive treatment in pregnancy (IPTp), insecticide-treated bed nets (ITNs), and antenatal iron supplementation, have shown promise in reducing morbidity and mortality in pediatric populations. Additionally, research efforts focused on developing new diagnostic tools, treatments, and vaccines are crucial for advancing our understanding and control of these twin troubles in pediatric health.<sup>41-46</sup>

## **Management Strategies**

Management strategies for addressing the twin troubles of anemia and malaria in pediatric populations require a multifaceted approach encompassing prevention, diagnosis, treatment, and supportive care. Here are some key strategies:

### **Prevention**

Vector control measures, such as insecticide-treated bed nets (ITNs) and indoor residual spraying (IRS), to reduce malaria transmission. Chemoprevention strategies, including intermittent preventive treatment in pregnancy (IPTp) and seasonal malaria chemoprevention (SMC) in children, to prevent malaria infection. Promotion of exclusive breastfeeding, adequate nutrition, and micronutrient supplementation to prevent nutritional anemia. Improved access to clean water, sanitation, and hygiene (WASH) facilities to reduce the risk of waterborne infections and intestinal parasites that can contribute to anemia.<sup>11</sup>

### **Early Diagnosis**

Use of rapid diagnostic tests (RDTs) to promptly diagnose malaria and differentiate between Plasmodium species. Screening for anemia through simple and cost-effective methods such as hemoglobin measurement using portable devices or laboratory tests. Integration of screening for anemia and malaria into routine pediatric healthcare visits, antenatal care, and community health programs.<sup>14</sup>

### **Treatment**

Antimalarial therapy with effective medications such as artemisinin-based combination therapies (ACTs) for uncomplicated malaria. Management of severe malaria with intravenous or intramuscular antimalarial drugs, such as artesunate, in hospital settings. Treatment of anemia with oral or parenteral iron supplementation, folate, vitamin B12, and other micronutrients as indicated. Blood transfusion for children with severe anemia or signs of impaired tissue perfusion.<sup>34</sup>

### **Supportive Care**

**Citation:** Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. *Elite Journal of Health Science*, 2024; 2(3): 36-43

Fluid resuscitation and electrolyte management for children with severe malaria or complications such as dehydration and metabolic acidosis. Monitoring and management of complications, including cerebral malaria, acute respiratory distress syndrome (ARDS), renal failure, and shock. Nutritional support, including therapeutic feeding programs and ready-to-use therapeutic foods (RUTF), for children with malnutrition or nutritional deficiencies contributing to anemia.<sup>45</sup>

### **Health Systems Strengthening**

Training healthcare providers in the diagnosis, treatment, and management of anemia and malaria, including recognition of severe disease and referral protocols. Ensuring reliable access to essential medications, diagnostic tools, and blood transfusion services. Community engagement and education on preventive measures, early recognition of symptoms, and prompt seeking of healthcare services. Integration of anemia and malaria control efforts with existing maternal and child health programs, immunization campaigns, and school-based health initiatives.<sup>40</sup>

### **Conclusion**

The coexistence of anemia and malaria presents significant challenges to pediatric health, particularly in regions where both diseases are endemic. Anemia, characterized by low levels of hemoglobin or red blood cells, and malaria, caused by Plasmodium parasites transmitted through mosquito bites, contribute to morbidity and mortality among children worldwide. The intertwined epidemiology, pathophysiology, clinical manifestations, and complications of these twin troubles underscore the importance of integrated management strategies that address both conditions simultaneously. Preventive measures such as vector control, chemoprevention, nutritional interventions, and improved access to clean water and sanitation are essential for reducing the incidence of anemia and malaria in pediatric populations. Early diagnosis through rapid diagnostic tests and screening for anemia can facilitate prompt initiation of appropriate treatment. Antimalarial therapy, iron supplementation, supportive care, and health systems strengthening are key components of effective management strategies aimed at reducing the burden of these diseases and mitigating their impact on child health.

### **References**

1. Verhoef H. Iron deficiency and malaria as determinants of anaemia in African children. Wageningen University and Research; 2001.
2. Namaste S. The Burden of Anemia, Iron Deficiency and Vitamin A Deficiency in Young Children Living in Kenya and Malawi. The George Washington University; 2014.
3. Gupta A, Gupta A. Aetiology of Iron Deficiency in Children. Nutritional Anemia in Preschool Children. 2017;47-118.
4. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):1-9.

5. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfectd with HIV in Umuahia. *Int. J. Curr. Res. Med. Sci.* 2017;3(5):100-104.
6. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. *Medicine.* 2023;102(46): e36166.
7. Obeagu EI, Chijioke UO, Ekelozie IS. Malaria rapid diagnostic test (RDTs). *Ann Clin Lab Res.* 2018;6(4):275.
8. Obeagu EI, Obeagu GU, Egba SI, Emeka-Obi OR. Combatting Anemia in Pediatric Malaria: Effective Management Strategies. *Int. J. Curr. Res. Med. Sci.* 2023;9(11):1-7.
9. Hassan AO, Oso OV, Obeagu EI, Adeyemo AT. Malaria Vaccine: Prospects and Challenges. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(2):22-40.
10. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(4):10-9.
11. Obeagu EI, Busari AI, Uduchi IO, Ogomaka IA, Ibekwe AM, Vincent CC, Chijioke UO, Okafor CJ, Okoroiwu HU, Adike CN. Age-Related Haematological Variations in Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(42B):218-24.
12. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(41B):380-386.
13. Ogbonna CO, Obeagu EI, Ufelle SA, Ogbonna LN. Evaluation of haematological alterations in children infected by *Plasmodium falciparum* Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(1):38-45.
14. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research.* 2019;2(1):25-36.
15. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9):1-4.
16. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with *Plasmodium falciparum* malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2016;3(8):40-6.
17. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research.* 2017;5(4):206.
18. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Irondi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal.* 2022;6(3):9-17.



19. Obeagu EI, Nimo OM, Bunu UO, Ugwu OP, Alum EU. Anaemia in children under five years: African perspectives. *Int. J. Curr. Res. Biol. Med.* 2023; 1:1-7.
20. Madekwe CC, Madekwe CC, Obeagu EI. Inequality of monitoring in Human Immunodeficiency Virus, Tuberculosis and Malaria: A Review. *Madonna University journal of Medicine and Health Sciences.* 2022;2(3):6-15.
21. Offie DC, Ibekwe AM, Agu CC, Esimai BN, Okpala PU, Obeagu EI, Ufelle SA, Ogbonna LN. Fibrinogen and C-Reactive Protein Significance in Children Infected by *Plasmodium falciparum* Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(15):1-8.
22. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences.* 2018;4(9):41-9.
23. Eberendu IF, Ozims SJ, Agu GC, Amah HC, Obasi CC, Obioma-Elemba JE, Ihekaire DE, Ibanga IE, Amah CC, Obeagu EI, Nwosu DC. Impact of human activities on the breeding of mosquitoes of human disease in Owerri metropolis, Imo state. *Int J Adv Res Biol Sci IJARBS.* 2017;4(12):98-106.
24. Obeagu EI, Ofodile AC, Okwuanaso CB. A review on socio economic and behavioral aspects of malaria and its control among children under 5 years of age in Africa. *J Pub Health Nutri.* 2023; 6 (1): 136.
25. Obeagu EI, Obeagu GU. Maternal Malaria and Neonatal Developmental Outcomes: A Life course Perspective. *Elite Journal of Health Science.* 2024;2(1):43-50.
26. Obeagu EI, Obeagu GU. Understanding Hematocrit Fluctuations in HIV-Malaria Coinfection for Improved Management. *Elite Journal of Public Health.* 2024;2(1):22-34.
27. Obeagu EI, Obeagu GU. Overcoming Hurdles: Anemia Management in Malaria-Affected Childhood. *Elite Journal of Laboratory Medicine.* 2024;2(1):59-69.
28. Obeagu EI, Ogunnaya FU, Obeagu GU. Integrated Approaches for Improving Pediatric Health: Addressing Anemia in Malaria Cases. *Journal home page: <http://www.journalijiar.com>;*12(01).
29. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. *Elite Journal of Laboratory Medicine.* 2024;2(1):33-45.
30. Obeagu EI, Obeagu GU. Hematocrit Fluctuations in HIV Patients Co-infected with Malaria Parasites: A Comprehensive Review. *Int. J. Curr. Res. Med. Sci.* 2024;10(1):25-36.
31. Obeagu EI, Obeagu GU. Advances in Understanding the Impact of Blood Transfusion on Anemia Resolution in HIV-Positive Children with Severe Malaria: A Comprehensive Review. *Elite Journal of Haematology.* 2024;2(1):26-41.
32. Obeagu EI, Obeagu GU. Malaria's Silent Partner: Anemia in Children. *Elite Journal of Public Health.* 2024;2(1):1-7.
33. Obeagu EI, Obeagu GU. Transfusion-Related Complications in Children Under 5 with Coexisting HIV and Severe Malaria: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2024;11(2):9-19.
34. Obeagu EI, Obeagu GU, Hauwa BA, Umar AI. Hematocrit Variations in HIV Patients Co-infected with Malaria: A Comprehensive Review. *Journal home page: <http://www.journalijiar.com>;*12(01).

**Citation:** Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. *Elite Journal of Health Science*, 2024; 2(3): 36-43

35. Obeagu EI, Obeagu GU, Hauwa BA, Umar AI. Neutrophil Dynamics: Unveiling Their Role in HIV Progression within Malaria Patients. Journal home page: <http://www.journalijar.com>.;12(01).
36. Obeagu EI, Obeagu GU. Synergistic Effects of Blood Transfusion and HIV in Children Under 5 Years with Severe Malaria: A Review. Elite Journal of HIV. 2024;2(1):31-50.
37. Kalu OA, Ukibe NR, Onah CE, Obiorah OC, Obeagu EI, Awalu CJ, Ukibe EG, Ukibe BC. Assessment of serum soluble intercellular adhesion molecule-1 and albumin in human immuno-deficiency virus-infected individuals with or without malaria parasite infection in Nauth, Nnewi, Nigeria. International Journal of Research in Medical Sciences. 2024 Feb;12(2):349.
38. Ukibe NR, Alex JC, Osakue NO, Ukibe EG, Ukibe BC, Ukibe VE, Obeagu EI. Linking Malaria and Hypertension: Unveiling the Interconnected Pathophysiological Nexus. mortality.;66:75.
39. Ajugwo AO, Bot YS, Obeagu EI, Mugabiza M. Anaemia and Plasmodiasis among children below five years attending OPD at Ishaka Adventist Hospital, Bushenyi District. Journal of Haematology. 2024;2(2):34-45.
40. Obeagu EI, Obeagu GU. Maternal Eosinophilic Responses in HIV-Positive Pregnant Women: Unraveling Immunological Dynamics for Improved Maternal-Fetal Health. Elite Journal of Immunology. 2024;2(1):47-64.
41. Obeagu EI, Obeagu GU. Implications of B Lymphocyte Dysfunction in HIV/AIDS. Elite Journal of Immunology. 2024;2(1):34-46.
42. Obeagu EI, Obeagu GU, Okwuanaso CB. Optimizing Immune Health in HIV Patients through Nutrition: A Review. Elite Journal of Immunology. 2024;2(1):14-33.
43. Obeagu EI, Obeagu GU. The Crucial Involvement of CD8 in HIV Progression: A Review. Int. J. Curr. Res. Med. Sci. 2024;10(2):15-25.
44. Obeagu EI, Obeagu GU. Counting Cells, Shaping Fates: CD4/CD8 Ratios in HIV. Elite Journal of Scientific Research and Review. 2024;2(1):37-50.
45. Obeagu EI, Obeagu GU. Platelet Distribution Width (PDW) as a Prognostic Marker for Anemia Severity in HIV Patients: A Comprehensive Review. Journal home page: <http://www.journalijar.com>.;12(01).
46. Obeagu EI, Obeagu GU. Eosinophil Dynamics in Pregnancy among Women Living with HIV: A Comprehensive Review. Int. J. Curr. Res. Med. Sci. 2024;10(1):11-24.