

Adverse Effects of Antimicrobial Therapeutic Agents in Common Use: A Review

S.O. Onemu^{1*}, M.O. Onemu-Metitiri², O. Odeyemi¹, P.O. Uyigue¹ and Emmanuel Ifeanyi Obeagu³

1. Department of Medical Laboratory Science, Achievers University, Owo, Nigeria; onemuso@achievers.edu.ng
2. School of Sciences, Engineering and Environment, University of Salford, United Kingdom. tefemetitiri@gmail.com
3. Department of Medical Laboratory Science, Kampala International University, Uganda

*Corresponding author: SO Onemu, Department of Medical Laboratory Science, Achievers University, Owo, Nigeria; onemuso@achievers.edu.ng

Abstract

Antimicrobial agents play critical roles in reducing the burden of bacterial infectious diseases and their associated fatalities but may also come with unintended consequences that in some cases may lead to other health burdens, disability and death in extreme cases. Unwanted effects occur commonly in the administration of a wide variety of drugs, but antimicrobial agents garner less attention as some of these agents pose extensive diversities of objectionable side effects. This review focuses on highlighting the dangers that may accompany the administration of antimicrobial agents. Literature search applying keys words on the subject matter was conducted and related materials were assembled. Adverse effects of antimicrobial agents ranged from hypersensitivity to penicillins, bleeding disorder with members of the fourth generation cephalosporins, nephrotoxicity and ototoxicity common to gentamycin, tendon rupture, liver toxicity and induction of hypoglycemic state with some fluoroquinolones. The interference and constitutional alteration in intestinal bacterial community and consequential overgrowth of resistant *Clostridioides difficile* is prominent in the administration some antimicrobial agents and especially with the clindamycins. The attendant side effects of each antimicrobial agent demand re-evaluation, expediency of personalized prescription and proper patient education to minimize and circumvent the adverse drug effects of antimicrobial agents.

Keywords: Adverse drug effects, antimicrobial agents, disease burden, morbidity, microbiome alteration, patient education, personalized prescription

1. Introduction.

Antimicrobial agents play a key role in preventing millions of deaths worldwide since their introduction into medical care in the 1940s. Antimicrobial agents also possess adverse drug effects or reactions, ADEs/ADRs that can immensely affect the outcomes of their administration [1,2]. Gerhard Domagk – a German Chemist initiated the first true antimicrobial therapy with prontosil (sulfonamide compound) in 1935 to treat different infections including syphilis but the patients experienced severe ADEs that included nausea and neurological disturbances [2]. Prontosil

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

became the precursor of modern sulfonamides [3] Alexander Flemming researching independently at the time of prontosil's introduction had previously observed the activity of penicillin against bacteria in 1928. Each of these early researchers were spurred into action by the horrific wound infections that were common place during the World War I, culminated in discovery of penicillin, eventual purification and availability for treatment of major human infections, marking a major turning point in treatment of bacterial infections [4]. The treatment of several bacterial illnesses that included pneumonia, septicemia, diphtheria, tetanus, typhoid and tuberculosis, all with elevated fatality rates became possible [5]. Following the introduction of penicillin into clinical practice there was a rapid discovery of other antimicrobial agents. This review highlights the adverse effects of common groups of antimicrobial agents regularly available for the management of bacterial infections.

2. Methodology

The execution of this review was with the aid of search engines applying the key words – antimicrobial agents, antibiotics, antibacterial agents, adverse effects/reactions/events, side effects combined with the different groups of antimicrobial agents. Literature that contained related information to the major review themes that included penicillins, cephalosporins, aminoglycosides, quinolones, macrolides, sulfonamides, tetracyclines, chloramphenicol, lincosamides, carbapenems, glycopeptides, monobactams oxazolidinones, rifampicin ethambutol and isoniazid in English language were extracted for compilation.

3. Antimicrobial agents and ADEs

3.1. Penicillins.

The availability of penicillin for the treatment of a wide variety bacterial infection in 1945 marked a great leap forward in human development and emancipation from the era of fatal infectious diseases [6]. Penicillin act by inhibiting the transpeptidation steps in bacterial cell intermediate macromolecules that are crucial for peptidoglycan formation [7, 8]. Extensive excretion of penicillins happens in the renal system and this can result in prolonged biological half-lives and elevated serum levels that can exacerbate the tendency for toxicity [9]. The kidneys are able to eliminate virtually all penicillins without the need for dosage adjustment to circumvent hepatotoxicity. The most prominent side effect of the penicillins is the propensity to bind to serum proteins forming hapten-protein complex to become antigenic as the basis of the hypersensitivity reactions commonly seen in piperacillin administration that may lead to hemolytic anemia if the drug binds to red blood cells [10, 11, 12]. This can occur at any age, but notably between the ages of 20 and 49 years, there is an increased risk for anaphylactic reaction during a second period in about 15% of patients administered with a penicillin agent [13]. The commonest adverse events in treatment with penicillins are usually intestinal discomfort involving nausea, vomiting, epigastric pain and antibiotic induced *Clostridioides difficile*. Hepatic effects and thrombophlebitis may spike transaminases with high doses of nafcillin in patients with renal limitations [14].

3.2. Cephalosporins

These agents have structural similarities with the penicillins by also possessing the beta-lactam ring and well tolerated like most penicillins and effective substitutes. They inhibit bacteria cell wall enzyme by binding to penicillin binding proteins, PBPs [15]. The ADEs of cephalosporins are similar to those of the penicillins with the exception of occasional biliary effect, bilirubinemia and a very rare fatal hemolysis observed with ceftriaxone [16]. Cephalosporins have low toxicity

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

without serious adverse events even in patients with evidence of previous hypersensitivity to penicillins [17]. The toxicity of cephalosporins are restricted to minor differences in the chemical and structural constitution of modern cephalosporins molecules that potentiate undesired reactions in up to 3.2% of patients [18]. The general reactions reported in cephalosporins include fever, arthralgia and exanthema in groups of children administered with cefaclor [16] and hypersensitivity in patients with liver disease and asthma [19,20]. Nephrotoxicity is uncommon in modern cephalosporins except for significant decline in renal activity with large regimens of ceftazidime, ceftriaxone and cefoperazone that each has a fraction of 3-methyl-thiotetrazole side-chain that interferes in the biosynthesis of prothrombin with consequential increase in the tendency for bleeding disorder with concomitant increase in disulfiram-like effects in individuals concurrently taking alcohol containing substances [21]. Cephalosporins ADEs are commoner in children aged four years and spikes with the age of the child especially with the use of ceftazidime, ceftriaxone and cefoperazone are sometimes fatal [22]. Neurological and psychological disturbances including epidermal and arthritic effects have been reported to be associated with the use of cefaclor [23, 24].

3.3. Aminoglycosides.

These agents are some of the foremost and very effective antibacterial agents that have continued to retain a key position in the treatment of serious Gram-negative bacterial infections including urinary tract infections and bacteremia [25]. Aminoglycosides inhibit 30S bacterial ribosomes thereby preventing the synthesis of essential proteins [26]. The glomeruli rapidly filter the agents unchanged within two hours. This can however be extended to 30-60 hours in patients with impaired renal function thereby requiring dosage modification [27]. Some of the most commonly used aminoglycosides include gentamycin, amikacin, tobramycin, streptomycin, neomycin and kanamycin. The initial application of streptomycin was for the treatment of tuberculosis while neomycin and kanamycin are effective in gut sterilization and wound infections on account their toxicity. Aminoglycosides are polar agents that are poorly absorbed through oral route of administration necessitating the parenteral route. The most pronounced ADEs of aminoglycosides are nephrotoxicity and ototoxicity. The nephrotoxicity is worrisome as it affects creatinine clearance causing a build-up of concentrations that can induce nephrotoxicity and the possibility of neurotoxicity that affects the kidneys due to the agents binding to phospholipid in the liposomes hindering permeability and leading to accumulation [27], and inhibition of their function [26]. Exposure to aminoglycosides in-utero can result in permanent deafness [28], whereas neurotoxicity can be reversible [25]. The mechanism of neurotoxicity is through the drugs ability to freely pass into hair cell and induce reactive oxygen species, ROS to damage the mitochondria resulting in cell death. Aminoglycosides nephrotoxicity is bothersome on the account of their significant effect on the proximal tubule epithelial cells arising from selective endocytosis and build-up of aminoglycosides in the renal cortex that can activate acute kidney injury [29]. Kidney injury initiated as the agents filter unchanged in the glomeruli, and reabsorbed by the proximal epithelia where they concentrate into liposomes after encounter with the phospholipids on the brush border membranes thereby interfering with phospholipid metabolism. This in addition, affects the functions of the cells of the kidneys greatly, influenced by the dosage and length of administration [27]. The pressure to use an aminoglycoside has dissipated immensely with the availability of less toxic and effective agents for treating Gram-negative bacterial infections [30].

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

3.4. Fluoroquinolones.

The arrival of the fluoroquinolones into the clinical arena with their wide spectrum of activity displaced some other frequently administered agents like the aminoglycosides administered parenterally with notable adverse effects. The fluoroquinolones combine the advantage of being available for oral use and outstanding bioavailability [31]. The fluoroquinolones inhibit bacterial gyrase and topoisomerases II and IV [32, 33]. The fluoroquinolones are associated with arthralgia in immature animals, which formed the basis of their restriction during pregnancy, and in the young [34]. These agents exhibit a range of intense and permanent forms of ADEs that include tendon atrophy, tendon rupture, and induction of hypoglycemic states, peripheral neurological toxicity, retinal detachment and aneurism of the aorta [35,36]. Teratogenic effects have with ciprofloxacin in mice even at low doses has been reported [37], and significant declines in spermatozoa concentration of infertile men treated with ciprofloxacin [38]. In 2008, the Food and Drug Administration, FDA issued a warning concerning the side effects of fluoroquinolones. This was followed by the European Medicine Agency, EMA and other countries' regulatory authorities issuing similar warnings on the ADEs involved in the use of some fluoroquinolones [39]. Gatifloxacin was a major quinolone that fell under the hammer due to its marked depression of glucose utilization leading to hypoglycemic state [40, 41]. Trovafloxacin is associated with induction of severe liver injury and fatal outcome [42]. In consequence, the two agents fell into disuse for oral or systemic administration. Studies have indicated that ciprofloxacin initiates DNA damage, chromatin abnormalities of sperm cells that could contribute to the observed low fertilization rates and retarded embryo development [43]. The risk of tendonopathy, peripheral neuropathy, retinal detachment, cardiac arrhythmia and central nervous system involvement are higher in individuals who are 60 years or older [36,39].

3.5. Sulfonamides.

These agents are in use in a wide array of human and animal situations that include the treatment of toxoplasmosis when used in combination with antimalarial agents [44]. Sulfonamides have a wide spectrum of activity and the foremost agents applied for the treatment of bacterial infections [45]. Some of the earlier sulfonamides (prontosil) are associated with grave ADEs. Sulfonamides with better tolerability have since been developed that can be administered for specific indications. Sulfonamides competitively inhibit the synthesis of protein with adverse reactions of modern sulfonamides including lethal toxidermia, severe liver injury, pulmonary reactions and blood dyscrasias [46]. The mechanism of the side effects remains unclear, it is presumed that the metabolites of sulfamethoxazole of the most regularly prescribed sulfonamide bind covalently to protein due to the chemical reactivity leading to the formation of unique adverse events [47]. Immune responses that follow could also trigger hematological complications antecedent to hemolytic anemia, agranulocytosis and or suppression of hematopoietic activity of the pluripotent stem cell in the bone marrow [48]. Renal involvement usually manifests from the crystallization of sulfonamide by-products in the renal tubules to precipitate kidney stones exhibiting as interstitial nephritis and or acute kidney injury, AKI [49,50].

3.6. Tetracyclines

The tetracyclines discovered in the 1940s possess activity against a wide spectrum of microorganisms including Gram-positive and Gram-negative bacteria, chlamydiae, mycoplasma, rickettsiae and protozoa [51]. They are inexpensive and used extensively as prophylactic agents in

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

human treatment and incorporated into livestock feeds at the sub-therapeutic levels as growth promoter thereby creating a high pressure for the emergence of resistance to these agents, this has severely limited their use in clinical settings [52]. Tetracyclines interfere with the 30S ribosomal subunit binding to the amino-acyl-tRNA to the receptor site on the mRNA-ribosome complex thereby hindering the process. Photosensitivity is a well-documented major ADE of the tetracyclines that involves the development of a complex with calcium ions capable of absorbing ultra-violet, UV light. This complex then generates reactive oxygen species, ROS that can damage DNA and cell membrane on exposure to sunlight leading to dermatological complications, initiation of lupus erythematosus, LE and pressure within cranium [53,54].

3.7. Macrolides

These agents have a spectrum of activity that includes viruses, fungi and protozoa [54] and recommended in the management of respiratory infections with anti-inflammatory activity [55-57]. Macrolides typically impair the synthesis of protein by attaching to 23S of the 50S ribosomal unit of bacteria [58]. Macrolides are very safe agents, with side effects usually limited to minor intestinal disturbances occurring in 15-20% of individuals taking erythromycin and less than 5% with the newer macrolides that rarely initiate motilin release *in-vivo*, such as clarithromycin, azithromycin, dirithromycin and rikamycin except for high doses of troleandomycin and erythromycin [59].

3.8. Lincosamides

This group consists of two main antibiotics that are used in human medicine and veterinary settings. Lincomycin and its derivative – clindamycin are agents with inhibitory effect on both Gram-positive and anaerobic Gram-negative bacteria as well as on protozoa especially in combination those of malaria [60]. Lincosamides hinder 23S on the RNA bigger subunit, thereby cutting off the path by which emerging intermediates substrates needed for the next step in protein synthesis [61-63]. The major side effects is the ability to create pseudomembranous colitis, nausea and abdominal cramps or vomiting following administration. The significant interference and modification of resident intestinal bacteria community (microbiome) by clindamycin confer selective advantage on resistant *Clostridioides difficile* to overgrow and produce toxins that lead to the development of micro-abscesses on the intestinal mucosae that coalesce into pseudo-membranes and may necessitate the discontinuation of the agent or intervention with treatment with vancomycin in extreme cases [64].

3.9. Chloramphenicol

Chloramphenicol is lowly priced antibiotic that binds to bacterial 50S subunit of the 70S ribosome thus interfering with the peptidyl-transferase chain of the nascent protein molecules [65]. The agent is particularly effective in treating cases of meningitis due to *Neisseria meningitides*, *Streptococcus pneumoniae* and *Haemophilus influenza* - the main bacterial agents of meningitis [66]. The key concern in prescribing chloramphenicol lies in its neurotoxicity and irreversible hematopoietic damage to pluripotent stem cells of the bone during extended period of use [66]. Chloramphenicol is still useful in life threatening infections especially in resource-poor countries and with the rising spate of multidrug resistant, MDR bacteria worldwide, there is a reawakening of attention on old antibiotics and intense focus on chloramphenicol as candidate for many infections [67-71].

3.10. Carbapenems

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

These are beta-lactam agents as the penicillins and cephalosporins representing the most efficacious agents in infections involving resistant bacterial strains as well as demonstrating a unique level of destructive effects on beta-lactamases [72-74]. The deployment of the agents is often as the antibiotics of last resort in the management of refractory Gram-negative bacterial infections [75]. The carbapenems act by blocking the synthesis of cell wall as the penicillins [73]. The general ADEs of this group of agents involve abdominal pain, nausea and vomiting in about 20% of patients with swelling and pain at the injection site. There are reports of nephrotoxicity about imipenem when not administered with cilastatin to shield the drug from renal dihydropeptidase degradation [73].

3.11. Glycopeptides

This are naturally occurring and semisynthetic agents with wide spectra of action, which are especially effective against Gram-positive and methicillin-resistant *Staphylococcus aureus*, MRSA [76-80]. The spiral of MDR bacteria in clinical settings all over the world has placed immense reliance on these agents as the drugs of last resort [76,80]. The ADEs associated are also partly responsible their restricted use [81]. The major members include vancomycin, teicoplanin, oritavancin and dalbavancin, all of which act by inhibiting cell wall synthesis [78,82-84]. The prominent ADEs experienced include infusion reaction or red man's syndrome, nausea, vomiting, diarrhea, rash pruritis and a rare hypersensitivity and *Clostridioides difficile* when given oral route [78].

3.12. Monobactams

Aztreonam is the main member of this group possessing a narrow range of activity with activity essentially against Gram-negative bacteria and remarkably effective in treating MDR bacterial infections with stability in the presence of metallo-beta-lactamases, MBLs except for the variant serine-metallo-beta-lactamases, SBLs that often occur concurrently with MBLs [85]. Aztreonam act by preventing the assembly peptidoglycan intermediates resulting in failure of cell wall synthesis [86]. High doses are associated with neutropenia in 11.3% of young patients, elevated transaminases and there are no reports of nephrotoxicity, ototoxicity or hematopoietic disorders [87].

3.13. Oxazolidinones

Linezolid is the first member of this group of synthetic antimicrobial agents introduced into human medical practice [88-91]. Linezolid acts by blocking both 30S and 50S ribosomes, halting the initiation of complexes during protein synthesis [90,91]. Linezolid is especially useful in vancomycin resistant *Enterococcus*, VRE and methicillin resistant *Staphylococcus aureus*, MRSA and in hospital acquired infections, HAIs, in addition to treating multidrug resistant tuberculosis, MDR-TB [92] and has an excellent bioavailability in both oral and parenteral routes [93,94]. Linezolid is well tolerated with low toxicity compared to the higher degree of adverse events associated with vancomycin. The main ADEs of linezolid involve diarrhea nausea, vomiting and headache that are often not severe [95]. Other ADEs of linezolid involve thrombocytopenia, leukocytopenia, anemia, hypoglycemia [95] and a rare coloration of the tongue during extended periods of administration [96].

3.14. Rifampicin, Ethambutol and Isoniazid

Rifampicin is a major agent for treating a wide range of bacterial infections especially Gram-positive bacteria including *Clostridioides difficile* [97] and an effective antituberculosis agent

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

arising from its ability to bind RNA-polymerase leading to the inhibition of essential protein formation [98,99]. The concurrent use of rifampicin with other antituberculosis agents as ethambutol and isoniazid are often the main cause of ADEs that alter metabolism and regularly involve the gastric and hepatic functions [97,100]. Rifampicin rarely elicits acute side effects, while ethambutol may induce dose related retrobulbar neuritis [101]. Ethambutol and isoniazid as first line agents in tuberculosis treatment capable of initiating nephrotoxicity particularly in individuals with nephrotic syndromes arising from a decline in renal elimination [102]. Rifampicin use may lead to urticaria, flu-like manifestations, thrombocytopenia states and orange colouring of body fluids [99].

4. Antimicrobial disturbance of microbiome

The administration of antimicrobial agents is a major cause in the ecological shift in the microbial communities of the gastrointestinal tract, GIT [103-108]. The impacts of antimicrobial agents on the GIT microbial community (microbiome/microbiota) leads to the alteration in the constitution of the microbiome and optimal function of the GIT [109-113]. Recent evidence indicates that most microbiome interactions during the first year of life is critical for infant development, as early life antibiotic exposure ultimately disrupts the conventional microbiome maturation and adversely affects well-being, in addition to increasing the abundance of antibiotic resistant bacteria [103,104,107,108,110]. The disturbance of the microbiome is associated with short-term and long-term implications [112,113]. Focusing on new source of antimicrobial agents that selectively targets the infective agent will diminish the constitutional change of gut microbiome and function [112].

Table 1. Summary of some prominent ADEs of antimicrobial therapeutic agents

Agents and specific members	ADEs
Penicillins	Hypersensitivity, hemolytic anemia, anaphylaxis in 15% 20-49 years group [13], vomiting, <i>Clostridioides difficile</i> infection [10,14].
Nafcillin	Spikes in transaminases, thrombophlebitis [14]
Cephalosporins	
Cefaclor	Fever, arthralgia, exanthema in 3.2% of children in age range of 4 years [16,21]
Ceftazidime/ceftriaxone/cefoperazone	Interference with prothrombin, bleeding disorder disulfiram-like reaction [22]
Aminoglycosides	Nephrotoxicity and ototoxicity [29,30] Irreversible ototoxicity or permanent deafness in embryonic administration [28].
Fluoroquinolones	Arthropathy in immature animals, tendonopathy tendon rupture, peripheral neuropathy and aortic aneurism [35,36].
Ciprofloxacin/ofloxacin	DNA damage and chromatin abnormalities [35] Decline in spermatozoa concentration [38]. Retardation of embryo growth [43]
Gatofloxacin	Hypoglycemia and liver injury [35,40-42].
Trovafloxacin	Fatal hepatic injury [43]. Increased risk of retinal detachment, arrhythmia, neurological disorder at 60+ years [36,39].
Sulfonamides	Fatal toxidermia, acute liver injury, AKI, Pulmonary reaction and blood dyscrasias [46,47,50].
Macrolides	Intestinal disturbances in 15-20% of individuals [58]
Domycin	Motilin release in high doses [59].
Lincosamides	Nausea, vomiting, <i>Clostridioides difficile</i> infection (or pseudomembranous colitis) [64].
Chloramphenicol	Neurotoxicity, irreversible depression of hematopoiesis with prolonged use [66]
Carbapenems	Abdominal pain nausea, vomiting, neuropathy site irritation or swelling in 20% of patients [73].
Glycopeptides	Red man's syndrome, nausea vomiting, diarrhea, rare hypersensitivity, <i>Clostridioides difficile</i> [78].
Monobactams	neutropenia in 11.3% of patients, elevated transaminases, no reports of nephrotoxicity or ototoxicity or hematopoietic disorders [87]
Oxazolidinones	Nausea, vomiting, diarrhea, headache and nephrotoxicity in high doses [95]. Other ADEs

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

	of linezolid – thrombocytopenia, leukocytopenia anemia, hypoglycemia [95,96] and rare colored coated tongue [96].
Rifampicin/Ethambutol/Isoniazid	
Rifampicin	Urticaria, flu-like illness, thrombocytopenia, orange colored body fluids [97,99,100].
Ethambutol	Retrobulbar neuritis [101].
Isoniazid	Nephrotoxicity in individuals with renal Syndromes, combination of ethambutol and isoniazid alter gastric and hepatic functions [97,100].

6. Addressing ADEs of antimicrobial agents

The deployment of antimicrobial therapeutic substances classically based on intervention in many infections or for prophylaxis in human medical care and in veterinary services globally. This sometimes necessitates the need for more accountability in their administration through deliberate effort to adhere to the general guidelines for their application [114]. Significant progress recorded in the last couple of years on the logical use of antimicrobial agents aims to harness the benefits, encourage sound use and stave off the occurrence of antimicrobial selective pressure in microbial populations, and as a measure to decrease the rising spate of resistant bacterial mutants as well as reduce ADEs [114]. Adverse drug effects are responsible for a noteworthy number of deaths globally estimated to be the fourth runner in causing death in the United States [115]. There is an abysmal level of ADEs reporting on antimicrobial agents indicating that death toll may be much higher in most African health settings [116]. It is crucially important to understand the adverse reaction that could follow each agent and drug-to-drug combination as a pathway to reducing the detrimental outcome of ADEs [115-120]. The application of personalized antimicrobial prescription based on history of ADEs, patients with a history of hypersensitivity reaction reduces the potential for allergies when the agents are avoided [10]. Agents exhibiting higher frequencies of ADEs at specific age groups as the fluoroquinolones for patients who are 60 years or older. The interference with prothrombin activity and bleeding disorder is thus avoidable with this pre-knowledge and as well as in ceftazidime, ceftriaxone or cefoperazone adverse event in children in the age range of 4 years or older becomes circumventable [22, 121].

6.1. Antimicrobial agent monitoring

Agents that have the potential for to induce renal or hepatic toxicity need monitoring as in the administration of aminoglycosides and glycopeptides with the sole purpose of obtaining a better outcome of therapy and minimizing harm to the patient has been proven to be rewarding [114,122-124]. Additionally, drug monitoring is essential for agents with narrow margins for toxicity as the inhibition of the pathogen relies on the concentration in the blood or tissue that is equal to or higher than the minimum inhibitory concentration, MIC for the pathogen [122].

6.2. Patient education

Education of the patient is pivotal to proper monitoring of ADEs, prevention and intervention [125,126]. The tetracyclines, especially doxycycline may cause photosensitivity that in most

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

cases are not so severe requiring no serious course of action [127]. It is expedient that the patient receive appropriate and timely information on the possibility of sunburn and use of sunscreen. Similarly, patients administered with clindamycin need to be aware of the unintended intestinal condition associated with overgrowth of *Clostridioides difficile* and toxin production when the microbiota is disturbed. A patient with the prior knowledge of a likely ADE has a better prognosis for resolution when it occurs.

Conclusion

Antibacterial agents remain the cornerstone in the treatment of bacterial infections since introduction in the 1940s with massive decline in the hitherto mortality rates of many infectious diseases. The use of antimicrobial therapeutic agents also come with some adverse drug effects requiring clear and critical evaluation in their deployment to circumvent harm that may lead to disability or fatality. The toxicity of some fourth generation cephalosporins – ceftazidime, ceftriaxone and cefoperazone leading to bleeding disorders; initiation of nephrotoxicity and ototoxicity of aminoglycosides; hypoglycemia, tendon rupture, retinal detachment and liver toxicity associated with some members of the fluoroquinolones or development of kidney stones, fatal toxicodermia, acute liver injury of sulfonamides are well documented. Disruption and functional modification of gut microbiome with short-range or enduring consequences conjoint with many orally administered agents is prominent with the lincosamides. Maximizing the benefits of antimicrobial therapeutic agents demands personalized focus with the knowledge of interactions in the host and based on the nature of the infective agent will markedly decrease ADEs incidences in the use of antimicrobial therapeutic agents.

References

1. Jesman R, Mludzik A, Cybulska M. History of antibiotics and sulphonamides discovery. Pol Merkur Lekarski, 2011; 30 (179): 320-322.
2. Lee Y, Bradley N. Overview and insight into carbapenem allergy. Pharmacy, 2019; 7(3): 110, doi. Org/10.3390/pharmacy.7030110.
3. Ovung A, Bhattacharyya J. Sulphonamide drugs: structure, antibacterial property, toxicity and biophysical interactions. Biophys Rev, 2021; 13(2): 259-272.
4. Bentley R. Different roads to discovery of prontosil (hence sulfa drugs) and penicillin (beta-lactams). J Ind Microbiol Biotechnol, 2009; 36(6): 775-781.
5. Christensen SB. Drugs that changed society, history and current status of the early antibiotic: salvarsan, sulphonamides and beta-lactams. Molecules, 2021; 26(19): 6057, doi.10.3390/molecules26196057.
6. Lobanovski M, Pilla G. Penicillin discovery and antibiotic resistance: lessons for future. Yale J Biol Med, 2017; 135-145.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

7. Cochraine SA, Lohans T. Breaking down the cell wall: strategies for antibiotic discovery targeting bacterial transpeptidation. *Eur J Medicinal Chemistry*, 2020; 112262, doi.org/10.1016/j.ejmech.2020.112262.
8. Puls SJ, Brajtenbach D, Schneider T, Kubitscheck U, Grein F. Inhibition of peptidoglycan synthesis is sufficient for total arrest of staphylococcal division. *Science Adv*, 2023; 9(12): doi.10.1126/sciadv.ade.9023.
9. Goh SJ, Tuomisto JE, Purcell AW, Mifsud NA, Illing PT. The complexity of T cell-mediated penicillin hypersensitivity reactions. *Eur J Allergy Clin Immunol*, 2020: doi.10.1111/all.14355.
10. Pichichero ME. Penicillin and cephalosporin allergy. *Animal Allergy, Asthma, Immunol*, 2014; 112(5): 404-412.
11. Roehmel J, Specht P, Staab D, Shwarz C, Salama A, Mayer B. Risk of piperacillin hemolytic anemia in patients with cystic fibrosis and anti-pseudomonal treatment of prospective observational study. *Transfusion*, 2019; 59(12): 3746-3754.
12. Kennard L, Kayode OS, Siew LQ, Makris M, Tsilochristou O Chytiroglou E, et al. Piperacilin-tazobactam hypersensitivity: a large multicenter analysis. *J Allergy Clin Immunol In Practice*, 2021; 9(5): 2001-2009
13. Wong JC, Au EY, Yeung HH, Lau CS, Li PH. Piperacillin-tazobactam allergy: an exception to usual penicillin allergy. *Allergy Asthma Immunol Res*, 2021; 13(12): 284-294.
14. Clifford KM, Selby AR, Reveles KR, Teng C, Hall RG, McCarell J, et al. The risk and clinical implications of antibiotic associated acute kidney injury: a review of the clinical data for agents with signals from the Food and Drug Administration,s Adverse Event Reporting System (FAERS) Database. *Antibiotics (Basel)*, 2022; 11(10): 1367, doi.10.3390/antibiotics11101367.
15. Lin X, Kuck U. Cephalosporins as key lead generation beta-lactam antibiotics. *Appl Microbiol Biotechnol*, 2022; 106(24): 8007-8020.
16. Fontana R, Cornaglia G, Ligozzi M, Mazzanol A. The final goal: penicillin-binding proteins and the target of cephalosporins. *Clin Microbiol Infect*, 2000, 6(S3): 34-40.
17. Macy E, Contreras MS. Adverse reactions associated with oral and parenteral use of cephalosporins: a retrospective population based analysis. *J Allergy Clin Immunol*, 2015; 135(3): 743-752
18. Harrison CJ, Bratcher B. Cephalosporins: a review. *Pediatric Rev*, 2008; 29(8): 265-273.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

19. Rhyou HI, Doo GE, Yoon J, Ha CY, Nam HJ, Woo SD, et al. Clinical characteristics and risk factors for cefaclor-induced immediate hypersensitivity: a retrospective observation at two university hospitals in Korea. *Allergy Asthma Clin Immunol*, 2021; 17(20): doi10.1186/s13223-020-00523-8.
20. Rhou HI. Current status of cefaclor adverse drug reaction in Dong-A University Hospital. *J Allergy Clin Microbiol*, 2020; 45(S2): AB94, doi.org/10.1016/j.jaci/2019.12.593.
21. Cai Y, Yang L, Shanguan X, Huang R. Status and safety signals of cephalosporins in children: a spontaneous reporting: database study. *Front Pharmacol*, 2021; 12: doi.org/10.3389/fphar.2021.736618.
22. Kancharlapalli PK, Donthu RK, Mohammed AS, Pasam RS. Disulfiram-like reaction with cefixime. *J Psychiatry Spectrum*, 2023; 2(2): 118-119.
23. Joo MK, Shin YJ, Kim BH. Cefaclor causes vagus nerve-mediated depression-like symptoms with gut dysbiosis in mice. *Sci Rep*, 2023; 13: 15529, doi.10.1038/s41598-023-42690-1.
24. King BA, Geelhoed GC. Adverse skin and joint reactions associated with oral antibiotics in children: the role of cefaclor in serum sickness-like reactions. *J Paediatric Child Health*, 2003; 39(9): 677-681.
25. Huth ME, Ricci AJ, Cheng AG. Mechanisms of aminoglycosides ototoxicity and target of hair cell protection. *Int J Otolaryngol*, 2011; 2011: e937861.
26. Rivetti S, Romano A, Mastrangelo S, Attina G, Maurini P, Ruggiero A. Aminoglycosides-related ototoxicity: mechanisms, risk factors and prevention in pediatric patients. *Pharmaceuticals (Basel)*, 2023; 6(10): 1353, doi.10.3390/ph1601353.
27. Parajuli NP, Mandava CS, Pavlov MY, Sanyal S. Mechanistic insights into translation inhibition by aminoglycosides antibiotic arbekacin. *Nucleic Acids Res*, 2021; 49(12): 6880-6892.
28. Fu X, Wan P, Li P, Guo S, Zhang Y, An Y, et al. Mechanism and prevention of ototoxicity induced by aminoglycosides. *Front Cell Neurosci*, 2021; 15: 692762. doi.10.3389/fncel.2021.692762.
29. Nayak-Rao S. Aminoglycosides use in renal failure. *Indian J Nephrol*, 2010; 20(3): 121-124.
30. McWilliams SJ, Antoine DJ, Smith RL, Pinnohamed M. Aminoglycosides-induced nephrotoxicity in children. *Pediatric Nephrol*, 2017; 3: 2015-2025.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

31. Krause KM, Sario AW, Kane TR, Connolly LE. Aminoglycosides: an overview. Cold Spring Harb Perspect, 2016; 6(6): e027029.
32. Millano AR, Mora AY, Villagra NA, Bcarey SA, Hillago AA. Biological effects of quinolones: a family of broad-spectrum agents. Molecules, 2021; 26(23): doi.10.3390/molecules26237153.
33. Hooper DC, Jacoby GA. Topoisomerase inhibitors: fluoroquinolone mechanisms of action and resistance. Cold Spring Harb Perspect Med, 2016; a025320.
34. Spencer AC, Pauda SS. DNA gyrase as a target of quinolones. Biomedicines, 2023; 11(2): 371, doi.10.3390/biomedicines1102037.
35. Zobeiri E, Sadkhanlou RA, Salami S, Mardani K, Ahmadi A. The effect of ciprofloxacin on spermatozoa DNA damage, infertility and potential early embryonic development in NMRI mice. Vet Res Forum, 2012; 3(2): 131-135
36. Baggio D. Anand-Rajah MR. Fluoroquinolones antibiotics and adverse events. Aust Prescr, 2021; 44:161-164.
37. Gazi Z. Teratogenic effect of ciprofloxacin. Alq J Med Appl Sci, 2001; 4(2)114-125.
38. Onemu SO, Ibeh IN. The effect of quinolone chemotherapy on spermatozoal concentration of Nigerian males. Nig J Biomed Engineering, 2001; 1(1): 12-15.
39. Rusu A, Munteanu AC, Arbaasi EM. Overview of the side effects of antimicrobial fluoroquinolones new drugs versus old drugs: a step forward in the safety profile. Pharmaceutics, 2023; 15(3): 804, doi.org/10.3390/pharmaceutics.15030804.
40. Mandell L, Tillotson G. Safety of fluoroquinolones: an update. Can J Infect Dis, 2002; 13(1):54-61.
41. Aspinall CL, Good CB, Jang R, McCaren M, Dung D, Cunningham HE. Severe dysglycemia with the fluoroquinolones: a class effects. Clin Infect Dis, 2009; 49(3): 402-408.
42. Daneman N, Lu H, Redelmeier DA. Fluoroquinolones associated with severe adverse events: a longitudinal cohort study. BMJ Open, 2015; 5(11): e010077
43. Kaden T, Graf K, Kennert K, Li R, Mosia SM, Raasch M. Evaluation of drug-induced liver toxicity of trovafloxacin in a human microphysiological model. Scientific Rep, 2023; 13: 13338, doi.org/10.1038/s41598-023-4004-z.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. Elite Journal of Medicine, 2024; 2(7): 1-20

44. Ovung A, Bhattacharyya J. Sulfonamide drugs: structure, antibacterial property, toxicity and biophysical interactions. *Biophys Rev*, 2021; 13(2): 259-272.
45. Venkatesan M, Fruci M, Verellen LA, Skarina T, Mesa N, Flick R, et al. Molecular mechanisms of plasmid-borne resistance to sulfonamide antibiotics. *Nat Comm*, 2023; 14: 4031, doi.org/10.1038/s41467-023-39778-7.
46. Mockenhaupt M, Viboud C, Dunant A, Naldi L, Halevy S, Bavinck JN, et al. Stevens-Johnson syndrome and toxic epidermal necrosis: assessment of medication risks with emphasis on recently marketed drugs: The EuroSCAR-study. *J Invest Dermatol*, 2008; 128(1): 35-44.
47. Kastylevsky GC, Vial T, Descotes J. Allergic adverse reactions to sulfonamides. *Curr Allergy Asthma Rep*, 2002; 2(1): 16-25
48. Savage E, Murphy M, Hoghes S. Hemolytic anemia due to sulfonamide-induced methemoglobinemia. *J Gen Internal Med*, 2016; 31(9): 1067-1071.
49. Yarlagadda SG, Perazella MA. Drug-induced crystal neuropathy: an update. *Exp Opin Drug Saf*, 2008; 7(2): 147-158.
50. Azencot R, Saint-Jacques C, Heymann JP, Frochot V, Daudon M, Letavernier E. Sulfamethoxazole-induced crystal neuropathy: characterization and prognosis in a case series. *Sci Rep*, 2024; 14(1): 6078, doi.10.1038/s41598-024-56322-9.
51. Grossman TH. Tetracycline antibiotics and resistance. *Cold Spring Harb Perspect Med*, 2016; 2016: a025387, doi.10.1101/cshperspect.a025387.
52. Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology and epidemiology of bacteria resistance. *Microbiol Rev*, 2001; 65(2): 232-260.
53. Del-Rosso JQ. Oral tetracycline in the management of acne-vulgaris: current perspective on clinical use and recent findings with a new double score small tablet formulation. *The J Clin Aesthetic Dermatol*, 2017; 10(5): 13-18.
54. Orylska-Ratynska M, Placek W. Tetracycline - an important therapeutic tool for dermatologists. *Int J Environ Res Public Health*, 2022; 19(12): 7246, doi.10.3390/ijerph1912746.
55. Lenz KD, Klosterman KE, Mukundan H, Sunderland JZ. Macrolides: from toxins to therapeutics. *Toxins (Basel)*, 2021; 13(5): 347, doi.10.3390/toxins13050347.
56. Siempos II, Dimopoulos G, Korbila IP, Manta K, Falagas MG. Macrolides, quinolones and amoxicillin-clavulanate: a meta-analysis. *Eur Res J*, 2011; 29: 1127-1137.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

57. Kanoh S, Rubin BK. Mechanism of action and clinical application of macrolides as immuno-modulatory medications. *Clin Microbiol Rev.* 2007; 23(3): 590-615.
58. Mazzei T, Mini E, Novelli A. Adverse effects of macrolides antibacterials. *Drug Safety*, 1993; 9: 346-364.
59. Vazquez-Laslop N, Mankin AS. How macrolide antibiotics work. *Trends In Biomed Sci*, 2018; 43(9): P668-684.
60. Spizek J, Rezanka T. Lincosamides, clinical structure, biosynthesis, mechanism of action, resistance and application. *Biochem Pharmacol*, 2017; 133: 20-28.
61. Matzor D, Eval Z, Benhamou RI. Shalev-Benani M, Halfor Y Krupin M, et al. Structural insights of lincosamides targeting the ribosome of *Staphylococcus aureus*. *Nucleic Acids Res*, 2017; 45(17): 10284-10292.
62. Tenson T, Lovmar M, Ehrenberg M. The mechanism of action of macrolides, lincosamides and streptogamin-B reveals the nascent peptide exit path in the ribosome. *J Mol Biol*, 2003; 330: 1005-1014.
63. Kukzycka-Mierzejewska K, Tryslka J, Sadlej J. Quantum mechanical studies of lincosamides. *J Molecular Model*, 2012; 18: 2727-2740.
64. Armillei MK, Lomakin IB, Del-Rosso JQ, Grada A, Bunick CG. Scientific rationale for clindamycin use in the treatment of dermatological disease. *Antibiotics*, 2024; 13(3): 270, doi.10.3390/antibiotics.13030270.
65. Chen CW, Pavlova JA, Lukianov AG, Tereshcenkov AG, Makarov GI, Khairullina ZZ, et al. Binding and action of triphenyl-phosphonium analog of chloramphenicol upon bacteria ribosome. *Antibiotics*, 2021; 10(4): 390. Doi.org/10.3390/antibiotics.10040390.
66. Dinos GP, Athanassopoulos CM, Giannopoulou AG, Vlachogiannis GE, Papioannou D, et al. Chloramphenicol derivatives as antibacterial and anticancer agents: historic problems and current solutions. *Antibiotics*, 2016; 5(2): doi.10.3390/antibiotics.5020020.
67. Dzipova O, Benes J. Role of chloramphenicol in current clinical practice. *Klin Mikrobiol Infekc Lek*, 2022; 28(4): 101-105.
68. Nitzan O, Suponitzky U, Kennes Y, Chazan B, Raul R, Colodner R. Is chloramphenicol making a comeback? *Isr Med Assoc J*, 2010; 12(6): 371-374.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

69. Nitzan O, Kennes Y, Colodner R, Saliba W, Edelsteinl H, Raz R, et al. Chloramphenicol use and susceptibility pattern in Israel: a national survey. *Isr Med Assoc J*, 2015; 17(1): 27-31.
70. Sood S. Chloramphenicol – a potent armament against multidrug resistant (MDR) Gram-negative bacilli. *J Clin Diag Res*, 2016; 10(2): DC01-3.
71. Rohana H, Hage-Cohen A, Azrad M, Perez A. Trends of changes in chloramphenicol during the years 2017-2020: a retrospective report from Israel. *Antibiotics (Basel)*, 2023; 12(2): 196, doi.10.3390/antibiotics12020196.
72. Papp-Wallace KM, Endimiani A Taracila MA, Bonomo RA. Carbapenems: past, present and future, 2011; 55(11): 4943-4960.
73. Armstrong T, Fenn SJ, Hardie KR. JMM profile: carbapenems: a broad-spectrum antibiotic. *J Med Microbiol*, 2021; 70(12): 001462, doi.10.1099/jmm.0.001462.
74. Aurillo C, Sansone P, Barbasi M, Pota V, Giaccari G, Coppolino F, et al. Mechanism of action of carbapenems resistance. *Antibiotics (Basel)*. 2022; 11(3): 421, doi.10.3390/antibiotics11030421.
75. Yang P, Chen Y, Tiang S, Shen P, Lu X, Xiao Y. Association between antibiotic consumption and the rate of carbapenem-resistant Gram-negative bacteria from China on 153 tertiary hospital data in 2014. *Antimicrobial Resistance Infect Control*, 2018; 7(37): doi.org/1186/s13756-018-0430-1.
76. Finch RG, Eliopoulos GM. Safety and efficacy of glycopeptide antibiotics. *J Antimicrob Chemother*, 2005; 55(Suppl 2): ii5-13.
77. Butler MS, Hansford KA, Blaskovich MA, Halai R, Cooper MA. Glycopeptide antibiotics: back to the future. *Antibiotics*, 2014; 67: 631-644.
78. Zeng D, Debabor D, Hartsell TL, Cano RJ, Adams S, Schuyler, et al. Approved glycopeptide antibacterial drugs. *Cold Spring Harb Perspect*, 2016; 6(12): a026098.
79. Blaskovich MA, Hansford KA, Butler MS, Jia ZG, Mark AE, Cooper MA. Development of glycopeptide antibiotics. *ACS Infect Dis*, 2018; 4(5): 715-735.
80. Huang V, Clayton NA, Welker KH. Glycopeptide hypersensitivity and adverse reactions. *Pharmacy (Basel)*, 2020; 8(2): 70; doi.10.3390/pharmacy80220070.
81. Treshchalin MI, Polozkova VA, Moiseenko B, Treshalina HM, Shchekotikhin AE, Pereverzeva ER. Evaluation of toxic properties of new glycopeptide flavancin on rats. *Pharmaceuticals*, 2022; 15(6): 661; doi.org/10.3390/ph15060661.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigwe PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

82. Van-Banbeke F, Van-Lacthem Y, Courvalin P, Tulkens PM. Glycopeptide antibiotics: from conventional molecules to new derivatives. *Drugs*, 2004; 64(9): 913-936.
83. Binda E, Marinelli F, Marcone GL. Old and new glycopeptide antibiotics and resistance. *Antibiotics*, 2014; 3(4): doi.10.3390/antibiotics/s3040572.
84. Singh M, Chang J, Coffman L, Kim SJ. Hidden mode of action of glycopeptide antibiotics. Inhibition of wall teichoic acid biosynthesis. *J Phys Chem B*, 2017; 121(16): 3925-3932.
85. Dean CR, Barkan DT, Bermingham A, Blais J, Casey F, Casarez A, et al. Mode of action of the monobactam LY228 and mechanisms: decreasing in-vitro susceptibility in *Escherichia coli* and *Klebsiella pneumonia*. *Antimicrob Agents Chemother*, 2018; 62(10): e01200-01218.
86. Finberg RW, Guharoy R. Monobactams. In: *Clinical use of anti-infective agents*. Springer, New York, NY, 2012; https://doi.org/10.1007/978-8-14614-1064-3_6.
87. Kapoor S, Gathwala G. Aztreonam. *Indian J Pediatr*, 2004; 41(4): 359-364
88. Stefani S, Bongiorno D, Mongelli G, Gampaile F. Linezolid resistance in staphylococci. *Pharmaceuticals*, 2010; 3(7): 1988-06.
89. Dryden MS. Linezolid pharmacokinetics and pharmacodynamics in clinical treatment. *Antimicrob Chemother*, 2011; 66(Suppl 4): iv7-iv15.
90. Bousa E, Munoz P. Linezolid: pharmacokinetic characteristics and clinical studies. *Clin Microbiol Infect*, 2012; 7(S4): 75-82.
91. Sadovsky R. Linezolid is the first of the new class of antibiotics. *Am Fam Physician*, 2003; 68(1): 166-7.
92. Hashemian SM, Farhadi T, Ganjparvar M, Linezolid: a review of its properties and use in critical care. *Drug Des Devel Ther*, 2018; 12: 1759-67.
93. Gan C, Ng HF, Ngeow YF. Mechanism of linezolid resistance in mycobacteria. *Pharmaceuticals*, 2023; 16(6): 784: doi.org/10.3390/ph16060784.
94. Thirot H, Briguet C, Fripiat F, Jacobs F, Holemans X, Tulkens PM, et al. Clinical use and adverse drug reactions of linezolid: a retrospective study in four Belgian Hospital Centers. *Antibiotics (Basel)*, 2021; 10(5): 530, doi.10.3390/antibiotics10050530.
95. Azzouz A, Preuss CV. Linezolid. StatPearls [updated March 01, 2024]. In: StatPearls [Internet], Treasure Island (FL): <https://ncbi.nlm.nih.gov/books/NBK53793>.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

96. Meera M, Kanna N, Chidambaram M. A rare adverse effect following prolonged linezolid therapy. *Indian J Pediatr*, 2023; 90: 618, doi.org/10.1007/s12098.
97. Sousa M, Pozniak A, Boffio M. Pharmacokinetics of drug interactions involving rifampicin and antimalarial drugs. *J Antimicrob Chemother*, 2008; 62(5): 872-8.
98. Campbell EA, Korzheva N, Mustaev A, Murakami K, Nair S, Goldfarb A, et al. Structural mechanism for rifampicin of bacterial RNA polymerase. *Cell*, 2001; 104(6): 901-12.
99. Suresh AB, Rosani A, Patel P, Wadwa R. Rifampicin In: StatPearls [Internet], Treasure Island (FL), StatPearls Publishing 2024. (Accessed March 5, 2024, PMID: 32491420).
100. Chen J, Raymond K, Roles of rifampicin in drug-drug interactions: underlying molecular mechanisms involving the nuclear pregnane-X receptor. *Ann Clin Microbiol Antimicrob*, 2006; 5(3): doi.10.1186/1476-0711-5-3.
101. Nicolini A Perazzo A, Gatto P, Piroddi IM, Barlascini S, Karamicheli S, et al. A rare adverse reaction to ethambutol: a drug-induced hemolytic anemia. *Int J Tuber Lung Dis*, 2016; 20(5):704-5.
102. Si M, Li H, Chen Y, Peng H. Ethambutol and isoniazid-induced severe neurotoxicity in a peritoneal dialysis. *BMJ Case Report*, 2018; bcr2017223187. Doi.10.1136/bcr-2017-223187.
103. Ramirez J, Guarner F, Fernandez LB, Maruy A, Sdepanam VL, Cohen H. Antibiotics as major disruptors of gut microbiota. *Front Cell Infect Microbiol*, 2020; 10: 572912, doi.10.3389/fcimb.2020.572912.
104. Patangia DV, Ryan CA, Dempsey E, Ross RP, Statin C. Impact of antibiotics on human microbiome and consequences for host health. *Microbiol Open*, 2022; 11(1): e1260.
105. Bich VT, Le NG, Barnett D, Chan J, Van-Best N, Tien TD, et al. Moderate and transient impact of antibiotic use on gut microbiota in a Vietnamese cohort. *Sci Rep*, 2022; 12: 20189, doi.org/10.1038/s41598-020-24488-9.
106. Matzaras R, Anagnostou N, Nikopoulou A, Tsiakas I, Christaki E. The role of probiotics in inflammation associated with major surgery: a narrative review. *Nutrients*, 2023; 15(6): 1331, doi.10.3390/nu15061331.
107. Lathakumari RH, Vajravelu LK, Satheesan A, Ravi S, Thulukanem J. Antibiotics and the gut microbiome: understanding the impact on human. *Med Microecology*, 2024, 20: 100106, doi.org/10.1016/medmicro.2024.100106.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

108. Thanert R, Swahney SS, Shwartz D, Dantos G. The resistance within; antibiotic disruption of the gut microbiome and resistance dynamics in infancy. *Cell Host Microbiome*, 2022; 30(5): 675-83.
109. Luchen CC, Chubuye M, Spiker R, Simyand M, Chisengo C, Chilengi, et al. Impact of antibiotics on low to middle-income countries: a systematic review. *PLoS Med*, 2023; 20(6): e1004235.
110. Ugwu OP, Alum EU, Okon AB, Obeagu EI. Mechanisms of microbiota modulation for health, disease and therapeutic intervention. *Medicine (Baltimore)*, 2024; 103(19): e38088.
111. Langdon A, Crook N, Dantas G. The effects of antibiotics on the microbiome throughout development and alternative approaches for therapeutic modulation. *Genome Med*, 2016; 8: 39, doi.10.1186/s13073-0294-2.
112. Yao J, Carter RA, Vuagniaux G, Barbier M, Rosch JW, Rock CO. A pathogen-selective antibiotic minimizes disturbance to the microbiome. *Antimicrob Agents Chemother*, 2016; 60(7): 4264-73.
113. Shwartz DJ, Langdon AE, Dantas G. Understanding the impact of antibiotic perturbation on the human microbiome. *Genome Med*, 2020; 12(82): doi.org/10.1186/s13073-020-00782-x.
114. Leckha S, Terrel CL, Edson RS. General principles of antimicrobial therapy. *Mayo Clin Pract*, 2011; 86(2): 156-67.
115. Tamma PD, Avdic E, Li DX, Dzintars K, Cosgrove SE. Association of adverse events with antibiotic use in hospitalized patients. *JAMA Intern Med*, 2017; 177(9): 1308-1.
116. Gordhon Y, Padayachee N, Evaluating the knowledge, attitude and practices in healthcare workers towards adverse drug reaction reporting at a public tertiary hospital in Johannesburg. *Int J Africa Nursing Science*, 2020; 12: doi.org/j.ijans.2020.100191.
117. Liang WS, Jones BB, Smalley S, Snyder M, Goetz LH, Schork N. Emerging therapeutic drug monitoring technology: considerations and opportunities in precision medicine. *Front Pharmacol*, 2024; 15: 1348112.
118. Siddiqui MK, Lazum J, Coenem M, Mahmoudpour SH, Pharmacogenomics of adverse drug reactions (ADRs). *Front Gene*, 2022; 13: doi.org/10.1089/fgene.2020.859909.
119. Yang J, Zhang L, Guan Y, Song C. Analysis of antimicrobial management and rational use of antibiotics. *Eur J Hosp Pharm*, 2020; 27(5): 286-91.

Citation: Onemu SO, Onemu-Metitiri MO, Odeyemi O, Uyigue PO, Obeagu EI. Unveiling Platelet Dynamics in ART-Treated HIV Patients: A Comprehensive Review. *Elite Journal of Medicine*, 2024; 2(7): 1-20

120. Wolf U, Baust H, Neef R, Steinke T. Individual pharmacotherapy management (IPM)-IV: optimized usage of approved antimicrobials addressing under-recognized adverse drug reactions and drug-drug interactions in polypharmacy. *Antibiotics*, 2022; 1381, doi.org/10.3390/antibiotics.11101381.
121. Small SM, Bacher RS, Jost SA. Disulfiram-like reaction involving ceftriaxone in a pediatric patient. *J Pedriaric Pharmacol Ther*, 2018; 23(3): 168-72.
122. Roberts JA, Norris R, Pareson DL, Martin JH. Therapeutic drug monitoring of antimicrobials. *Br J Clin Pharmacol*, 2011; 73(1): 27-36.
123. Abdul-Aziz MH, Alfenaar IW. Antimicrobbial therapeutic drug monitoring in critically ill adult patients: a position paper. *Intensive Care Med*, 2020; 46: 1127-53.
124. Sjoval F, Lanckohr C, Bracht H. What's new in therapeutic drug monitoring? *Intensive Care Med*, 2023; 49: 857-59.
125. Satterfield J, Miesner Ar, Percival KM. The role of education in antimicrobial stewardship. *J Hosp Infect*, 2020; 105(2): 130-41.
126. Watkins RR. Antibiotic stewardship in the era of precision medicine. *JAC Antimicrob Resist*, 2022; 4(3): dlac066, doi.org/10.1093/jacamr/dlac066.
127. Veluscek M, Bajrovic FF, Strle F, Stupico D. Doxycycline photosensitivity in patients treated for erythema migrans. *BMC Infect Dis*, 2018; 18; 365, doi.org/10.1186/s12879-018-3270-y.