

Malaria and Low Birth Weight: Understanding the Linkages

*Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²

¹Department of Medical Laboratory Science, Kampala International University, Ishaka, Uganda.

²School of Nursing Science, Kampala International University, Ishaka, Uganda.

*Corresponding authour: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, emmanuelobeagu@yahoo.com, ORCID: 0000-0002-4538-0161](#)

Abstract

Malaria is a significant public health concern, particularly affecting pregnant women and their unborn children in tropical and subtropical regions. This review investigates the relationship between malaria and low birth weight (LBW), a critical determinant of neonatal and infant health. LBW, defined as a birth weight of less than 2,500 grams, is closely linked to increased risks of infant morbidity and mortality. The pathophysiological mechanisms connecting malaria to LBW include placental malaria, maternal anemia, systemic inflammation, and nutrient deficiency. Placental malaria disrupts nutrient and oxygen transfer to the fetus, while malaria-induced anemia reduces oxygen delivery, both contributing to intrauterine growth restriction (IUGR). Additionally, systemic inflammation triggered by malaria and resulting maternal nutrient deficiencies further compromise fetal development. Epidemiological evidence from malaria-endemic regions consistently shows a higher prevalence of LBW, emphasizing the need for targeted interventions. Effective public health strategies are crucial to mitigate the adverse effects of malaria on birth weight. Interventions such as intermittent preventive treatment in pregnancy (IPTp), insecticide-treated bed nets (ITNs), prompt diagnosis and treatment of malaria, and maternal nutrition programs have shown promise in reducing the incidence of LBW. By addressing these linkages through comprehensive health initiatives, it is possible to improve maternal and child health outcomes in regions heavily burdened by malaria.

Keywords: *Malaria, Low Birth Weight, Placental Malaria, Maternal Health, Infant Mortality, Anemia, Intrauterine Growth Restriction (IUGR), Public Health, Preventive Strategies*

Introduction

Malaria remains one of the most formidable public health challenges, particularly in tropical and subtropical regions where the disease is endemic. Caused by Plasmodium parasites, malaria is transmitted to humans through the bites of infected Anopheles mosquitoes. According to the World
Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

Health Organization (WHO), there were an estimated 229 million cases of malaria worldwide in 2019, resulting in over 400,000 deaths, with a significant proportion occurring in children under five years old and pregnant women. The burden of malaria on pregnant women is of particular concern due to the severe health implications for both the mother and the developing fetus. Pregnant women are more susceptible to malaria infection due to changes in their immune system and increased attractiveness to mosquitoes. The immune changes during pregnancy, particularly the modulation of cellular immunity, reduce the mother's ability to mount an effective response against the malaria parasite. This vulnerability is further exacerbated in regions with high malaria transmission, where repeated exposure to the parasite can result in cumulative adverse health outcomes. One of the most significant consequences of malaria during pregnancy is low birth weight (LBW), which is a critical factor influencing neonatal and infant morbidity and mortality. Low birth weight, defined by the WHO as a birth weight of less than 2,500 grams, is a major public health concern. Infants born with LBW face an increased risk of neonatal complications, developmental delays, and higher mortality rates. The association between malaria and LBW is well-documented, particularly in malaria-endemic regions. Malaria can directly affect fetal growth and development, leading to intrauterine growth restriction (IUGR) and preterm birth, both of which are primary causes of LBW. Understanding the mechanisms through which malaria impacts fetal growth is crucial for developing effective interventions.¹⁻¹⁰

Placental malaria is a key mechanism linking malaria infection to LBW. When a pregnant woman is infected with *Plasmodium falciparum*, the deadliest malaria parasite, the infected erythrocytes sequester in the placenta. This sequestration leads to placental inflammation, impaired placental function, and reduced nutrient and oxygen transfer to the fetus. As a result, the fetus may suffer from IUGR, leading to LBW. Additionally, malaria-induced maternal anemia, a common complication of malaria during pregnancy, reduces the oxygen-carrying capacity of the blood, further restricting fetal growth. Systemic inflammation is another pathway through which malaria contributes to LBW. Malaria infection triggers the release of pro-inflammatory cytokines and other immune mediators that can cross the placental barrier and affect the fetus. This inflammatory response can impair placental function and disrupt fetal development. Chronic malaria infection can also lead to maternal malnutrition and micronutrient deficiencies, which are essential for fetal growth and development. Nutrient deficiencies exacerbate the risk of LBW, highlighting the multifactorial impact of malaria on pregnancy outcomes. Epidemiological studies consistently show a strong association between malaria infection during pregnancy and increased incidence of LBW. Research in sub-Saharan Africa, Southeast Asia, and other malaria-endemic regions reveals that pregnant women infected with malaria are more likely to give birth to LBW infants compared to their uninfected counterparts. These studies underscore the importance of malaria prevention and treatment during pregnancy to improve birth outcomes. Interventions such as intermittent preventive treatment in pregnancy (IPTp) with antimalarial drugs and the use of insecticide-treated bed nets (ITNs) have been shown to significantly reduce the risk of malaria and associated adverse pregnancy outcomes.¹¹⁻¹⁵

The global public health community has recognized the need for comprehensive strategies to address the burden of malaria in pregnant women. The WHO recommends a combination of IPTp, ITNs, and prompt diagnosis and treatment of malaria to protect pregnant women and their fetuses.

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

These interventions aim to reduce the incidence of malaria, prevent complications such as anemia and placental malaria, and ultimately improve birth outcomes. Additionally, maternal nutrition programs that address micronutrient deficiencies can complement malaria prevention efforts by supporting optimal fetal growth and development. Despite the availability of effective interventions, challenges remain in achieving widespread implementation and coverage. Health system constraints, including limited access to healthcare services, inadequate infrastructure, and insufficient funding, hinder the delivery of malaria prevention and treatment programs. Cultural beliefs and practices, as well as lack of awareness about the importance of malaria prevention during pregnancy, also pose barriers to intervention uptake. Addressing these challenges requires a multifaceted approach, including strengthening health systems, enhancing community engagement, and increasing investment in maternal and child health programs.¹⁶⁻²⁰

Pathophysiological Mechanisms

The pathophysiological mechanisms linking malaria to low birth weight (LBW) are complex and multifaceted. They involve direct effects of the malaria parasite on the placenta and fetus, as well as indirect effects mediated through maternal health. Placental malaria is a primary mechanism by which malaria contributes to LBW. When a pregnant woman is infected with *Plasmodium falciparum*, the infected red blood cells (iRBCs) adhere to the placental tissue. This adhesion is mediated by specific parasite proteins, such as VAR2CSA, which bind to chondroitin sulfate A on the placental surface. The accumulation of iRBCs in the placenta leads to inflammation, characterized by the infiltration of immune cells and the production of pro-inflammatory cytokines. This inflammatory response impairs placental function by disrupting the architecture of the placental tissue and interfering with the exchange of nutrients and oxygen between the mother and fetus. The resulting hypoxia and nutrient deficiency can severely restrict fetal growth, leading to intrauterine growth restriction (IUGR) and ultimately LBW. Additionally, placental malaria can cause structural changes in the placenta, such as thickening of the placental membrane, further hindering nutrient and oxygen transfer. Malaria-induced maternal anemia is another significant contributor to LBW. Anemia in pregnant women is primarily caused by the destruction of red blood cells by the malaria parasite, as well as by hemolysis and the bone marrow's inability to compensate for the loss of red blood cells. Anemia reduces the oxygen-carrying capacity of the blood, leading to decreased oxygen delivery to the fetus. Chronic hypoxia can impair fetal growth and development, resulting in LBW.²⁵⁻³⁰

In addition to oxygen deprivation, anemia can exacerbate placental insufficiency, further limiting nutrient transfer to the fetus. The severity of anemia correlates with the risk of LBW, highlighting the importance of preventing and treating anemia in pregnant women to improve birth outcomes. Systemic inflammation triggered by malaria infection plays a critical role in the pathogenesis of LBW. The immune response to malaria involves the release of pro-inflammatory cytokines, such as tumor necrosis factor-alpha (TNF- α), interleukin-1 (IL-1), and interferon-gamma (IFN- γ). These cytokines can cross the placental barrier and affect fetal development. Elevated levels of pro-inflammatory cytokines can lead to endothelial dysfunction and increased vascular permeability, resulting in placental edema and reduced blood flow to the fetus. This inflammatory milieu can impair placental function and contribute to IUGR and LBW. Additionally, chronic

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

inflammation can affect the mother's overall health, further compromising fetal growth. Malaria can exacerbate maternal malnutrition and micronutrient deficiencies, which are critical for fetal growth and development. Infected women may experience reduced appetite and increased metabolic demands, leading to inadequate nutrient intake. Furthermore, malaria can interfere with the absorption and utilization of essential nutrients, such as iron, folate, and zinc. Nutrient deficiencies during pregnancy can impair placental function and fetal growth, increasing the risk of LBW. For example, iron deficiency can lead to anemia, while folate deficiency can affect DNA synthesis and cell division, both of which are crucial for fetal development. Addressing these deficiencies through nutritional interventions can help mitigate the impact of malaria on pregnancy outcomes.³¹⁻³⁵

Malaria infection can lead to hypoglycemia, particularly in severe cases. Hypoglycemia occurs when blood glucose levels drop to abnormally low levels, depriving the fetus of a critical energy source. This condition can result from increased glucose utilization by the malaria parasite, impaired gluconeogenesis, or the effects of antimalarial treatments. Hypoglycemia during pregnancy can contribute to fetal distress and growth restriction, further increasing the risk of LBW. Malaria infection induces oxidative stress, characterized by the excessive production of reactive oxygen species (ROS) and the depletion of antioxidants. Oxidative stress can damage placental cells and tissues, disrupt placental function, and impair fetal growth. The placenta's role as a barrier and mediator of nutrient and oxygen transfer makes it particularly vulnerable to oxidative damage. Strategies to reduce oxidative stress, such as antioxidant supplementation, may help protect against LBW in malaria-infected pregnancies. Normal fetal development requires proper angiogenesis, the formation of new blood vessels. Malaria infection can disrupt angiogenesis by altering the balance of pro-angiogenic and anti-angiogenic factors in the placenta. For instance, malaria-induced inflammation and oxidative stress can upregulate anti-angiogenic factors, such as soluble fms-like tyrosine kinase-1 (sFlt-1), while downregulating pro-angiogenic factors, like vascular endothelial growth factor (VEGF). This imbalance can impair placental blood flow and nutrient transfer, contributing to LBW. Recent research suggests that malaria infection during pregnancy can lead to epigenetic modifications, such as DNA methylation and histone modification, which can influence fetal growth and development. These epigenetic changes can alter the expression of genes involved in placental function and fetal growth, potentially leading to long-term health consequences for the child. Understanding the epigenetic impacts of malaria can provide new insights into the mechanisms underlying LBW and inform the development of novel therapeutic strategies.³⁶⁻⁴⁵

Epidemiological Evidence

The relationship between malaria infection during pregnancy and low birth weight (LBW) has been well-documented through numerous epidemiological studies. These studies highlight the significant impact of malaria on maternal and fetal health, particularly in regions where the disease is endemic. Epidemiological data consistently show a higher prevalence of LBW in regions with high malaria transmission. Sub-Saharan Africa, which bears the highest burden of malaria, also reports some of the highest rates of LBW. Studies have demonstrated that in malaria-endemic areas, pregnant women are at an increased risk of contracting malaria, which directly correlates

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. *Elite Journal of Nursing and Health Science*, 2024; 2(7):41-55

with higher rates of LBW. For instance, in a study conducted in Malawi, it was found that pregnant women with malaria were significantly more likely to give birth to LBW infants compared to those without malaria. Longitudinal studies tracking pregnant women over time provide robust evidence of the impact of malaria on birth weight. One such study in Tanzania followed a cohort of pregnant women from early pregnancy through delivery, documenting malaria episodes and birth outcomes. The study found that women who experienced malaria during pregnancy were more likely to deliver LBW infants compared to women who remained malaria-free. Cross-sectional studies conducted in various malaria-endemic countries, including Kenya and Nigeria, have also reported similar findings, reinforcing the association between malaria infection and LBW. Intervention studies have provided compelling evidence that malaria prevention and treatment can significantly reduce the incidence of LBW. Intermittent preventive treatment in pregnancy (IPTp) with sulfadoxine-pyrimethamine (SP) has been shown to reduce the risk of malaria and improve birth outcomes. A landmark randomized controlled trial in Mozambique demonstrated that women who received IPTp had a lower risk of delivering LBW infants compared to those who did not receive the intervention. Similarly, the use of insecticide-treated bed nets (ITNs) has been associated with reduced malaria incidence and improved birth weights in pregnant women.⁴⁶⁻⁵⁵

Regional variations in the impact of malaria on LBW highlight the importance of local context in understanding and addressing the problem. For example, studies in Southeast Asia have shown that while malaria is a significant risk factor for LBW, the specific strains of *Plasmodium* and local health practices can influence the outcomes. In Thailand, research indicated that *Plasmodium vivax*, though less deadly than *P. falciparum*, still posed a significant risk for LBW, emphasizing the need for comprehensive malaria control measures targeting all malaria species. The seasonality of malaria transmission also plays a role in the incidence of LBW. In regions where malaria transmission is seasonal, pregnant women are at increased risk of malaria infection during the high transmission seasons, leading to higher rates of LBW during these periods. Studies from Burkina Faso and Ghana have documented seasonal variations in birth outcomes, with higher rates of LBW observed during and immediately after the peak malaria transmission season. Epidemiological evidence suggests that maternal age and parity influence the risk of malaria-associated LBW. Younger pregnant women and primigravidae (women in their first pregnancy) are at higher risk of malaria infection and its adverse outcomes, including LBW. This is partly due to the lack of acquired immunity to malaria that typically develops with repeated exposure. Studies in Uganda and Malawi have shown that primigravidae have higher malaria prevalence and higher rates of LBW compared to multigravidae (women with previous pregnancies). Socioeconomic factors, such as education level, income, and access to healthcare, significantly affect the incidence of malaria and LBW. Women from lower socioeconomic backgrounds often have limited access to malaria prevention and treatment services, increasing their risk of infection and adverse pregnancy outcomes. A study in Nigeria highlighted that women with higher education levels and better socioeconomic status had lower rates of malaria and LBW, underscoring the importance of addressing social determinants of health in malaria control programs. The epidemiological evidence underscores the critical need for integrated public health strategies to reduce the burden of malaria and improve birth outcomes. Preventive measures, such as IPTp, ITNs, and prompt diagnosis and treatment of malaria, are essential components of these strategies. Additionally, improving maternal nutrition and addressing socioeconomic barriers to healthcare access are

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. *Elite Journal of Nursing and Health Science*, 2024; 2(7):41-55

crucial for mitigating the impact of malaria on pregnancy outcomes. Public health campaigns that raise awareness about the risks of malaria in pregnancy and promote preventive behaviors are also vital.⁵⁶⁻⁶⁵

Public Health Interventions

Addressing the link between malaria and low birth weight (LBW) requires a comprehensive approach that integrates various public health interventions. These strategies are designed to prevent malaria infection in pregnant women, mitigate its adverse effects, and improve maternal and fetal health outcomes. This section discusses key public health interventions that have proven effective in reducing the incidence of malaria-associated LBW. Intermittent preventive treatment in pregnancy (IPTp) involves administering antimalarial drugs at scheduled intervals during pregnancy, regardless of whether the woman is infected with malaria. The World Health Organization (WHO) recommends IPTp with sulfadoxine-pyrimethamine (SP) for all pregnant women in areas of moderate to high malaria transmission. This intervention has been shown to significantly reduce the risk of malaria, maternal anemia, and LBW. IPTp works by clearing existing malaria infections and providing a prophylactic effect against new infections. Studies have demonstrated that IPTp with SP reduces placental malaria, increases mean birth weight, and decreases the incidence of LBW. For example, a randomized controlled trial in Tanzania found that women who received IPTp had a 38% lower risk of delivering LBW infants compared to those who did not receive the treatment. The use of insecticide-treated bed nets (ITNs) is a highly effective method for preventing malaria infection. ITNs create a physical barrier and kill or repel mosquitoes, reducing the incidence of malaria. WHO recommends that all pregnant women in malaria-endemic areas sleep under ITNs. ITNs have been shown to significantly reduce malaria morbidity and mortality, including during pregnancy. A study in Kenya found that pregnant women who used ITNs had a 25% lower risk of malaria infection and a 23% lower risk of delivering LBW infants compared to those who did not use ITNs. The widespread distribution and use of ITNs have been a cornerstone of malaria control programs and have contributed to substantial reductions in malaria transmission and related adverse pregnancy outcomes. Prompt diagnosis and effective treatment of malaria are crucial for preventing severe disease and complications in pregnant women. Rapid diagnostic tests (RDTs) and microscopy are commonly used to diagnose malaria, and treatment with effective antimalarial drugs should be initiated as soon as possible after diagnosis. Artemisinin-based combination therapies (ACTs) are the recommended first-line treatment for uncomplicated malaria in pregnant women in the second and third trimesters. Timely treatment of malaria in pregnancy can prevent the progression to severe disease, reduce the risk of maternal anemia, and improve birth outcomes. A study in Uganda showed that prompt treatment of malaria with ACTs significantly reduced the risk of LBW and improved maternal hemoglobin levels.⁶⁶⁻⁷⁵

Health Education and Community Engagement

Health education and community engagement are vital components of public health interventions aimed at reducing malaria and improving pregnancy outcomes. Educating pregnant women and communities about the importance of malaria prevention, the use of IPTp and ITNs, and the need

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

for prompt diagnosis and treatment can enhance the uptake and effectiveness of these interventions. Community health workers (CHWs) play a crucial role in delivering health education, distributing ITNs, and supporting pregnant women to access IPTp and treatment services. Community-based programs that involve CHWs have been successful in increasing the use of preventive measures and improving birth outcomes in malaria-endemic areas. For example, a program in Zambia that trained CHWs to provide malaria education and distribute ITNs resulted in increased ITN usage and reduced incidence of LBW. Strengthening health systems is essential for the successful implementation and sustainability of malaria interventions. This includes ensuring the availability and accessibility of IPTp, ITNs, diagnostic tools, and effective antimalarial treatments. Health system strengthening also involves training healthcare providers, improving supply chain management, and enhancing data collection and surveillance systems. Investment in health infrastructure, such as antenatal care (ANC) clinics, is critical for delivering malaria prevention and treatment services to pregnant women. Integrated ANC services that offer comprehensive care, including malaria prevention, nutritional support, and routine health check-ups, can improve maternal and fetal health outcomes. A study in Ghana demonstrated that integrating IPTp delivery into routine ANC services increased IPTp coverage and reduced the incidence of malaria and LBW.⁷⁶⁻⁸⁴

Nutritional interventions are important for addressing the indirect effects of malaria on maternal and fetal health. Malaria infection can exacerbate nutrient deficiencies, which are critical for fetal growth and development. Providing nutritional supplements, such as iron and folic acid, can help prevent anemia and support optimal fetal growth. Programs that combine malaria prevention with nutritional interventions have shown promise in improving birth outcomes. For example, a study in Malawi found that providing pregnant women with both IPTp and iron-folic acid supplements significantly reduced the incidence of LBW compared to providing either intervention alone. Monitoring and evaluation are crucial for assessing the effectiveness of public health interventions and making necessary adjustments. Regular monitoring of malaria incidence, IPTp and ITN coverage, treatment outcomes, and birth weights can provide valuable data for guiding program implementation and policy decisions. Evaluation studies can identify best practices and areas for improvement, ensuring that interventions are tailored to local contexts and needs. For instance, a multi-country evaluation of IPTp implementation highlighted the importance of continuous training for healthcare providers and community engagement to maintain high coverage and adherence rates. Policy and advocacy efforts are essential for sustaining political commitment and securing funding for malaria control programs. National and international policies that prioritize malaria prevention and maternal health can drive the allocation of resources and support the implementation of effective interventions. Advocacy campaigns can raise awareness about the impact of malaria on pregnancy and the importance of preventive measures. Engaging stakeholders, including governments, non-governmental organizations, and international agencies, can foster collaboration and coordination in the fight against malaria and its adverse effects on pregnancy.⁸⁵⁻⁸⁹

Conclusion

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

The intricate relationship between malaria and low birth weight (LBW) underscores the profound impact of malaria on maternal and fetal health, particularly in malaria-endemic regions. This review has highlighted the pathophysiological mechanisms, epidemiological evidence, and public health interventions that link malaria to adverse pregnancy outcomes, especially LBW. Malaria-induced placental changes, maternal anemia, systemic inflammation, nutrient deficiencies, hypoglycemia, oxidative stress, impaired angiogenesis, and epigenetic modifications are key mechanisms that contribute to LBW. These mechanisms disrupt the normal growth and development of the fetus, leading to intrauterine growth restriction and ultimately LBW. Epidemiological studies consistently show higher rates of LBW in malaria-endemic areas, with substantial variations based on regional, seasonal, and socioeconomic factors. These studies provide robust evidence that malaria prevention and treatment can significantly reduce the incidence of LBW.

Public health interventions, such as intermittent preventive treatment in pregnancy (IPTp), insecticide-treated bed nets (ITNs), prompt diagnosis and treatment, health education, health system strengthening, nutritional support, and policy advocacy, are essential components of a comprehensive approach to combat malaria and its adverse effects on pregnancy. These interventions have been proven effective in reducing malaria incidence, improving maternal health, and increasing birth weights. However, their successful implementation requires sustained political commitment, adequate funding, community engagement, and integrated health systems. The fight against malaria and its impact on pregnancy is far from over. Continued research is needed to further elucidate the mechanisms linking malaria to LBW, identify new therapeutic targets, and develop innovative interventions. Moreover, scaling up and sustaining effective public health strategies will be crucial to achieving significant reductions in malaria-associated LBW and improving overall maternal and child health in malaria-endemic regions.

References

1. Uneke CJ. Impact of placental *Plasmodium falciparum* malaria on pregnancy and perinatal outcome in sub-Saharan Africa: part III: placental malaria, maternal health, and public health. *The Yale journal of biology and medicine*. 2008;81(1):1.
2. Gontie GB, Wolde HF, Baraki AG. Prevalence and associated factors of malaria among pregnant women in Sherkole district, Benishangul Gumuz regional state, West Ethiopia. *BMC Infectious Diseases*. 2020; 20:1-8.
3. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri*. 2023; 6 (1). 2023;138. [links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf](https://doi.org/10.63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf).
4. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci*. 2016;3(3):10-8. [links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf](https://doi.org/10.5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf).
5. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria.

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. *Elite Journal of Nursing and Health Science*, 2024; 2(7):41-55

- J Res Med Dent Sci. 2021;9(4):145-8. [links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf](https://epjournals.com/journals/EJNHS/links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf).
6. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. International Research in Medical and Health Sciences. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
 7. Jakheng SP, Obeagu EI. Seroprevalence of human immunodeficiency virus based on demographic and risk factors among pregnant women attending clinics in Zaria Metropolis, Nigeria. J Pub Health Nutri. 2022; 5 (8). 2022;137. [links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf](https://epjournals.com/journals/EJNHS/links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf).
 8. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. Madonna University journal of Medicine and Health Sciences. 2022;2(2):1-9.
 9. Bonilla FA, Oettgen HC. Adaptive immunity. Journal of Allergy and Clinical Immunology. 2010;125(2): S33-40.
 10. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):1-9.
 11. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfectd with HIV in Umuahia. Int. J. Curr. Res. Med. Sci. 2017;3(5):100-104.
 12. Feeney ME. The immune response to malaria in utero. Immunological reviews. 2020 ;293(1):216-229.
 13. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. Medicine. 2023;102(46): e36166.
 14. Obeagu EI, Chijioke UO, Ekelozie IS. Malaria rapid diagnostic test (RDTs). Ann Clin Lab Res. 2018;6(4):275.
 15. Ogomaka IA, Obeagu EI. Methods of Breast Feeding as Determinants of Malaria Infections among Babies in IMO State, Nigeria. International Journal of Medical Science and Dental Research. 2019;2(01):17-24.
 16. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. Madonna University Journal of Medicine and Health Sciences. 2022;2(1):292-8. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
 17. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. J. Bio. Innov11 (3). 2022:936-941.
 18. Obeagu EI, Obeagu GU, Egba SI, Emeka-Obi OR. Combatting Anemia in Pediatric Malaria: Effective Management Strategies. Int. J. Curr. Res. Med. Sci. 2023;9(11):1-7.
 19. Hassan AO, Oso OV, Obeagu EI, Adeyemo AT. Malaria Vaccine: Prospects and Challenges. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):22-40.

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. Elite Journal of Nursing and Health Science, 2024; 2(7):41-55

20. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-9.
21. Moya-Alvarez V, Abellana R, Cot M. Pregnancy-associated malaria and malaria in infants: an old problem with present consequences. *Malaria journal*. 2014; 13:1-10.
22. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2022;9(9): 21-6.DOI: [10.22192/ijcrpcps.2022.09.09.003](https://doi.org/10.22192/ijcrpcps.2022.09.09.003)
23. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
24. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State. Nigeria. *Journal of Research in Medical and Dental Science*. 2021;9(12):462-. [links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://doi.org/10.21961/jrmads.61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf).
25. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022 Jun 21;5(2):113-121.
26. Obeagu EI. Gestational Thrombocytopaenia. *J Gynecol Women's Health*. 2023;25(3):556163. [links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf](https://doi.org/10.21961/jgmh.64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf).
27. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
28. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
29. Obeagu EI, Busari AI, Uduchi IO, Ogomaka IA, Ibekwe AM, Vincent CC, Chijioke UO, Okafor CJ, Okoroiwu HU, Adike CN. Age-Related Haematological Variations in Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):218-24.
30. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
31. Ogbonna CO, Obeagu EI, Ufelle SA, Ogbonna LN. Evaluation of haematological alterations in children infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(1):38-45.
32. Appay V. The physiological role of cytotoxic CD4+ T-cells: the holy grail? *Clinical & Experimental Immunology*. 2004;138(1):10-13.

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. *Elite Journal of Nursing and Health Science*, 2024; 2(7):41-55

33. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(1):25-36.
34. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med*. 2018;3(9):1-4.
35. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with *Plasmodium falciparum* malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2016;3(8):40-46.
36. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology*. 2019;41(3): e12573.
37. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elamba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research*. 2017;5(4):206.
38. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Irondi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal*. 2022;6(3):9-17.
39. Madekwe CC, Madekwe CC, Obeagu EI. Inequality of monitoring in Human Immunodeficiency Virus, Tuberculosis and Malaria: A Review. *Madonna University journal of Medicine and Health Sciences*. 2022;2(3):6-15.
40. Offie DC, Ibekwe AM, Agu CC, Esimai BN, Okpala PU, Obeagu EI, Ufelle SA, Ogbonna LN. Fibrinogen and C-Reactive Protein Significance in Children Infected by *Plasmodium falciparum* Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(15):1-8.
41. Obeagu EI, Ogunnaya FU. PREGNANCYINDUCED HAEMATOLOGICAL CHANGES: A KEY TO MATERNAL AND CHILD HEALTH. *European Journal of Biomedical*. 2023;10(8):42-43. [links/64c890bddd38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf](https://epjournals.com/links/64c890bddd38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf)
42. Obeagu EI, Ofodile AC, Okwuanaso CB. A review of urinary tract infections in pregnant women: Risks factors. *J Pub Health Nutri*. 2023; 6 (1). 2023; 137:26-35. [links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf](https://epjournals.com/links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf).
43. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. *Int. J. Curr. Res. Med. Sci*. 2023;9(2): 14-17.DOI: 10.22192/ijcrms.2023.09.02.003
44. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8): 14-21.DOI: 10.22192/ijcrps.2023.10.08.002

45. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
46. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9): 1-4.DOI: [10.22192/ijcrbm.2018.03.09.001](https://doi.org/10.22192/ijcrbm.2018.03.09.001)
47. Onyenweaku FC, Amah HC, Obeagu EI, Nwandikor UU, Onwuasoanya UF. Prevalence of asymptomatic bacteriuria and its antibiotic susceptibility pattern in pregnant women attending private ante natal clinics in Umuahia Metropolitan. *Int J Curr Res Biol Med.* 2017;2(2): 13-23.DOI: [10.22192/ijcrbm.2017.02.02.003](https://doi.org/10.22192/ijcrbm.2017.02.02.003)
48. Okoroiwu IL, Chinedu-Madu JU, Obeagu EI, Vincent CC, Ochiabuto OM, Ibekwe AM, Amaechi CO, Agu CC, Anoh NV, Amadi NM. Evaluation of Iron Status, Haemoglobin and Protein Levels of Pregnant Women in Owerri Metropolis. *Journal of Pharmaceutical Research International*. 2021;33(27A):36-43.
49. Obeagu EI, Njar VE, Obeagu GU. Infertility: Prevalence and Consequences. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(7):43-50.
50. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
51. Broen K, Brustoski K, Engelmann I, Luty AJ. Placental Plasmodium falciparum infection: causes and consequences of in utero sensitization to parasite antigens. *Molecular and biochemical parasitology*. 2007;151(1):1-8.
52. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9):41-49.
53. Ogbonna LN, Ezeoru VC, Ofodile AC, Ochiabuto OM, Obi-Ezeani CN, Okpala PU, Okafor CJ, Obeagu GU, Busari AI, Obeagu EI. Gender Based Variations of Haematological Parameters of Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(8):75-80.
54. Eberendu IF, Ozims SJ, Agu GC, Amah HC, Obasi CC, Obioma-Elemba JE, Ihekaire DE, Ibanga IE, Amah CC, Obeagu EI, Nwosu DC. Impact of human activities on the breeding of mosquitoes of human disease in Owerri metropolis, Imo state. *Int J Adv Res Biol Sci IJARBS*. 2017;4(12):98-106.
55. Obeagu EI, Ofodile AC, Okwuanaso CB. A review on socio economic and behavioral aspects of malaria and its control among children under 5 years of age in Africa. *J Pub Health Nutri.* 2023; 6 (1): 136.
56. Djontu JC, Siewe Siewe S, Mpeke Edene YD, Nana BC, Chomga Foko EV, Bigoga JD, Leke RF, Megnekou R. Impact of placental Plasmodium falciparum malaria infection on the Cameroonian maternal and neonate's plasma levels of some cytokines known to regulate T cells differentiation and function. *Malaria journal*. 2016; 15:1-1.
57. Obeagu EI, Faduma MH, Uzoma G. Ectopic Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(4): 40-4.DOI: [10.22192/ijcrps.2023.10.04.004](https://doi.org/10.22192/ijcrps.2023.10.04.004)
58. Obeagu EI, Gamade SM, Obeagu GU. The roles of Neutrophils in pregnancy. *Int. J. Curr. Res. Med. Sci.* 2023;9(5): 31-35.DOI: [10.22192/ijcrms.2023.09.05.005](https://doi.org/10.22192/ijcrms.2023.09.05.005)

Citation: Obeagu EI, Obeagu GU. Malaria and Low Birth Weight: Understanding the Linkages. *Elite Journal of Nursing and Health Science*, 2024; 2(7):41-55

59. Obeagu EI, Obeagu GU. Molar Pregnancy: Update of prevalence and risk factors. *Int. J. Curr. Res. Med. Sci.* 2023;9(7): 25-28.DOI: [10.22192/ijcrms.2023.09.07.005](https://doi.org/10.22192/ijcrms.2023.09.07.005)
60. Kabyemela E, Gonçalves BP, Prevots DR, Morrison R, Harrington W, Gwamaka M, Kurtis JD, Fried M, Duffy PE. Cytokine profiles at birth predict malaria severity during infancy. *PloS one.* 2013;8(10):e77214.
61. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Barriers to utilization of maternal health services in southern senatorial district of Cross Rivers state, Nigeria. *International Journal of Advanced Multidisciplinary Research.* 2017;4(8): 1-9.DOI: [10.22192/ijamr.2017.04.08.001](https://doi.org/10.22192/ijamr.2017.04.08.001)
62. Emmanuel G, Martin O, Peter OS, Obeagu EI, Daniel K. Factors Influencing Early Neonatal Adverse Outcomes among Women with HIV with Post Dated Pregnancies Delivering at Kampala International University Teaching Hospital, Uganda. *Asian Journal of Pregnancy and Childbirth.* 2023;6(1):203-211.
<http://research.sdpublishers.net/id/eprint/2819/>.
63. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences.* 2018;4(9): 41-9.DOI: [10.22192/ijcrms.2018.04.09.006](https://doi.org/10.22192/ijcrms.2018.04.09.006)
64. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 22-6.DOI: [10.22192/ijcrmps.2023.10.08.003](https://doi.org/10.22192/ijcrmps.2023.10.08.003)
65. Obeagu E, Eze RI, Obeagu EI, Nnatuanya IN, Dara EC. ZINC LEVEL IN APPARENTLY PREGNANT WOMEN IN URBAN AREA. *Madonna University journal of Medicine and Health Sciences.* 2022;2(1):134-48.
<https://www.journal.madonnauniversity.edu.ng/index.php/medicine/article/view/40>.
66. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(41B):380-386.
67. Obeagu EI, Ogunnaya FU, Obeagu GU, Ndidi AC. SICKLE CELL ANAEMIA: A GESTATIONAL ENIGMA. *migration.* 2023; 17:18.
68. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology.* 2019;41(3):e12573.
69. Ifeanyi OE, Uzoma OG. A review on erythropietin in pregnancy. *J. Gynecol. Womens Health.* 2018;8(3):1-4.
https://www.academia.edu/download/56538560/A_Review_on_Erythropietin_in_Pregnancy.pdf.
70. Ifeanyi OE. A review on pregnancy and haematology. *Int. J. Curr. Res. Biol. Med.* 2018;3(5): 26-8.DOI: [10.22192/ijcrbm.2018.03.05.006](https://doi.org/10.22192/ijcrbm.2018.03.05.006)
71. Nwosu DC, Nwanjo HU, Obeagu EI, Ibebuike JE, Ezeama MC. Ihekireh. Changes in liver enzymes and lipid profile of pregnant women with malaria in Owerri, Nigeria. *International Journal of Current Research and Academic Review.* 2015;3(5):376-383.
72. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Factors that influence women's utilization of

- primary health care services in Calabar Cross river state, Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2017;4(7):28-33.
73. Elemchukwu Q, Obeagu EI, Ochei KC. Prevalence of Anaemia among Pregnant Women in Braithwaite Memorial Specialist Hospital (BMSH) Port Harcourt. *IOSR Journal of Pharmacy and Biological Sciences.* 2014;9(5):59-64.
 74. Natama HM, Moncunill G, Rovira-Vallbona E, Sanz H, Sorgho H, Aguilar R, Coulibaly-Traoré M, Somé MA, Scott S, Valéa I, Mens PF. Modulation of innate immune responses at birth by prenatal malaria exposure and association with malaria risk during the first year of life. *BMC medicine.* 2018; 16:1-5.
 75. Akandinda M, Obeagu EI, Katonera MT. Non Governmental Organizations and Women's Health Empowerment in Uganda: A Review. *Asian Research Journal of Gynaecology and Obstetrics.* 2022;8(3):12-26.
 76. Gamde MS, Obeagu EI. IRON DEFICIENCY ANAEMIA: ENEMICAL TO PREGNANCY. *European Journal of Biomedical.* 2023;10(9):272-275.
[links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf](https://www.ejbs.com/links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf).
 77. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International.* 2021;33(42A):53-65.
 78. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Studies of Some Haemostatic Variables in Preeclamptic Women in Owerri, Imo State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(42B):39-48.
 79. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8):22-26.
 80. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences.* 2023;6(2):10-13.
 81. Mutabingwa TK, Bolla MC, Li JL, Domingo GJ, Li X, Fried M, Duffy PE. Maternal malaria and gravidity interact to modify infant susceptibility to malaria. *PLoS medicine.* 2005;2(12):e407.
 82. Gamble C, Ekwari PJ, Garner P, Ter Kuile FO. Insecticide-treated nets for the prevention of malaria in pregnancy: a systematic review of randomised controlled trials. *PLoS medicine.* 2007;4(3):e107.
 83. Okoko BJ, Enwere G, Ota MO. The epidemiology and consequences of maternal malaria: a review of immunological basis. *Acta tropica.* 2003;87(2):193-205.
 84. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal.* 2018; 17:1-3.
 85. Umbers AJ, Stanisic DI, Ome M, Wangnapi R, Hanieh S, Unger HW, Robinson LJ, Lufele E, Baiwog F, Siba PM, King CL. Does malaria affect placental development? Evidence from in vitro models. *PLoS One.* 2013;8(1):e55269.

86. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
87. Gbedande K, Carpio VH, Stephens R. Using two phases of the CD 4 T cell response to blood-stage murine malaria to understand regulation of systemic immunity and placental pathology in *Plasmodium falciparum* infection. *Immunological reviews*. 2020;293(1):88-114.
88. Lindsay SW, Thomas MB, Kleinschmidt I. Threats to the effectiveness of insecticide-treated bednets for malaria control: thinking beyond insecticide resistance. *The Lancet Global Health*. 2021;9(9):e1325-1331.
89. Akinleye SO, Falade CO, Ajayi IO. Knowledge and utilization of intermittent preventive treatment for malaria among pregnant women attending antenatal clinics in primary health care centers in rural southwest, Nigeria: a cross-sectional study. *BMC pregnancy and childbirth*. 2009; 9:1-9.