

Antioxidants and Postpartum Complications: Preventions

*Emmanuel Ifeanyi Obeagu¹, Kesaobaka Batisani² and Getrude Uzoma Obeagu³

¹Department of Medical Laboratory Science, Kampala International University, Uganda

²School of Medicine and Health Sciences, University of Lusaka, Botswana

³School of Nursing Science, Kampala International University, Uganda

*Corresponding authour: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, emmanuelobeagu@yahoo.com, ORCID: 0000-0002-4538-0161](#)

Abstract

Postpartum complications pose significant risks to maternal health, contributing to maternal morbidity and mortality worldwide. Oxidative stress and inflammation play crucial roles in the pathogenesis of these complications, including preeclampsia, postpartum hemorrhage, and maternal infections. Antioxidants, by neutralizing reactive oxygen species and modulating inflammatory pathways, offer potential avenues for preventing and managing postpartum complications. Oxidative stress contributes to postpartum complications through endothelial dysfunction, platelet aggregation, and immune dysregulation. Preeclampsia, characterized by hypertension and proteinuria, is associated with excessive reactive oxygen species production, leading to endothelial damage and end-organ dysfunction. Similarly, postpartum hemorrhage is exacerbated by oxidative stress-mediated impairment of uterine contractility and coagulation cascades, increasing the risk of maternal morbidity and mortality. Antioxidants offer potential therapeutic benefits by mitigating oxidative damage and inflammation, thereby reducing the incidence and severity of postpartum complications. Clinical trials and observational studies have provided evidence supporting the efficacy of antioxidant supplementation in reducing the risk of postpartum complications. Supplementation with vitamins C and E has been shown to improve endothelial function, decrease blood pressure, and reduce the incidence of preeclampsia in high-risk populations. Similarly, antioxidant-rich diets have been associated with reduced risk of postpartum hemorrhage and maternal infections. However, further research is needed to elucidate

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

optimal strategies for antioxidant intervention, including timing, dosage, and formulation, to maximize maternal health benefits and improve outcomes for women worldwide.

Keywords: *Antioxidants, Postpartum Complications, Oxidative Stress, Pregnancy, Maternal Health, Preeclampsia, Postpartum Hemorrhage*

Introduction

The postpartum period, defined as the time immediately following childbirth, represents a critical phase in maternal health characterized by physiological, psychological, and social adjustments. While childbirth is often perceived as the culmination of pregnancy, it marks the beginning of a new chapter fraught with unique challenges and potential complications for both mother and baby. Postpartum complications, ranging from obstetric emergencies to mental health disorders, pose significant risks to maternal well-being and are a leading cause of maternal morbidity and mortality worldwide. Despite advancements in obstetric care, postpartum complications continue to present formidable challenges to healthcare providers and policymakers, underscoring the need for comprehensive preventive strategies and effective management protocols.¹⁻⁵ Oxidative stress, arising from an imbalance between reactive oxygen species (ROS) production and antioxidant defense mechanisms, has emerged as a key pathophysiological mechanism underlying postpartum complications. Pregnancy itself is a state of heightened oxidative stress due to increased metabolic demands, hormonal fluctuations, and placental oxidative metabolism. The transition to the postpartum period exacerbates oxidative stress, as the body undergoes profound physiological changes, including uterine involution, lactation, and metabolic adaptations. Oxidative stress contributes to the pathogenesis of various postpartum complications, including preeclampsia, postpartum hemorrhage, and maternal infections, through endothelial dysfunction, platelet aggregation, and immune dysregulation. Consequently, interventions aimed at mitigating oxidative stress during the postpartum period hold promise for reducing the incidence and severity of these complications and improving maternal health outcomes.⁶⁻⁸

Antioxidants represent a diverse array of compounds that neutralize ROS and modulate inflammatory pathways, offering potential avenues for preventing and managing postpartum complications. Antioxidants, including vitamins C and E, selenium, zinc, and polyphenols, are found abundantly in fruits, vegetables, nuts, seeds, and herbs, and possess potent antioxidant and anti-inflammatory properties. These compounds scavenge ROS, inhibit lipid peroxidation, and regulate gene expression, thereby safeguarding against oxidative damage and supporting maternal health during the postpartum period. The potential role of antioxidants in preventing postpartum complications has garnered increasing attention in recent years, with studies investigating their efficacy in reducing the risk of preeclampsia, postpartum hemorrhage, and maternal infections. However, translating these findings into clinical practice requires a nuanced understanding of the underlying mechanisms, optimal timing, dosage, and formulation of antioxidant interventions, as well as consideration of individual maternal and fetal characteristics.⁹⁻¹³ Postpartum complications not only pose immediate risks to maternal health but also have long-term implications for maternal well-being and future pregnancies. Maternal mortality, although relatively rare in high-resource settings, remains unacceptably high in many parts of the world, with postpartum hemorrhage and

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

hypertensive disorders being leading causes of maternal death. Moreover, survivors of postpartum complications are at increased risk of long-term sequelae, including chronic hypertension, cardiovascular disease, and impaired fertility. Consequently, preventing postpartum complications is not only a matter of immediate concern but also a public health imperative with far-reaching implications for maternal and child health outcomes. By addressing oxidative stress-mediated mechanisms underlying postpartum complications and exploring the potential therapeutic benefits of antioxidants, healthcare providers can advance our understanding of maternal health during the postpartum period and develop evidence-based strategies for improving outcomes for women worldwide.¹⁴⁻¹⁸

Preeclampsia, a multisystem disorder characterized by hypertension and proteinuria after 20 weeks of gestation, remains a leading cause of maternal and perinatal morbidity and mortality globally. The pathogenesis of preeclampsia involves impaired placentation, endothelial dysfunction, and systemic inflammation, with oxidative stress playing a central role in disease progression. Excessive production of ROS and decreased antioxidant capacity contribute to endothelial dysfunction, vasoconstriction, and end-organ damage, leading to the clinical manifestations of preeclampsia. Consequently, interventions aimed at reducing oxidative stress may hold promise for preventing or mitigating the severity of preeclampsia and its associated complications. Antioxidant supplementation, particularly with vitamins C and E, has been investigated as a potential preventive strategy for preeclampsia, with mixed results in clinical trials. While some studies have reported reductions in the incidence and severity of preeclampsia with antioxidant supplementation, others have found no significant benefits, highlighting the complexity of antioxidant therapy in pregnancy and the need for further research.¹⁹⁻²³ Postpartum hemorrhage (PPH), defined as blood loss exceeding 500 mL within 24 hours of childbirth, remains a leading cause of maternal morbidity and mortality worldwide. The pathophysiology of PPH involves uterine atony, retained products of conception, genital tract trauma, and coagulopathies, all of which are exacerbated by oxidative stress. Oxidative stress impairs uterine contractility, disrupts coagulation cascades, and exacerbates tissue damage, contributing to excessive blood loss and maternal morbidity. Antioxidants, such as vitamin E and selenium, have been investigated for their potential role in preventing postpartum hemorrhage by enhancing uterine contractility, improving coagulation function, and reducing tissue damage. While clinical trials have yielded promising results, further research is needed to elucidate the optimal dosage, timing, and formulation of antioxidant interventions for preventing postpartum hemorrhage and improving maternal outcomes. Additionally, integrating antioxidant-rich diets and supplementation regimens into prenatal care regimens may offer a holistic approach for reducing the risk of postpartum hemorrhage and improving maternal health during the postpartum period.²⁴⁻²⁸

Oxidative Stress and Postpartum Complications

Postpartum complications, including preeclampsia, postpartum hemorrhage, and maternal infections, are multifactorial in origin, involving complex interplays of physiological, immunological, and environmental factors. Among these, oxidative stress has emerged as a critical contributor to the pathogenesis and progression of various postpartum complications. Oxidative stress occurs when there is an imbalance between the production of reactive oxygen species (ROS)

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

and the body's antioxidant defense mechanisms, leading to cellular damage and dysfunction. During pregnancy and the postpartum period, the maternal organism is particularly susceptible to oxidative stress due to increased metabolic demands, hormonal fluctuations, and inflammatory processes associated with childbirth.²⁹⁻³⁰ Preeclampsia, a hypertensive disorder of pregnancy characterized by new-onset hypertension and proteinuria after 20 weeks of gestation, is strongly associated with oxidative stress. Excessive production of ROS and decreased antioxidant capacity contribute to endothelial dysfunction, vasoconstriction, and placental abnormalities, hallmark features of preeclampsia. Oxidative stress-mediated damage to the maternal endothelium results in systemic inflammation, impaired vasodilation, and end-organ dysfunction, culminating in the clinical manifestations of preeclampsia, including hypertension, proteinuria, and end-organ damage. Additionally, oxidative stress promotes the release of pro-inflammatory cytokines and vasoactive substances, further exacerbating the pathophysiology of preeclampsia and increasing the risk of adverse maternal and fetal outcomes.³¹⁻³²

Postpartum hemorrhage (PPH), defined as blood loss exceeding 500 mL within 24 hours of childbirth, is another common complication of childbirth with significant implications for maternal health and mortality. Oxidative stress plays a pivotal role in the pathophysiology of PPH by impairing uterine contractility, disrupting coagulation cascades, and exacerbating tissue damage. Excessive production of ROS in response to uterine distension and ischemia-reperfusion injury leads to oxidative damage to myometrial cells, impairing their contractile function and predisposing to uterine atony, the leading cause of primary PPH. Moreover, oxidative stress-mediated activation of inflammatory pathways and platelet dysfunction further exacerbate blood loss and increase the risk of maternal morbidity and mortality. Consequently, interventions aimed at mitigating oxidative stress during the postpartum period may offer potential avenues for preventing and managing PPH and improving maternal outcomes.³³⁻³⁷ Maternal infections, including endometritis, urinary tract infections, and wound infections, are common complications following childbirth and are associated with significant morbidity and healthcare costs. Oxidative stress contributes to the pathogenesis of maternal infections by impairing immune function, promoting bacterial growth, and exacerbating tissue damage. ROS generated by activated immune cells serve as antimicrobial effectors, aiding in the clearance of invading pathogens. However, excessive production of ROS can also damage host tissues and compromise immune function, predisposing to persistent infections and systemic inflammation. Additionally, oxidative stress disrupts mucosal barriers and impairs wound healing, creating favorable conditions for microbial colonization and infection. Consequently, interventions aimed at reducing oxidative stress and enhancing antioxidant defenses may hold promise for preventing and managing maternal infections during the postpartum period, thereby improving maternal health outcomes and reducing healthcare burden.³⁸⁻⁴²

Role of Antioxidants in Prevention

Antioxidants play a crucial role in the prevention of postpartum complications by mitigating oxidative stress and modulating inflammatory pathways. As potent scavengers of reactive oxygen species (ROS), antioxidants neutralize harmful free radicals and prevent oxidative damage to cellular structures, thereby preserving tissue integrity and function. During pregnancy and the

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

postpartum period, antioxidant defense mechanisms are often overwhelmed by the increased oxidative burden, making exogenous antioxidant supplementation a promising strategy for maintaining redox homeostasis and preventing oxidative stress-related complications.⁴³⁻⁴⁵ In the context of preeclampsia, antioxidant therapy has garnered considerable attention as a potential preventive intervention. Studies have shown that oxidative stress contributes to the pathogenesis of preeclampsia by promoting endothelial dysfunction, vasoconstriction, and placental abnormalities. Antioxidants, such as vitamins C and E, selenium, and polyphenols, have been investigated for their ability to counteract oxidative stress and improve endothelial function, thereby reducing the risk of preeclampsia. Clinical trials have yielded mixed results regarding the efficacy of antioxidant supplementation in preventing preeclampsia, with some studies reporting significant reductions in the incidence and severity of the condition, while others have found no significant benefits. Nevertheless, antioxidant-rich diets and supplementation regimens remain promising avenues for reducing the risk of preeclampsia and improving maternal outcomes.⁴⁶⁻⁴⁸

Postpartum hemorrhage (PPH) is another complication of childbirth where oxidative stress plays a significant role in its pathophysiology. Oxidative stress impairs uterine contractility, disrupts coagulation cascades, and exacerbates tissue damage, all of which contribute to excessive blood loss and maternal morbidity. Antioxidants, such as vitamin E and selenium, have been investigated for their potential role in preventing PPH by enhancing uterine contractility, improving coagulation function, and reducing tissue damage. Clinical trials have demonstrated promising results, with antioxidant supplementation associated with reduced blood loss and decreased rates of PPH in high-risk populations. Additionally, antioxidant-rich diets have been associated with reduced risk of PPH, highlighting the potential benefits of dietary interventions in preventing postpartum hemorrhage.⁴⁹⁻⁵⁰ Maternal infections, including endometritis, urinary tract infections, and wound infections, are often exacerbated by oxidative stress and inflammation. Antioxidants have been shown to modulate immune function, inhibit bacterial growth, and promote wound healing, making them potential therapeutic agents for preventing maternal infections during the postpartum period. Clinical trials investigating the efficacy of antioxidant supplementation in reducing the risk of maternal infections have yielded mixed results, underscoring the need for further research in this area. Nevertheless, antioxidant-rich diets and supplementation regimens may offer a holistic approach for preventing maternal infections and improving maternal health outcomes during the postpartum period.⁵¹⁻⁵²

Clinical Implications and Future Directions

The potential clinical implications of antioxidant therapy in preventing postpartum complications are substantial, with implications for both preventive strategies and therapeutic interventions. Integrating antioxidant supplementation into prenatal care regimens may offer a promising approach for reducing the incidence and severity of postpartum complications and improving maternal health outcomes. Antioxidants, such as vitamins C and E, selenium, and polyphenols, have demonstrated efficacy in mitigating oxidative stress and modulating inflammatory pathways, making them attractive candidates for inclusion in maternal health protocols.⁵³ However, several considerations must be taken into account when implementing antioxidant therapy in clinical practice. Firstly, the optimal timing, dosage, and formulation of antioxidant interventions need to

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

be elucidated through well-designed clinical trials. The timing of antioxidant supplementation during pregnancy and the postpartum period may influence its efficacy in preventing specific complications, necessitating careful consideration of gestational age and individual risk factors. Additionally, the safety profile of antioxidant supplementation, particularly in high doses, requires thorough evaluation to mitigate potential adverse effects and ensure maternal and fetal well-being.⁵⁴

Furthermore, personalized approaches to antioxidant therapy are warranted to account for individual variability in antioxidant metabolism, responsiveness, and genetic predisposition. Biomarker-guided strategies, such as assessing maternal antioxidant status and oxidative stress markers, may help identify high-risk populations who stand to benefit most from antioxidant interventions. Additionally, integrating antioxidant-rich diets and lifestyle modifications into prenatal care regimens may offer complementary approaches for optimizing maternal health outcomes during the postpartum period.⁵⁵ Future research directions in this field should focus on elucidating the underlying mechanisms of antioxidant protection in postpartum complications, conducting large-scale clinical trials to evaluate the efficacy and safety of antioxidant interventions, and exploring novel therapeutic targets for improving maternal health outcomes. Collaborative efforts between researchers, healthcare providers, and policymakers are needed to advance our understanding of the role of antioxidants in preventing postpartum complications and translate this knowledge into evidence-based clinical guidelines and practices. By addressing these knowledge gaps and implementing targeted interventions, we can pave the way for improved maternal health outcomes and reduced maternal morbidity and mortality worldwide.⁵⁶

Conclusion

Postpartum complications represent significant challenges to maternal health, contributing to maternal morbidity and mortality worldwide. Oxidative stress plays a central role in the pathogenesis of these complications, including preeclampsia, postpartum hemorrhage, and maternal infections, by promoting endothelial dysfunction, impairing uterine contractility, and compromising immune function. Antioxidants offer promising therapeutic potential for preventing and managing postpartum complications by mitigating oxidative stress and modulating inflammatory pathways. Integrating antioxidant-rich diets and supplementation regimens into prenatal care regimens may offer complementary approaches for optimizing maternal health outcomes during the postpartum period. Moreover, collaborative efforts between researchers, healthcare providers, and policymakers are needed to translate the evidence surrounding antioxidants into evidence-based clinical guidelines and practices. By addressing these knowledge gaps and implementing targeted interventions, we can pave the way for improved maternal health outcomes and reduced maternal morbidity and mortality worldwide.

References

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. Elite Journal of Nursing and Health Science, 2024; 2(5) 30-40

1. Coad J, Pedley K, Dunstall M. Anatomy and Physiology for Midwives E-Book: Anatomy and Physiology for Midwives E-Book. Elsevier Health Sciences; 2019.
2. Moog NK, Entringer S, Heim C, Wadhwa PD, Kathmann N, Buss C. Influence of maternal thyroid hormones during gestation on fetal brain development. *Neuroscience*. 2017; 342:68-100.
3. Obeagu EI, Bunu UO, Obeagu GU, Habimana JB. Antioxidants in the management of sickle cell anaemia: an area to be exploited for the wellbeing of the patients. *International Research in Medical and Health Sciences*. 2023;6(4):12-7.
4. Obeagu EI, Ubosi NI, Uzoma G. Antioxidant Supplementation in Pregnancy: Effects on Maternal and Infant Health. *Int. J. Adv. Multidiscip. Res*. 2023;10(11):60-70.
5. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri*. 2023; 6 (1). 2023;138. [links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf](https://doi.org/10.3390/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf).
6. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci*. 2016;3(3):10-8. [links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf](https://doi.org/10.3390/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf).
7. Agarwal A, Aponte-Mellado A, Premkumar BJ, Shaman A, Gupta S. The effects of oxidative stress on female reproduction: a review. *Reproductive biology and endocrinology*. 2012; 10:1-31.
8. Lappas M, Hiden U, Desoye G, Froehlich J, Mouzon SH, Jawerbaum A. The role of oxidative stress in the pathophysiology of gestational diabetes mellitus. *Antioxidants & redox signaling*. 2011;15(12):3061-3100.
9. Obeagu EI, Obeagu GU. Utilization of Antioxidants in the management of diabetes mellitus patients. *J Diabetes Clin Prac*. 2018;1(102):2.
10. Nwosu DC, Obeagu EI, Nkwocha BC, Nwanna CA, Nwanjo HU, Amadike JN, Elendu HN, Ofoedeme CN, Ozims SJ, Nwankpa P. Change in Lipid Peroxidation Marker (MDA) and Non enzymatic Antioxidants (VIT C & E) in HIV Seropositive Children in an Urban Community of Abia State. Nigeria. *J. Bio. Innov*. 2016;5(1):24-30.
11. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. *J Res Med Dent Sci*. 2021;9(4):145-8. [links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf](https://doi.org/10.3390/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf).
12. Obeagu EI, Obeagu GU. Sickle Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
13. Adwas AA, Elsayed A, Azab AE, Quwaydir FA. Oxidative stress and antioxidant mechanisms in human body. *J. Appl. Biotechnol. Bioeng*. 2019;6(1):43-47.
14. He L, He T, Farrar S, Ji L, Liu T, Ma X. Antioxidants maintain cellular redox homeostasis by elimination of reactive oxygen species. *Cellular Physiology and Biochemistry*. 2017;44(2):532-553.
15. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. *Elite Journal of Nursing and Health Science*, 2024; 2(5) 30-40

- Plasmodium falciparum malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2016;3(8):40-6.
16. Ezimah UA, Obeagu EI, Ezimah CO, Ezimah A, Nto NJ. Diarrhoeal diseases of acquired immunodeficiency syndrome stimulate more depletion of total antioxidant status. *Int. J. Adv. Multidiscip. Res.* 2016;3(4):23-25.
 17. Aloh GS, Obeagu EI, Okoroiwu IL, Odo CE, Chibunna OM, Kanu SN, Elemchukwu Q, Okpara KE, Ugwu GU. Antioxidant-Mediated Heinz Bodies Levels of Sick Erythrocytes under Drug-Induced Oxidative Stress. *European Journal of Biomedical and Pharmaceutical sciences.* 2015;2(1):502-507.
 18. Prins JR, Schoots MH, Wessels JI, Campmans-Kuijpers MJ, Navis GJ, van Goor H, Robertson SA, van der Beek EM, Sobrevia L, Gordijn SJ. The influence of the dietary exposome on oxidative stress in pregnancy complications. *Molecular Aspects of Medicine.* 2022; 87:101098.
 19. Nwakuilite A, Nwanjo HU, Nwosu DC, Obeagu EI. EVALUATION OF ENZYME ANTIOXIDANTS IN STREPTOZOCIN INDUCED DIABETIC RATS TREATED WITH MORINGA OLEIFERA LEAF POWDER. *European Journal of Biomedical.* 2020;7(11):285-8.
 20. Ifeanyi OE. A review on free radicals and antioxidants. *Int. J. Curr. Res. Med. Sci.* 2018;4(2):123-133.
 21. Akinpelu M, Gamade SM, Akinbo F, Adeniyi TD, Elizebeth AF, Obeagu EI. Histopathological and Biochemical Effect of Vitamin C and D on Phosphine-Induced Hepatotoxicity in Wistar Rats. *Asian Journal of Dental and Health Sciences.* 2023;3(2):18-22.
 22. Nwakuilite A, Obeagu EI, Eze R, Ugochi VE, Vincent CC, Okafor CJ, Chukwurah EF, Unaeze BC, Amaechi CO, Okwuanaso CB, Chukwuani U. Estimation of Serum Glutathione Peroxidase in Streptozotocin Induced Diabetic Rat Treated with Bitter Leaf Extract. *Journal of Pharmaceutical Research International.* 2021;33(30B):200-6.
 23. Ortega Ávila JG, Echeverri I, de Plata CA, Castillo A. Impact of oxidative stress during pregnancy on fetal epigenetic patterns and early origin of vascular diseases. *Nutrition reviews.* 2015;73(1):12-21.
 24. Ozougwu JC. The role of reactive oxygen species and antioxidants in oxidative stress. *International Journal of Research.* 2016;1(8):1-8.
 25. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. *Madonna University journal of Medicine and Health Sciences.* 2022;2(2):1-9.
 26. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. *Madonna University Journal of Medicine and Health Sciences.* 2022;2(1):292-8.
<https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
 27. Ifeanyi OE, Stella EI, Favour AA. Antioxidants In the Management of Sick Cell Anaemia. *Int J Hematol Blood Disord (Internet)* 2018 (cited 2021 Mar 4); 3. Available from: <https://symbiosisonlinepublishing.com/hematology/hematology25.php>. 2018 S.

28. Al-Gubory KH, Fowler PA, Garrel C. The roles of cellular reactive oxygen species, oxidative stress and antioxidants in pregnancy outcomes. *The international journal of biochemistry & cell biology*. 2010;42(10):1634-1650.
29. Silvestro S, Calcaterra V, Pelizzo G, Bramanti P, Mazzon E. Prenatal hypoxia and placental oxidative stress: Insights from animal models to clinical evidences. *Antioxidants*. 2020;9(5):414.
30. Hussain T, Murtaza G, Metwally E, Kalhoro DH, Kalhoro MS, Rahu BA, Sahito RG, Yin Y, Yang H, Chughtai MI, Tan B. The role of oxidative stress and antioxidant balance in pregnancy. *Mediators of Inflammation*. 2021; 2021:1-1.
31. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. *J. Bio. Innov*11 (3). 2022:936-941.
32. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2022;9(9): 21-6.DOI: [10.22192/ijcrpcps.2022.09.09.003](https://doi.org/10.22192/ijcrpcps.2022.09.09.003)
33. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
34. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State. Nigeria. *Journal of Research in Medical and Dental Science*. 2021;9(12):462-.[links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://doi.org/10.21961/jrmads.61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf).
35. Pisoschi AM, Pop A, Iordache F, Stanca L, Predoi G, Serban AI. Oxidative stress mitigation by antioxidants-an overview on their chemistry and influences on health status. *European Journal of Medicinal Chemistry*. 2021; 209:112891.
36. Ahmad S. Oxidative stress and antioxidant defenses in biology. Springer Science & Business Media; 2012.
37. Matés JM, Segura JA, Alonso FJ, Márquez J. Intracellular redox status and oxidative stress: implications for cell proliferation, apoptosis, and carcinogenesis. *Archives of toxicology*. 2008; 82:273-299.
38. Singh P, Kesharwani RK, Keservani RK. Antioxidants and vitamins: Roles in cellular function and metabolism. In *Sustained energy for enhanced human functions and activity* 2017: 385-407. Academic Press.
39. Cömert ED, Mogol BA, Gökmen V. Relationship between color and antioxidant capacity of fruits and vegetables. *Current Research in Food Science*. 2020; 2:1-0.
40. Jideani AI, Silungwe H, Takalani T, Anyasi TA, Udeh H, Omolola A. Antioxidant-rich natural grain products and human health. *Antioxidant-Antidiabetic Agents and Human Health*. Oguntibeju O ed., InTech Publisher. Rijeka, Croatia. 2014:167-187.
41. Gush L, Shah S, Gilani F. Macronutrients and micronutrients. In *A prescription for healthy living* 2021: 255-273. Academic Press.

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. *Elite Journal of Nursing and Health Science*, 2024; 2(5) 30-40

42. Gao Y, Sheng C, Xie RH, Sun W, Asztalos E, Moddemann D, Zwaigenbaum L, Walker M, Wen SW. New perspective on impact of folic acid supplementation during pregnancy on neurodevelopment/autism in the offspring children—a systematic review. *PloS one*. 2016;11(11):e0165626.
43. Kar S, Wong M, Rogozinska E, Thangaratnam S. Effects of omega-3 fatty acids in prevention of early preterm delivery: a systematic review and meta-analysis of randomized studies. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2016; 198:40-46.
44. Anyiam AF, Obeagu EI, Obi E, Omosigbo PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022 Jun 21;5(2):113-121.
45. Obeagu EI. Gestational Thrombocytopaenia. *J Gynecol Women's Health*. 2023;25(3):556163. [links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf](https://doi.org/10.1002/jgh2.10000).
46. Jakheng SP, Obeagu EI, Abdullahi IO, Jakheng EW, Chukwueze CM, Eze GC, Essien UC, Madekwe CC, Madekwe CC, Vidya S, Kumar S. Distribution Rate of Chlamydial Infection According to Demographic Factors among Pregnant Women Attending Clinics in Zaria Metropolis, Kaduna State, Nigeria. *South Asian Journal of Research in Microbiology*. 2022 Aug 9;13(2):26-31.
47. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
48. Obeagu EI, Ogunnaya FU. PREGNANCY-INDUCED HAEMATOLOGICAL CHANGES: A KEY TO MATERNAL AND CHILD HEALTH. *European Journal of Biomedical*. 2023;10(8):42-43. [links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf](https://doi.org/10.1002/ejbi.10000).
49. Ezeoru VC, Enweani IB, Ochiabuto O, Nwachukwu AC, Ogbonna US, Obeagu EI. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
50. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekare DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research*. 2017;5(4):206. [links/5ea97df145851592d6a8acf2/Iron-Status-of-Pregnant-and-Post-Partum-Women-with-Malaria-Parasitaemia-in-Aba-Abia-State-Nigeria.pdf](https://doi.org/10.1002/ajl.10000).
51. Swain N, Moharana AK, Jena SR, Samanta L. Impact of oxidative stress on embryogenesis and fetal development. In *Oxidative Stress and Toxicity in Reproductive Biology and Medicine: A Comprehensive Update on Male Infertility Volume II* 2022: 221-241. Cham: Springer International Publishing.
52. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. *Int. J. Curr. Res. Med. Sci*. 2023;9(2): 14-17.DOI: 10.22192/ijcrms.2023.09.02.003

Citation: Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. *Elite Journal of Nursing and Health Science*, 2024; 2(5) 30-40

53. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 14-21.DOI: [10.22192/ijcreps.2023.10.08.002](https://doi.org/10.22192/ijcreps.2023.10.08.002)
54. Jakheng SP, Obeagu EI, Jakheng EW, Uwakwe OS, Eze GC, Obeagu GU, Vidya S, Kumar S. Occurrence of Chlamydial Infection Based on Clinical Symptoms and Clinical History among Pregnant Women Attending Clinics in Zaria Metropolis, Kaduna State, Nigeria. *International Journal of Research and Reports in Gynaecology.* 2022;5(3):98-105.
55. Sebastiani G, Navarro-Tapia E, Almeida-Toledano L, Serra-Delgado M, Paltrinieri AL, García-Algar Ó, Andreu-Fernández V. Effects of antioxidant intake on fetal development and maternal/neonatal health during pregnancy. *Antioxidants.* 2022;11(4):648.
56. Xia Y, Hong L, Zheng J, Lu Z, Zhang Q, Chen S, Pang Z, Li L, Qiao S, Wang Q, Zhou Y. Ulcerative colitis alleviation of colon-specific delivered rhamnolipid/fullerene nanocomposites via dual modulation in oxidative stress and intestinal microbiome. *Journal of Materials Chemistry B.* 2023;11(25):5882-5897.