Elite Journal of Laboratory Medicine. Volume 2 issue 5(2024), Pp. 24-31 https://epjournals.com/journals/EJLM

Understanding Transfusion-Related Changes in Coagulation Parameters and Disseminated Intravascular Coagulation in Pediatric Severe Malaria Cases with HIV: A Comprehensive Review

*Emmanuel Ifeanyi Obeagu

Department of Medical Laboratory Science, Kampala International University, Uganda

*Corresponding authour: Emmanuel Ifeanyi Obeagu, <u>Department of Medical Laboratory Science</u>, <u>Kampala International University, Uganda, emmanuelobeagu@yahoo.com, ORCID:</u> 0000-0002-4538-0161

Abstract

Pediatric severe malaria cases complicated by HIV co-infection present a significant clinical challenge, characterized by a complex interplay of severe anemia, coagulation abnormalities, and an increased risk of disseminated intravascular coagulation (DIC). Blood transfusion, a cornerstone intervention in managing severe anemia, introduces additional complexities by potentially altering coagulation parameters and predisposing patients to transfusion-related complications. Despite the clinical significance, the impact of transfusion on coagulation dynamics and DIC development in pediatric severe malaria cases with HIV remains poorly understood. This review aims to address this knowledge gap by critically evaluating current evidence and providing insights into tailored management strategies to optimize outcomes in this vulnerable population. Alterations in coagulation parameters, including thrombocytopenia, prolonged PT and aPTT, and elevated FDP levels, are frequently observed in pediatric severe malaria cases, reflecting underlying coagulation dysregulation. The presence of HIV infection further exacerbates coagulation abnormalities, potentially predisposing pediatric patients to DIC, a life-threatening complication associated with increased morbidity and mortality. Transfusion therapy, while essential for correcting severe anemia and improving tissue oxygenation, may exacerbate coagulation abnormalities and contribute to DIC development through various mechanisms, including dilution of coagulation factors and platelets, immunomodulatory effects, and proinflammatory cytokine release.

Keywords: Transfusion, Coagulation Parameters, Disseminated Intravascular Coagulation, Pediatric, Severe Malaria, HIV

Introduction

Pediatric severe malaria cases complicated by HIV infection represent a formidable challenge in clinical practice, particularly in regions with high malaria endemicity and HIV prevalence. Severe malaria is characterized by a spectrum of complications, including severe anemia, cerebral malaria, respiratory distress, and metabolic derangements, contributing significantly to morbidity and mortality in pediatric populations. The coexistence of HIV infection further exacerbates the clinical complexity, as HIV-associated immune dysfunction and hematologic alterations may predispose pediatric patients to more severe manifestations of malaria, including coagulation abnormalities and disseminated intravascular coagulation (DIC). Thus, understanding the interplay between severe malaria, HIV, and transfusion-related changes in coagulation parameters is imperative for optimizing clinical management and improving outcomes in this vulnerable population. 1-5 Pediatric severe malaria cases often present with alterations in coagulation parameters, reflecting the dysregulation of both procoagulant and anticoagulant pathways. Thrombocytopenia, prolonged prothrombin time (PT), activated partial thromboplastin time (aPTT), and elevated levels of fibrin degradation products (FDPs) are commonly observed in severe malaria, indicative of ongoing coagulopathy and endothelial activation. The presence of HIV infection further exacerbates coagulation abnormalities, as HIV-associated immune dysfunction and endothelial injury may potentiate coagulation activation and predispose pediatric patients to DIC. Consequently, pediatric severe malaria cases with HIV co-infection represent a high-risk group for coagulation-related complications, highlighting the need for tailored management strategies to address these complex pathophysiologic processes. 6-10

Transfusion therapy plays a crucial role in managing severe anemia and improving tissue oxygenation in pediatric severe malaria cases, but its impact on coagulation parameters and DIC development in the context of HIV co-infection remains poorly understood. Blood transfusion introduces exogenous erythrocytes, plasma, and coagulation factors into the recipient's circulation, potentially altering coagulation dynamics and predisposing patients to transfusion-related complications, including DIC. Moreover, the immunomodulatory effects of transfusion and the release of pro-inflammatory cytokines may further exacerbate coagulation activation and endothelial injury, complicating the management of coagulation abnormalities in pediatric patients with severe malaria and HIV co-infection. 11-15 The development of DIC represents a critical complication in pediatric severe malaria cases with HIV co-infection, as it is associated with increased morbidity and mortality. DIC is characterized by systemic activation of the coagulation cascade, leading to widespread microvascular thrombosis, consumption of coagulation factors and platelets, and ultimately, organ dysfunction. Pediatric patients with severe malaria and HIV coinfection may be particularly susceptible to DIC due to the complex interplay between malariainduced coagulopathy, HIV-associated immune dysfunction, and transfusion-related changes in coagulation parameters. Thus, early recognition and management of DIC are paramount in optimizing clinical outcomes and reducing mortality in this vulnerable population. ¹⁶⁻¹⁸ Despite the significant clinical implications, there is a paucity of data regarding transfusion-related changes in coagulation parameters and DIC development in pediatric severe malaria cases with HIV coinfection. Existing literature predominantly focuses on adult populations, with limited studies Citation: Obeagu EI. Understanding Transfusion-Related Changes in Coagulation Parameters and Disseminated Intravascular Coagulation in Pediatric Severe Malaria Cases with HIV: A Review. Elite Journal of Laboratory Medicine, 2024; 2(5): 24-31

investigating the specific challenges and management strategies relevant to pediatric patients. Therefore, this review aims to bridge this gap in knowledge by comprehensively evaluating current evidence and providing insights into tailored management strategies for pediatric severe malaria cases with HIV co-infection. By elucidating the complex interplay between severe malaria, HIV, transfusion-related changes in coagulation parameters, and DIC development, this review seeks to inform evidence-based clinical practice and improve outcomes in this vulnerable patient population. ¹⁹⁻²⁰

Coagulation Parameters in Pediatric Severe Malaria Cases with HIV

In pediatric severe malaria cases with HIV co-infection, alterations in coagulation parameters represent a significant aspect of the clinical presentation and management. Severe malaria often induces a dysregulated coagulation system, characterized by thrombocytopenia, prolonged prothrombin time (PT), activated partial thromboplastin time (aPTT), and elevated levels of fibrin degradation products (FDPs). These changes reflect the activation of both procoagulant and anticoagulant pathways, contributing to the pathogenesis of coagulopathy. In the context of HIV co-infection, the presence of immunosuppression and endothelial dysfunction further exacerbates coagulation abnormalities, increasing the propensity for thrombotic complications and disseminated intravascular coagulation (DIC). Therefore, monitoring coagulation parameters is essential in the management of pediatric severe malaria cases with HIV, guiding treatment strategies and mitigating the risk of thromboembolic events. ²¹⁻²⁵ Thrombocytopenia, characterized by a reduction in platelet count, is a common finding in pediatric severe malaria cases with HIV co-infection and is indicative of platelet sequestration, consumption, and destruction. The reduction in platelet count contributes to the hemorrhagic diathesis observed in severe malaria and increases the risk of bleeding complications, particularly in the setting of coexisting thrombocytopenia due to HIV-associated immune dysfunction. Moreover, thrombocytopenia may serve as a prognostic indicator of disease severity and mortality in pediatric patients with severe malaria and HIV co-infection, necessitating close monitoring and appropriate management to prevent hemorrhagic complications. 26-28 Prolonged PT and aPTT, along with elevated levels of FDPs, reflect the activation of the coagulation cascade and fibrinolytic system in pediatric severe malaria cases with HIV co-infection. These abnormalities indicate a state of hypercoagulability and ongoing fibrinolysis, which may contribute to microvascular thrombosis, organ dysfunction, and DIC. Additionally, the presence of HIV infection exacerbates endothelial dysfunction and systemic inflammation, further potentiating coagulation activation and thrombotic complications. Therefore, monitoring coagulation parameters, along with clinical assessment, is essential in identifying patients at risk of DIC and guiding therapeutic interventions, including anticoagulation therapy and supportive care, to mitigate the risk of thromboembolic events and improve outcomes in this vulnerable population.²⁹⁻³⁰

Transfusion-Related Changes in Coagulation Parameters

Transfusion-related changes in coagulation parameters represent a critical aspect of clinical management in pediatric severe malaria cases with HIV co-infection, particularly given the **Citation**: Obeagu EI. Understanding Transfusion-Related Changes in Coagulation Parameters and Disseminated Intravascular Coagulation in Pediatric Severe Malaria Cases with HIV: A Review. Elite Journal of Laboratory Medicine, 2024; 2(5): 24-31

frequent need for blood transfusions to manage severe anemia. Blood transfusion introduces exogenous erythrocytes, plasma, and coagulation factors into the recipient's circulation, potentially altering coagulation dynamics and predisposing patients to transfusion-related complications. While transfusion therapy aims to improve tissue oxygenation and prevent mortality, its impact on coagulation parameters must be carefully monitored to mitigate the risk of thrombotic and hemorrhagic complications. Transfusion-related changes in coagulation parameters may include dilutional effects, alterations in coagulation factor levels, and immune-mediated reactions, all of which can influence hemostasis and contribute to coagulopathy in pediatric patients with severe malaria and HIV co-infection. 31-35

Dilutional effects are a primary mechanism underlying transfusion-related changes in coagulation parameters, particularly in pediatric patients receiving large-volume transfusions. Transfusion of packed red blood cells (PRBCs) may lead to dilution of coagulation factors and platelets, resulting in a relative deficiency of clotting factors and impaired hemostasis. Additionally, dilutional thrombocytopenia may exacerbate pre-existing thrombocytopenia in severe malaria cases, increasing the risk of bleeding complications. Therefore, close monitoring of coagulation parameters, including PT, aPTT, and platelet count, is essential in pediatric patients receiving blood transfusions to detect dilutional effects and guide transfusion strategies to mitigate the risk of coagulopathy. 36-37 In addition to dilutional effects, transfusion-related changes in coagulation parameters may also involve alterations in coagulation factor levels and immune-mediated reactions. Transfusion of plasma-containing blood products, such as fresh frozen plasma (FFP) or cryoprecipitate, may introduce exogenous coagulation factors into the recipient's circulation, potentially correcting coagulation abnormalities and improving hemostasis. However, transfusionrelated immune reactions, such as transfusion-related acute lung injury (TRALI) or transfusionassociated circulatory overload (TACO), can lead to systemic inflammation, endothelial dysfunction, and further exacerbation of coagulation abnormalities. Therefore, careful selection of blood products, along with vigilant monitoring for transfusion-related complications, is essential in optimizing transfusion therapy and minimizing the risk of adverse outcomes in pediatric severe malaria cases with HIV co-infection. 38-40

Implications for DIC Development and Management

The implications of transfusion-related changes in coagulation parameters extend to the development and management of disseminated intravascular coagulation (DIC) in pediatric severe malaria cases with HIV co-infection. DIC represents a critical complication characterized by systemic activation of the coagulation cascade, leading to widespread microvascular thrombosis, consumption of coagulation factors and platelets, and ultimately, organ dysfunction. Transfusion therapy, while essential for correcting severe anemia, may exacerbate DIC development through various mechanisms, including dilutional effects, alterations in coagulation factor levels, and immune-mediated reactions. Transfusion-related dilutional effects can contribute to DIC development by impairing hemostasis and exacerbating pre-existing coagulopathy in pediatric severe malaria cases with HIV co-infection. The transfusion of packed red blood cells (PRBCs) may lead to dilution of coagulation factors and platelets, resulting in a relative deficiency of Citation: Obeagu EI. Understanding Transfusion-Related Changes in Coagulation Parameters and Disseminated Intravascular Coagulation in Pediatric Severe Malaria Cases with HIV: A Review. Elite Journal of Laboratory Medicine, 2024; 2(5): 24-31

clotting factors and increased bleeding tendency. Furthermore, the introduction of plasmacontaining blood products, such as fresh frozen plasma (FFP) or cryoprecipitate, may alter the balance between procoagulant and anticoagulant factors, promoting coagulation activation and thrombotic complications. Therefore, careful monitoring of coagulation parameters and prompt recognition of DIC are essential in guiding transfusion practices and optimizing outcomes in this vulnerable population. 43-44 Management of DIC in pediatric severe malaria cases with HIV coinfection requires a multidisciplinary approach involving pediatricians, hematologists, infectious disease specialists, and transfusion medicine experts. Early recognition and treatment of DIC are paramount in preventing progression to multiorgan failure and reducing mortality. Therapeutic interventions may include supportive care, such as volume resuscitation and vasopressor therapy, along with targeted treatments to address the underlying coagulopathy, such as anticoagulation therapy and transfusion of blood products. However, the management of DIC in pediatric patients with severe malaria and HIV co-infection presents unique challenges, given the complex interplay between disease pathophysiology, immune status, and transfusion-related complications. Therefore, further research is needed to elucidate optimal management strategies and improve outcomes in this vulnerable population. 45-47

Conclusion

The management of coagulation parameters and disseminated intravascular coagulation (DIC) in pediatric severe malaria cases with HIV co-infection represents a complex clinical challenge. Transfusion therapy, while essential for correcting severe anemia, introduces additional complexities by potentially altering coagulation dynamics and predisposing patients to DIC development. Dilutional effects, alterations in coagulation factor levels, and immune-mediated reactions associated with blood transfusion can exacerbate pre-existing coagulopathy and increase the risk of thrombotic and hemorrhagic complications in this vulnerable population. Early recognition and management of DIC are paramount in optimizing outcomes in pediatric severe malaria cases with HIV co-infection. Close monitoring of coagulation parameters, along with vigilant assessment of clinical status, is essential in identifying patients at risk of DIC and guiding therapeutic interventions. A multidisciplinary approach involving pediatricians, hematologists, infectious disease specialists, and transfusion medicine experts is essential in developing comprehensive management strategies tailored to the individual needs of each patient.

References

- 1. Manning L, Laman M, Davis WA, Davis TM. Clinical features and outcome in children with severe Plasmodium falciparum malaria: a meta-analysis. PLoS One. 2014;9(2):e86737.
- 2. Patel H, Dunican C, Cunnington AJ. Predictors of outcome in childhood Plasmodium falciparum malaria. Virulence. 2020;11(1):199-221.
- 3. Obeagu EI, Obeagu GU. Transfusion-Related Complications in Children Under 5 with Coexisting HIV and Severe Malaria: A Review. Int. J. Curr. Res. Chem. Pharm. Sci. 2024;11(2):9-19.

- 4. Obeagu EI, Obeagu GU. Counting Cells, Shaping Fates: CD4/CD8 Ratios in HIV. Elite Journal of Scientific Research and Review. 2024;2(1):37-50.
- 5. Obeagu EI, Obeagu GU. Eosinophil Dynamics in Pregnancy among Women Living with HIV: A Comprehensive Review. Int. J. Curr. Res. Med. Sci. 2024;10(1):11-24.
- 6. Balaji SN, Deshmukh R, Trivedi V. Severe malaria: Biology, clinical manifestation, pathogenesis and consequences. Journal of Vector Borne Diseases. 2020;57(1):1-3.
- 7. White NJ. Anaemia and malaria. Malaria journal. 2018;17(1):371.
- 8. Perkins DJ, Were T, Davenport GC, Kempaiah P, Hittner JB, Ong'echa JM. Severe malarial anemia: innate immunity and pathogenesis. International journal of biological sciences. 2011;7(9):1427.
- 9. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035. 2022;2(2):1-9.
- 10. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfected with HIV in Umuahia. Int. J. Curr. Res. Med. Sci. 2017;3(5):100-104.
- 11. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. Medicine. 2023;102(46): e36166.
- 12. Obeagu EI, Obeagu GU, Paul-Chima UO. Stigma Associated With HIV. AIDS: A Review. Newport International Journal of Public Health and Pharmacy (Nijpp). 2023;3(2):64-67.
- 13. Viola N, Kimono E, Nuruh N, Obeagu EI. Factors Hindering Elimination of Mother to Child Transmission of HIV Service Uptake among HIV Positive Women at Comboni Hospital Kyamuhunga Bushenyi District. Asian J Dental Health Sci [Internet]. 2023;3(2):7-14. http://ajdhs.com/index.php/journal/article/view/39
- 14. Tzounakas VL, Valsami SI, Kriebardis AG, Papassideri IS, Seghatchian J, Antonelou MH. Red cell transfusion in paediatric patients with thalassaemia and sickle cell disease: current status, challenges and perspectives. Transfusion and apheresis science. 2018 Jun 1:57(3):347-57.
- 15. Singh U. Hematological Diseases and Syndromes in the Neonate: Haemoglobin, Haemoglobinopathies, and Oxygen Therapy. InClinical Anesthesia for the Newborn and the Neonate 2023; 291-313. Singapore: Springer Nature Singapore.
- 16. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. Elite Journal of Laboratory Medicine, 2024; 2(1): 33-45
- 17. Obeagu EI, Obeagu GU. The Role of Blood Transfusion Strategies in HIV Management: Current Insights and Future Directions. Elite Journal of Medicine, 2024; 2(1):10-22
- 18. Obeagu EI, Okwuanaso CB, Edoho SH, Obeagu GU. Under-nutrition among HIV-exposed Uninfected Children: A Review of African Perspective. Madonna University journal of Medicine and Health Sciences. 2022;2(3):120-127.
- 19. Obeagu EI. A Review of Challenges and Coping Strategies Faced by HIV/AIDS Discordant Couples. Madonna University journal of Medicine and Health Sciences. 2023

;3(1):7-12. https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/91.

- 20. Obeagu EI, Obeagu GU. An update on premalignant cervical lesions and cervical cancer screening services among HIV positive women. J Pub Health Nutri. 2023; 6 (2). 2023; 141:1-2. links/63e538ed64252375639dd0df/An-update-on-premalignant-cervical-lesions-and-cervical-cancer-screening-services-among-HIV-positive-women.pdf.
- 21. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. Elite Journal of Laboratory Medicine. 2024;2(1):33-45.
- 22. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. Elite Journal of Laboratory Medicine. 2024;2(1):33-45.
- 23. Obeagu EI, Obeagu GU. The Role of Blood Transfusion Strategies in HIV Management: Current Insights and Future Directions. Elite Journal of Medicine. 2024;2(1):10-22.
- 24. Obeagu EI, Obeagu GU. Advances in Understanding the Impact of Blood Transfusion on Anemia Resolution in HIV-Positive Children with Severe Malaria: A Comprehensive Review. Elite Journal of Haematology. 2024;2(1):26-41.
- 25. Obeagu EI, Babar Q, Obeagu GU. Allergic blood Transfusion reaction: A Review. Int. J. Curr. Res. Med. Sci. 2021;7(5):25-33.
- 26. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. Medicine. 2023;102(46): e36166.
- 27. Obeagu EI, Obeagu GU, Ukibe NR, Oyebadejo SA. Anemia, iron, and HIV: decoding the interconnected pathways: A review. Medicine. 2024;103(2): e36937.
- 28. Obeagu EI, Obeagu GU. Tackling childhood anemia in malaria zones: comprehensive strategies, challenges, and future directions. Academia Medicine. 2024; 1:1-6.
- 29. Nhampossa T, Sigaúque B, Machevo S, Macete E, Alonso P, Bassat Q, Menéndez C, Fumadó V. Severe malnutrition among children under the age of 5 years admitted to a rural district hospital in southern Mozambique. Public health nutrition. 2013;16(9):1565-1574.
- 30. Wasserheit JN, Holmes KK. Reproductive tract infections: challenges for international health policy, programs, and research. InReproductive tract infections: Global impact and priorities for women's reproductive health 1992; 7-33. Boston, MA: Springer US.
- 31. Obeagu EI, Obeagu GU. Hematological Changes Following Blood Transfusion in Young Children with Severe Malaria and HIV: A Critical Review. Elite Journal of Laboratory Medicine. 2024;2(1):33-45.
- 32. Obeagu EI, Obeagu GU. Advances in Understanding the Impact of Blood Transfusion on Anemia Resolution in HIV-Positive Children with Severe Malaria: A Comprehensive Review. Elite Journal of Haematology. 2024;2(1):26-41.
- 33. Obeagu EI, Obeagu GU. Synergistic Effects of Blood Transfusion and HIV in Children Under 5 Years with Severe Malaria: A Review. Elite Journal of HIV. 2024;2(1):31-50.
- 34. Obeagu EI, Obeagu GU. Hematologic Considerations in Breast Cancer Patients with HIV: Insights into Blood Transfusion Strategies. Elite Journal of Health Science. 2024;2(2):20-35.

- 35. Obeagu EI, Obeagu GU. The Nexus Between Obesity and Leukemia Progression in HIV-Positive Individuals: A Review. Elite Journal of Haematology. 2024;2(4):180-98.
- 36. Akoolo L, Rocha SC, Parveen N. Protozoan co-infections and parasite influence on the efficacy of vaccines against bacterial and viral pathogens. Frontiers in Microbiology. 2022; 13:1020029.
- 37. Obeagu EI, Obeagu GU. Anemia and Malaria: Twin Troubles in Pediatric Health. Elite Journal of Health Science. 2024;2(3):36-43.
- 38. Wintergerst ES, Maggini S, Hornig DH. Contribution of selected vitamins and trace elements to immune function. Annals of nutrition and metabolism. 2007;51(4):301-323.
- 39. Haldar K, Mohandas N. Malaria, erythrocytic infection, and anemia. ASH Education Program Book. 2009;2009(1):87-93.
- 40. Tsai AG, Hofmann A, Cabrales P, Intaglietta M. Perfusion vs. oxygen delivery in transfusion with "fresh" and "old" red blood cells: the experimental evidence. Transfusion and apheresis science. 2010;43(1):69-78.
- 41. Jelkmann W. Erythropoietin: structure, control of production, and function. Physiological reviews. 1992;72(2):449-489.
- 42. Ezeoru VC, Enweani IB, Ochiabuto O, Nwachukwu AC, Ogbonna US, Obeagu EI. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. Journal of Pharmaceutical Research International. 2021;33(4):10-19.
- 43. Omo-Emmanuel UK, Chinedum OK, Obeagu EI. Evaluation of laboratory logistics management information system in HIV/AIDS comprehensive health facilities in Bayelsa State, Nigeria. Int J Curr Res Med Sci. 2017;3(1): 21-38.DOI: 10.22192/ijcrms.2017.03.01.004
- 44. Obeagu EI, Obeagu GU. An update on survival of people living with HIV in Nigeria. J Pub Health Nutri. 2022; 5 (6). 2022;129. links/645b4bfcf3512f1cc5885784/An-update-on-survival-of-people-living-with-HIV-in-Nigeria.pdf.
- 45. Obeagu EI, Obeagu GU. Advances in Understanding the Impact of Blood Transfusion on Anemia Resolution in HIV-Positive Children with Severe Malaria: A Comprehensive Review. Elite Journal of Haematology. 2024;2(1):26-41.
- 46. Pradesh A. 4th ISTM Annual Conference, TRANSMEDCON 2015, Kolkata. Asian Journal of Transfusion Science. 2016; 10:1.
- 47. Obeagu EI. Understanding the Impact of Blood Transfusion on Erythropoiesis in Pediatric Severe Malaria Cases with HIV: A Review. *Elite Journal of Haematology, 2024; 2(5):* 147-154