

## Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review

\*Emmanuel Ifeanyi Obeagu<sup>1</sup> and Getrude Uzoma Obeagu<sup>2</sup>

<sup>1</sup>Department of Medical Laboratory Science, Kampala International University, Ishaka, Uganda.

<sup>2</sup>School of Nursing Science, Kampala International University, Ishaka, Uganda.

\*Corresponding author: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, \[emmanuelobeagu@yahoo.com\]\(mailto:emmanuelobeagu@yahoo.com\), ORCID: 0000-0002-4538-0161](#)

### Abstract

Malaria, anemia, and iron deficiency during pregnancy represent a critical triad of health challenges with profound implications for both maternal and fetal well-being. Malaria, caused by *PLASMODIUM* species, exacerbates anemia and interacts with iron deficiency, creating a complex clinical scenario that impacts pregnancy outcomes. This integrated review explores the interplay between these conditions, focusing on their combined effects on maternal health, fetal development, and overall pregnancy outcomes. The review highlights the pathophysiological mechanisms through which malaria-induced inflammation and iron deficiency contribute to anemia, and how these factors compound the risks of low birth weight, preterm birth, and other adverse outcomes. Epidemiological evidence underscores the high prevalence of malaria, anemia, and iron deficiency in pregnant women, particularly in malaria-endemic regions. Studies reveal that these conditions often coexist and interact, leading to increased maternal morbidity and adverse fetal outcomes. Effective public health interventions, including intermittent preventive treatment in pregnancy (IPTp), insecticide-treated bed nets (ITNs), and routine iron supplementation, are crucial for mitigating the impact of these conditions. Integrating malaria and anemia management within maternal and child health services is essential for improving health outcomes and reducing the burden of these prevalent issues.

**Keywords:** *malaria, anemia, iron deficiency, pregnancy, maternal health, fetal development, public health interventions*

### Introduction

Pregnancy represents a unique physiological state during which maternal health is intricately linked to fetal development. Among the myriads of health concerns that can impact pregnant women, malaria, anemia, and iron deficiency stand out as major contributors to adverse pregnancy outcomes. These conditions are particularly prevalent in low-resource settings, where the burden of infectious diseases and nutritional deficiencies is high. Understanding the interactions between

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

malaria, anemia, and iron deficiency is crucial for developing effective interventions and improving maternal and fetal health. Malaria, caused by *PLASMODIUM* parasites transmitted through *Anopheles* mosquitoes, is a significant public health issue in many tropical and subtropical regions. In pregnant women, malaria can lead to severe complications, including anemia, preterm birth, and low birth weight. The presence of malaria during pregnancy often exacerbates anemia, creating a complex interplay that affects both maternal and fetal health. The inflammatory response induced by malaria can interfere with fetal development and lead to poor pregnancy outcomes. Anemia, defined as a reduction in hemoglobin concentration below the normal range, is a prevalent condition during pregnancy. It can result from multiple causes, including iron deficiency, folate deficiency, and chronic diseases. Iron deficiency anemia (IDA) is the most common type, resulting from insufficient iron to meet the increased demands of pregnancy. Anemia during pregnancy is associated with increased maternal morbidity and mortality, as well as adverse outcomes such as low birth weight and impaired fetal development. Iron is a crucial nutrient for various physiological functions, including oxygen transport and cellular metabolism. During pregnancy, the demand for iron increases to support the growing fetus and expanded maternal blood volume. Insufficient iron intake or absorption can lead to iron deficiency, which in turn can contribute to anemia. Iron deficiency during pregnancy can impair cognitive development in the fetus, increase the risk of preterm birth, and negatively impact maternal health.<sup>1-10</sup>

The interplay between malaria, anemia, and iron deficiency creates a complex health scenario. Malaria-induced inflammation can exacerbate anemia by increasing red blood cell destruction and impairing iron metabolism. Conversely, iron deficiency can compromise the body's ability to combat malaria effectively, leading to more severe infections and further worsening anemia. This cyclical relationship underscores the need for integrated approaches to manage these interconnected conditions. Epidemiological studies reveal that malaria, anemia, and iron deficiency often co-occur, particularly in malaria-endemic regions. For example, research indicates high prevalence rates of these conditions in sub-Saharan Africa, where malaria transmission is intense and access to healthcare services may be limited. The coexistence of malaria and anemia complicates the clinical management of pregnant women, requiring tailored interventions to address both conditions simultaneously. Effective public health interventions are crucial for addressing malaria, anemia, and iron deficiency during pregnancy. Strategies such as intermittent preventive treatment in pregnancy (IPTp), use of insecticide-treated bed nets (ITNs), and routine iron supplementation are essential components of a comprehensive approach. These interventions aim to reduce the incidence of malaria, prevent and treat anemia, and improve overall maternal and fetal health outcomes. Integrating malaria and anemia management within maternal and child health services is essential for providing comprehensive care. This includes ensuring that antenatal care incorporates both malaria prevention and iron supplementation, as well as addressing other related health issues. Coordinated care approaches help streamline health services, improve adherence to interventions, and enhance overall health outcomes for pregnant women.<sup>11-20</sup>

## **Malaria and Pregnancy**

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

Malaria during pregnancy represents a critical public health issue, particularly in regions where the disease is endemic. The infection, caused by *PLASMODIUM* species transmitted through the bite of *Anopheles* mosquitoes, poses significant risks to both the mother and the fetus. The unique physiological changes occurring during pregnancy make pregnant women more susceptible to malaria, and the presence of malaria can exacerbate pregnancy-related complications, leading to adverse outcomes. Pregnant women are more vulnerable to malaria due to changes in immune function and increased blood volume. Malaria infection during pregnancy can lead to severe anemia, which results from both the destruction of red blood cells by the malaria parasites and the inflammatory response induced by the infection. Severe anemia can cause fatigue, weakness, and increase the risk of maternal mortality. Additionally, malaria can trigger other complications such as placental malaria, which can disrupt the delivery of nutrients and oxygen to the fetus, further impacting maternal health. The effects of malaria on fetal health are profound and multifaceted. The infection can lead to adverse fetal outcomes such as low birth weight, preterm birth, and intrauterine growth restriction (IUGR). Placental malaria, characterized by the accumulation of malaria parasites in the placenta, impairs the placental function and disrupts fetal nutrient and oxygen supply. This can result in poor fetal growth and development. Malaria has also been associated with an increased risk of stillbirth and neonatal death.<sup>21-25</sup>

The pathophysiology of malaria in pregnancy involves complex interactions between the malaria parasites, the host's immune system, and the placenta. *PLASMODIUM FALCIPARUM*, the most severe malaria species, can adhere to the placenta through a process known as cytoadherence. This adhesion of infected red blood cells to the placental tissue causes inflammation and impairs blood flow. The inflammatory response leads to the production of cytokines and other mediators that can exacerbate maternal anemia and negatively affect fetal development. Pregnant women experience changes in immune function that can influence their susceptibility to malaria. The immune system during pregnancy undergoes a shift to accommodate the developing fetus, which can make pregnant women less able to mount an effective response to malaria. Additionally, the malaria parasites have evolved mechanisms to evade the immune system, such as antigenic variation and sequestration in the placenta, making it challenging for the body to clear the infection. Epidemiological studies highlight the high prevalence of malaria among pregnant women in endemic regions, with significant variations based on geographic location and season. In sub-Saharan Africa, where malaria transmission is intense, pregnant women are at increased risk of malaria and its complications. Studies have shown that effective malaria control measures, including intermittent preventive treatment (IPTp) and insecticide-treated bed nets (ITNs), are crucial in reducing the incidence of malaria and improving maternal and fetal health outcomes. Addressing malaria in pregnancy requires a multifaceted approach. Intermittent preventive treatment in pregnancy (IPTp) with sulfadoxine-pyrimethamine (SP) is recommended by the World Health Organization (WHO) to prevent malaria and reduce its complications. Insecticide-treated bed nets (ITNs) provide protection against mosquito bites and reduce malaria transmission. Prompt diagnosis and treatment of malaria are essential to prevent severe outcomes and manage the infection effectively.<sup>26-35</sup>

### **Anemia in Pregnancy**

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

Anemia is a common condition during pregnancy that can have significant implications for both maternal and fetal health. Defined as a reduction in hemoglobin concentration below normal levels, anemia affects the oxygen-carrying capacity of the blood, which is crucial for meeting the increased physiological demands of pregnancy. The condition is often multifactorial, with iron deficiency being the most prevalent cause, although other factors such as folate deficiency, vitamin B12 deficiency, and chronic diseases can also contribute. Anemia in pregnancy is a widespread issue, particularly in developing countries where nutritional deficiencies and infections are more common. According to the World Health Organization (WHO), approximately 40% of pregnant women in low- and middle-income countries are affected by anemia. The most common cause of anemia during pregnancy is iron deficiency, which results from increased iron requirements due to the expanding blood volume and fetal growth. Other causes include folate deficiency, which is essential for red blood cell production, and vitamin B12 deficiency, which affects red blood cell maturation. The pathophysiology of anemia during pregnancy involves several mechanisms. Iron deficiency impairs the production of hemoglobin, leading to a reduced number of red blood cells. This, in turn, affects the oxygen delivery to tissues and organs. Folate and vitamin B12 deficiencies disrupt red blood cell formation and maturation, leading to ineffective erythropoiesis. Additionally, chronic diseases and infections can contribute to anemia by affecting red blood cell production or increasing red blood cell destruction.<sup>36-40</sup>

Anemia during pregnancy is associated with a range of adverse health outcomes for the mother. Severe anemia can lead to fatigue, weakness, and decreased exercise tolerance, which can impact daily functioning and quality of life. In severe cases, anemia increases the risk of maternal mortality and complications such as postpartum hemorrhage. Anemia also makes pregnant women more susceptible to infections and can exacerbate the effects of other pregnancy-related conditions. The effects of maternal anemia extend to the fetus, with potential consequences for fetal growth and development. Anemia can lead to low birth weight, preterm birth, and intrauterine growth restriction (IUGR). In severe cases, it can result in stillbirth or neonatal death. Iron deficiency in particular can impair cognitive and motor development in children, with potential long-term effects on academic performance and overall health. Diagnosing anemia during pregnancy typically involves measuring hemoglobin levels through a complete blood count (CBC). Additional tests may be required to determine the underlying cause of anemia, including serum ferritin levels to assess iron stores, and tests for folate and vitamin B12 levels. Identifying the specific cause of anemia is crucial for selecting appropriate treatment and management strategies.<sup>41-45</sup>

The management of anemia in pregnancy depends on its underlying cause. For iron deficiency anemia, oral iron supplements are commonly prescribed, often in combination with folic acid to address concurrent folate deficiency. In cases of severe anemia or when oral supplementation is ineffective, intravenous iron or blood transfusions may be necessary. Addressing folate and vitamin B12 deficiencies involves supplementation with these nutrients. Nutritional counseling and education on dietary sources of iron, folate, and vitamin B12 are also important components of management. Public health interventions to address anemia during pregnancy include routine screening and supplementation programs. The WHO recommends iron and folic acid supplementation for all pregnant women, particularly in regions with high rates of anemia. Programs aimed at improving nutritional status through dietary diversification and fortification of

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

staple foods can help prevent anemia. Additionally, integrating anemia management into antenatal care services ensures that pregnant women receive timely diagnosis and treatment. Iron deficiency is a prevalent nutritional issue that significantly impacts maternal health during pregnancy. As an essential component of hemoglobin, iron plays a crucial role in oxygen transport and cellular metabolism. During pregnancy, the demand for iron increases to support fetal growth, expanded maternal blood volume, and placental development. Insufficient iron intake or absorption can lead to iron deficiency, which is a common cause of anemia in pregnant women and has important implications for both maternal and fetal health. Iron deficiency occurs when the body's iron stores are depleted, impairing hemoglobin production and leading to reduced oxygen delivery to tissues and organs. During pregnancy, the physiological expansion of blood volume increases the iron requirements, making pregnant women more susceptible to iron deficiency. If iron deficiency progresses, it can lead to iron deficiency anemia (IDA), characterized by low hemoglobin levels and reduced red blood cell production. The condition can affect maternal health by diminishing the body's ability to cope with the increased demands of pregnancy.<sup>46-55</sup>

### **Impact on Maternal Health**

Iron deficiency can have a range of adverse effects on maternal health. The most common manifestations include fatigue, weakness, and decreased exercise tolerance, which can impact daily functioning and quality of life. Severe iron deficiency anemia can increase the risk of complications such as preterm labor, postpartum hemorrhage, and impaired immune function, making pregnant women more susceptible to infections. Additionally, untreated iron deficiency anemia is associated with increased maternal morbidity and mortality. Iron deficiency during pregnancy not only affects the mother but also has significant implications for fetal health. Insufficient maternal iron levels can lead to poor fetal growth, low birth weight, and intrauterine growth restriction (IUGR). In severe cases, iron deficiency can increase the risk of preterm birth and stillbirth. Long-term effects on the child may include cognitive and motor developmental delays, as iron is crucial for brain development and function. Addressing maternal iron deficiency is essential for supporting optimal fetal development and reducing the risk of adverse outcomes. Diagnosing iron deficiency involves assessing serum ferritin levels, which reflect the body's iron stores. Other diagnostic tests may include measuring hemoglobin levels, serum iron, total iron-binding capacity (TIBC), and transferrin saturation. A comprehensive evaluation helps determine the severity of iron deficiency and guides appropriate treatment strategies. Routine screening for iron deficiency and anemia is recommended during antenatal care to identify and address issues early. The primary treatment for iron deficiency involves oral iron supplementation. Ferrous sulfate is commonly prescribed due to its high bioavailability and efficacy in increasing iron levels. In cases of severe anemia or when oral iron is not tolerated or effective, intravenous iron may be administered. Alongside iron supplements, dietary counseling is important to encourage the intake of iron-rich foods such as lean meats, legumes, and fortified cereals. Vitamin C can enhance iron absorption, so dietary recommendations often include foods rich in vitamin C. Public health interventions play a critical role in addressing iron deficiency during pregnancy. The World Health Organization (WHO) recommends routine iron and folic acid supplementation for all pregnant women, particularly in regions with high prevalence of anemia and iron deficiency. Programs that promote iron fortification of staple foods and improve access to iron supplements can help prevent

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76



and manage iron deficiency. Additionally, education on the importance of iron-rich diets and adherence to supplementation is essential for effective management.<sup>56-64</sup>

### Pathophysiological Mechanisms

Iron deficiency during pregnancy involves complex pathophysiological mechanisms that affect both maternal and fetal health. The interplay between increased iron requirements during pregnancy and the body's ability to meet these needs can lead to iron deficiency and its associated complications. During pregnancy, the demand for iron increases significantly due to several physiological changes. These include the expansion of maternal blood volume, which requires additional iron for hemoglobin synthesis, and the development of the placenta, which needs iron for fetal nourishment. Additionally, iron is essential for the growth and development of the fetus. The increased iron requirement is particularly critical in the second and third trimesters when fetal growth accelerates. Iron is absorbed primarily in the duodenum and upper jejunum of the small intestine. The absorption process is regulated by several factors, including dietary intake, the form of iron (heme versus non-heme), and the presence of enhancing or inhibiting substances in the diet. Vitamin C, for example, enhances non-heme iron absorption, while phytates and polyphenols inhibit it. In pregnancy, increased iron demands can lead to an imbalance between iron intake and absorption, especially if dietary intake is insufficient or if there are malabsorption issues.<sup>65-70</sup>

Once absorbed, iron is transported in the bloodstream bound to transferrin, a protein that delivers iron to various tissues and the developing fetus. Iron is stored in the liver, spleen, and bone marrow in the form of ferritin and hemosiderin. During pregnancy, iron is mobilized from these stores to meet the increased demands. In cases of inadequate dietary intake or absorption, these stores can become depleted, leading to iron deficiency. Iron deficiency anemia develops when there is an insufficient supply of iron for hemoglobin synthesis. Hemoglobin, the oxygen-carrying component of red blood cells, is crucial for maintaining adequate oxygen delivery to tissues. In iron deficiency anemia, the production of hemoglobin is impaired, leading to a reduced number of red blood cells (microcytic anemia) and decreased oxygen-carrying capacity. This results in symptoms such as fatigue, weakness, and decreased exercise tolerance. Iron deficiency anemia during pregnancy can have several adverse effects on maternal health. Reduced oxygen delivery to tissues can impair maternal function and increase the risk of complications such as preterm labor and postpartum hemorrhage. Additionally, severe anemia can compromise immune function, making pregnant women more susceptible to infections. The increased workload on the cardiovascular system can also exacerbate the effects of anemia, leading to potential cardiovascular complications.<sup>71-75</sup>

Iron deficiency during pregnancy can negatively impact fetal health and development. Adequate iron levels are crucial for fetal growth, brain development, and overall health. Iron deficiency can lead to low birth weight, intrauterine growth restriction (IUGR), and preterm birth. Long-term effects on the child may include cognitive and developmental delays, as iron is essential for neurodevelopment. The developing brain is particularly sensitive to iron deficiency, which can affect learning and behavioral outcomes. Iron deficiency can be exacerbated by inflammatory and metabolic factors. Chronic inflammation, often associated with infections or underlying diseases, can increase the production of hepcidin, a hormone that regulates iron homeostasis by inhibiting

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

iron absorption and mobilization. This can worsen iron deficiency by reducing the availability of iron for hematopoiesis. Additionally, metabolic stress and chronic diseases can interfere with iron utilization and exacerbate deficiency. Accurate diagnosis of iron deficiency involves assessing various biomarkers, including serum ferritin, serum iron, transferrin saturation, and total iron-binding capacity (TIBC). Monitoring these indicators helps to evaluate the severity of deficiency and guide appropriate treatment strategies. Regular screening during pregnancy is essential for early detection and management of iron deficiency anemia. Treatment of iron deficiency in pregnancy typically involves oral iron supplementation, which helps to replenish iron stores and correct anemia. In cases of severe anemia or when oral iron is not effective, intravenous iron may be required. Nutritional counseling to increase dietary intake of iron-rich foods and strategies to enhance iron absorption (such as combining iron with vitamin C) are also important components of treatment.<sup>76-80</sup>

## **Epidemiological Evidence**

Epidemiological studies provide valuable insights into the prevalence, risk factors, and outcomes associated with iron deficiency and anemia during pregnancy. These studies help in understanding the scope of the problem, identifying vulnerable populations, and evaluating the effectiveness of interventions. Iron deficiency and anemia are widespread issues in many parts of the world, particularly in low- and middle-income countries. According to the World Health Organization (WHO), approximately 40% of pregnant women in these regions are affected by anemia, with iron deficiency being the most common cause. Prevalence rates can vary based on geographic location, socioeconomic status, and access to healthcare. In regions with high malaria transmission, such as sub-Saharan Africa, the prevalence of anemia is often higher due to the combined effects of malaria and iron deficiency. Epidemiological data show significant regional variations in the prevalence of iron deficiency and anemia during pregnancy. For example, studies from sub-Saharan Africa and South Asia report high rates of anemia, with prevalence ranging from 30% to 60% among pregnant women. In contrast, prevalence rates are generally lower in high-income countries, although iron deficiency still poses a problem in certain populations. Variations in prevalence are influenced by factors such as dietary intake, access to healthcare, and the prevalence of other health conditions.<sup>81-83</sup>

Low dietary intake of iron-rich foods, poor absorption due to dietary inhibitors (e.g., phytates), and inadequate intake of complementary nutrients like vitamin C can contribute to iron deficiency. Lower socioeconomic status is often associated with higher rates of anemia due to limited access to nutritious foods and healthcare services. Living in areas with high malaria transmission or where other infections are prevalent can increase the risk of anemia due to the combined effects of infection and iron deficiency. Women with multiple pregnancies or closely spaced births are at higher risk of iron deficiency due to the increased iron demands associated with each pregnancy. Epidemiological studies have demonstrated the adverse effects of iron deficiency and anemia on both maternal and fetal health. Iron deficiency during pregnancy is linked to increased risks of maternal morbidity and mortality, including complications such as preterm birth, postpartum hemorrhage, and impaired immune function. For the fetus, iron deficiency can lead to low birth weight, intrauterine growth restriction (IUGR), and developmental delays. The long-term effects

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

on the child's cognitive and motor development have also been documented, emphasizing the importance of addressing iron deficiency early. Public health interventions, such as iron and folic acid supplementation programs, have been shown to be effective in reducing the prevalence of iron deficiency and anemia among pregnant women. Epidemiological evidence supports the implementation of routine supplementation as part of antenatal care in high-prevalence areas. Studies also highlight the benefits of combining supplementation with other strategies, such as iron fortification of staple foods and malaria prevention measures, to address the multifaceted nature of anemia.<sup>84-85</sup>

### **Public Health Interventions**

Addressing iron deficiency and anemia during pregnancy requires a comprehensive approach involving various public health interventions. These interventions aim to prevent, diagnose, and treat iron deficiency and anemia effectively, thereby improving maternal and fetal health outcomes. The cornerstone of public health interventions for iron deficiency anemia (IDA) in pregnancy is the provision of iron and folic acid supplements. The World Health Organization (WHO) recommends daily supplementation of iron (30–60 mg) and folic acid (400 µg) for all pregnant women, particularly in regions with high prevalence rates. This intervention helps replenish iron stores, correct anemia, and support the increased iron demands of pregnancy. In high-risk populations or areas with particularly high rates of anemia, targeted supplementation strategies may be implemented. This includes providing higher doses of iron or additional micronutrients as needed based on individual risk factors and anemia severity. Ensuring adherence to supplementation is crucial for achieving optimal health outcomes. Providing education on iron-rich foods and balanced diets is an important component of managing iron deficiency. Nutritional counseling helps pregnant women understand the importance of dietary iron and how to incorporate iron-rich foods such as lean meats, legumes, fortified cereals, and green leafy vegetables into their diets. Counseling also addresses factors that enhance or inhibit iron absorption, such as the role of vitamin C in improving non-heme iron absorption. Iron fortification of staple foods, such as flour, rice, and salt, is a widely used public health strategy to combat iron deficiency. Fortification programs aim to increase the iron content of commonly consumed foods, thus reaching a broad population and improving overall iron intake. Fortification efforts should be tailored to local dietary patterns and needs.<sup>86-87</sup>

In regions where malaria is endemic, malaria prevention and treatment are crucial for reducing anemia prevalence. Interventions such as insecticide-treated bed nets (ITNs), intermittent preventive treatment in pregnancy (IPTp) with sulfadoxine-pyrimethamine, and prompt diagnosis and treatment of malaria help prevent the combined effects of malaria and iron deficiency anemia. De-worming programs for pregnant women, particularly in areas with high prevalence of intestinal parasites, can help reduce anemia. Parasitic infections can contribute to anemia by causing blood loss and impairing nutrient absorption. Routine de-worming as part of antenatal care can help mitigate these effects. Integrated antenatal care services that include routine screening for anemia, regular monitoring, and targeted interventions are essential for effective management. Antenatal care should encompass anemia screening through hemoglobin measurement, assessment of iron status, and personalized treatment plans. This approach ensures that anemia is identified early and

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76



managed appropriately. Engaging community health workers and local organizations in anemia prevention and management can enhance the reach and effectiveness of public health interventions. Community-based programs can provide education, distribute supplements, and support adherence to treatment, particularly in remote or underserved areas. Developing and implementing national guidelines for anemia prevention and management during pregnancy is crucial. These guidelines should include recommendations for supplementation, dietary interventions, and integration with other health services. Policymakers should ensure that guidelines are based on local epidemiological data and resources. Regular monitoring and evaluation of anemia prevention programs are essential for assessing their impact and effectiveness. Data collection on anemia prevalence, supplement coverage, and health outcomes helps to identify gaps, improve strategies, and ensure that interventions are meeting their objectives.<sup>88-9</sup>

## Conclusion

Iron deficiency and anemia during pregnancy are significant public health challenges with profound implications for both maternal and fetal health. The increasing demand for iron during pregnancy, combined with potential barriers to adequate intake and absorption, underscores the need for comprehensive and targeted public health interventions. Epidemiological evidence highlights the widespread nature of the problem, with varying prevalence rates influenced by geographic, socioeconomic, and health factors. Effective management of iron deficiency and anemia requires a multifaceted approach that includes routine iron and folic acid supplementation, dietary improvements, and strategies to control infections like malaria. Public health programs must integrate these interventions into comprehensive antenatal care to ensure early detection and appropriate treatment. Additionally, addressing socioeconomic and cultural barriers, improving access to healthcare, and strengthening health systems are crucial for enhancing the reach and impact of these programs.

## References

1. Uneke CJ. Impact of placental *Plasmodium falciparum* malaria on pregnancy and perinatal outcome in sub-Saharan Africa: part III: placental malaria, maternal health, and public health. *The Yale journal of biology and medicine*. 2008;81(1):1.
2. Gontie GB, Wolde HF, Baraki AG. Prevalence and associated factors of malaria among pregnant women in Sherkole district, Benishangul Gumuz regional state, West Ethiopia. *BMC Infectious Diseases*. 2020; 20:1-8.
3. Obeagu EI, Agree FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri*. 2023; 6 (1). 2023;138. [links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf](https://doi.org/10.3390/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf).
4. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci*. 2016;3(3):10-8. [links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf](https://doi.org/10.3390/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf).

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

5. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. *J Res Med Dent Sci*. 2021;9(4):145-8. [links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf](https://epjournals.com/links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf).
6. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
7. Jakheng SP, Obeagu EI. Seroprevalence of human immunodeficiency virus based on demographic and risk factors among pregnant women attending clinics in Zaria Metropolis, Nigeria. *J Pub Health Nutri*. 2022; 5 (8). 2022;137. [links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf](https://epjournals.com/links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf).
8. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. *Madonna University journal of Medicine and Health Sciences*. 2022;2(2):1-9.
9. Bonilla FA, Oettgen HC. Adaptive immunity. *Journal of Allergy and Clinical Immunology*. 2010;125(2): S33-40.
10. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(2):1-9.
11. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfectd with HIV in Umuahia. *Int. J. Curr. Res. Med. Sci*. 2017;3(5):100-104.
12. Feeney ME. The immune response to malaria in utero. *Immunological reviews*. 2020 ;293(1):216-229.
13. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. *Medicine*. 2023;102(46): e36166.
14. Obeagu EI, Chijioke UO, Ekelozie IS. Malaria rapid diagnostic test (RDTs). *Ann Clin Lab Res*. 2018;6(4):275.
15. Ogomaka IA, Obeagu EI. Methods of Breast Feeding as Determinants of Malaria Infections among Babies in IMO State, Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(01):17-24.
16. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(1):292-8. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
17. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. *J. Bio. Innov*11 (3). 2022:936-941.
18. Obeagu EI, Obeagu GU, Egba SI, Emeka-Obi OR. Combatting Anemia in Pediatric Malaria: Effective Management Strategies. *Int. J. Curr. Res. Med. Sci*. 2023;9(11):1-7.

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

19. Hassan AO, Oso OV, Obeagu EI, Adeyemo AT. Malaria Vaccine: Prospects and Challenges. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(2):22-40.
20. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-9.
21. Moya-Alvarez V, Abellana R, Cot M. Pregnancy-associated malaria and malaria in infants: an old problem with present consequences. *Malaria journal*. 2014; 13:1-10.
22. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2022;9(9): 21-6.DOI: [10.22192/ijcrrcps.2022.09.09.003](https://doi.org/10.22192/ijcrrcps.2022.09.09.003)
23. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
24. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State. Nigeria. *Journal of Research in Medical and Dental Science*. 2021;9(12):462-.[links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://doi.org/10.21961/links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf).
25. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022 Jun 21;5(2):113-121.
26. Obeagu EI. Gestational Thrombocytopaenia. *J Gynecol Women's Health*. 2023;25(3):556163. [links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf](https://doi.org/10.21961/links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf).
27. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
28. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
29. Obeagu EI, Busari AI, Uduchi IO, Ogomaka IA, Ibekwe AM, Vincent CC, Chijioke UO, Okafor CJ, Okoroiwu HU, Adike CN. Age-Related Haematological Variations in Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):218-24.
30. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

31. Ogbonna CO, Obeagu EI, Ufelle SA, Ogbonna LN. Evaluation of haematological alterations in children infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(1):38-45.
32. Appay V. The physiological role of cytotoxic CD4+ T-cells: the holy grail? *Clinical & Experimental Immunology*. 2004;138(1):10-13.
33. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(1):25-36.
34. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med*. 2018;3(9):1-4.
35. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with Plasmodium falciparum malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2016;3(8):40-46.
36. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology*. 2019;41(3): e12573.
37. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research*. 2017;5(4):206.
38. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Ironi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal*. 2022;6(3):9-17.
39. Madekwe CC, Madekwe CC, Obeagu EI. Inequality of monitoring in Human Immunodeficiency Virus, Tuberculosis and Malaria: A Review. *Madonna University journal of Medicine and Health Sciences*. 2022;2(3):6-15.
40. Offie DC, Ibekwe AM, Agu CC, Esimai BN, Okpala PU, Obeagu EI, Ufelle SA, Ogbonna LN. Fibrinogen and C-Reactive Protein Significance in Children Infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(15):1-8.
41. Obeagu EI, Ogunnaya FU. PREGNANCY-INDUCED HAEMATOLOGICAL CHANGES: A KEY TO MATERNAL AND CHILD HEALTH. *European Journal of Biomedical*. 2023;10(8):42-43. [links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf](https://epjournals.com/links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf)
42. Obeagu EI, Ofodile AC, Okwuanaso CB. A review of urinary tract infections in pregnant women: Risks factors. *J Pub Health Nutri*. 2023; 6 (1). 2023; 137:26-35. [links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf](https://epjournals.com/links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf).
43. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. *Int. J. Curr. Res. Med. Sci*. 2023;9(2): 14-17.DOI: 10.22192/ijcrms.2023.09.02.003

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

44. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 14-21.DOI: [10.22192/ijcreps.2023.10.08.002](https://doi.org/10.22192/ijcreps.2023.10.08.002)
45. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
46. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9): 1-4.DOI: [10.22192/ijcrbm.2018.03.09.001](https://doi.org/10.22192/ijcrbm.2018.03.09.001)
47. Onyenweaku FC, Amah HC, Obeagu EI, Nwandikor UU, Onwuasoanya UF. Prevalence of asymptomatic bacteriuria and its antibiotic susceptibility pattern in pregnant women attending private ante natal clinics in Umuahia Metropolitan. *Int J Curr Res Biol Med.* 2017;2(2): 13-23.DOI: [10.22192/ijcrbm.2017.02.02.003](https://doi.org/10.22192/ijcrbm.2017.02.02.003)
48. Okoroiwu IL, Chinedu-Madu JU, Obeagu EI, Vincent CC, Ochiabuto OM, Ibekwe AM, Amaechi CO, Agu CC, Anoh NV, Amadi NM. Evaluation of Iron Status, Haemoglobin and Protein Levels of Pregnant Women in Owerri Metropolis. *Journal of Pharmaceutical Research International*. 2021;33(27A):36-43.
49. Obeagu EI, Njar VE, Obeagu GU. Infertility: Prevalence and Consequences. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(7):43-50.
50. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
51. Broen K, Brustoski K, Engelmann I, Luty AJ. Placental Plasmodium falciparum infection: causes and consequences of in utero sensitization to parasite antigens. *Molecular and biochemical parasitology*. 2007;151(1):1-8.
52. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9):41-49.
53. Ogbonna LN, Ezeoru VC, Ofodile AC, Ochiabuto OM, Obi-Ezeani CN, Okpala PU, Okafor CJ, Obeagu GU, Busari AI, Obeagu EI. Gender Based Variations of Haematological Parameters of Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(8):75-80.
54. Eberendu IF, Ozims SJ, Agu GC, Amah HC, Obasi CC, Obioma-Elemba JE, Ihekare DE, Ibanga IE, Amah CC, Obeagu EI, Nwosu DC. Impact of human activities on the breeding of mosquitoes of human disease in Owerri metropolis, Imo state. *Int J Adv Res Biol Sci IJARBS*. 2017;4(12):98-106.
55. Obeagu EI, Ofodile AC, Okwuanaso CB. A review on socio economic and behavioral aspects of malaria and its control among children under 5 years of age in Africa. *J Pub Health Nutri*. 2023; 6 (1): 136.
56. Djontu JC, Siewe Siewe S, Mpeke Edene YD, Nana BC, Chomga Foko EV, Bigoga JD, Leke RF, Megnekou R. Impact of placental Plasmodium falciparum malaria infection on the Cameroonian maternal and neonate's plasma levels of some cytokines known to regulate T cells differentiation and function. *Malaria journal*. 2016; 15:1-1.

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76



57. Obeagu EI, Faduma MH, Uzoma G. Ectopic Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(4): 40-4.DOI: [10.22192/ijcrps.2023.10.04.004](https://doi.org/10.22192/ijcrps.2023.10.04.004)
58. Obeagu EI, Gamade SM, Obeagu GU. The roles of Neutrophils in pregnancy. *Int. J. Curr. Res. Med. Sci.* 2023;9(5): 31-35.DOI: [10.22192/ijcrms.2023.09.05.005](https://doi.org/10.22192/ijcrms.2023.09.05.005)
59. Obeagu EI, Obeagu GU. Molar Pregnancy: Update of prevalence and risk factors. *Int. J. Curr. Res. Med. Sci.* 2023;9(7): 25-28.DOI: [10.22192/ijcrms.2023.09.07.005](https://doi.org/10.22192/ijcrms.2023.09.07.005)
60. Kabyemela E, Gonçalves BP, Prevots DR, Morrison R, Harrington W, Gwamaka M, Kurtis JD, Fried M, Duffy PE. Cytokine profiles at birth predict malaria severity during infancy. *PloS one.* 2013;8(10):e77214.
61. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Barriers to utilization of maternal health services in southern senatorial district of Cross Rivers state, Nigeria. *International Journal of Advanced Multidisciplinary Research.* 2017;4(8): 1-9.DOI: [10.22192/ijamr.2017.04.08.001](https://doi.org/10.22192/ijamr.2017.04.08.001)
62. Emmanuel G, Martin O, Peter OS, Obeagu EI, Daniel K. Factors Influencing Early Neonatal Adverse Outcomes among Women with HIV with Post Dated Pregnancies Delivering at Kampala International University Teaching Hospital, Uganda. *Asian Journal of Pregnancy and Childbirth.* 2023;6(1):203-211.  
<http://research.sdpublishers.net/id/eprint/2819/>.
63. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences.* 2018;4(9): 41-9.DOI: [10.22192/ijcrms.2018.04.09.006](https://doi.org/10.22192/ijcrms.2018.04.09.006)
64. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 22-6.DOI: [10.22192/ijcrps.2023.10.08.003](https://doi.org/10.22192/ijcrps.2023.10.08.003)
65. Obeagu E, Eze RI, Obeagu EI, Nnatuanya IN, Dara EC. ZINC LEVEL IN APPARENTLY PREGNANT WOMEN IN URBAN AREA. *Madonna University journal of Medicine and Health Sciences.* 2022;2(1):134-48.  
<https://www.journal.madonnauniversity.edu.ng/index.php/medicine/article/view/40>.
66. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(41B):380-386.
67. Obeagu EI, Ogunnaya FU, Obeagu GU, Ndidi AC. SICKLE CELL ANAEMIA: A GESTATIONAL ENIGMA. *migration.* 2023; 17:18.
68. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology.* 2019;41(3):e12573.
69. Ifeanyi OE, Uzoma OG. A review on erythropietin in pregnancy. *J. Gynecol. Womens Health.* 2018;8(3):1-4.  
[https://www.academia.edu/download/56538560/A\\_Review\\_on\\_Erythropietin\\_in\\_Pregnancy.pdf](https://www.academia.edu/download/56538560/A_Review_on_Erythropietin_in_Pregnancy.pdf).
70. Ifeanyi OE. A review on pregnancy and haematology. *Int. J. Curr. Res. Biol. Med.* 2018;3(5): 26-8.DOI: [10.22192/ijcrbm.2018.03.05.006](https://doi.org/10.22192/ijcrbm.2018.03.05.006)

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

71. Nwosu DC, Nwanjo HU, Obeagu EI, Ibebuike JE, Ezeama MC. Ihekireh. Changes in liver enzymes and lipid profile of pregnant women with malaria in Owerri, Nigeria. *International Journal of Current Research and Academic Review*. 2015;3(5):376-383.
72. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Factors that influence women's utilization of primary health care services in Calabar Cros river state, Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2017;4(7):28-33.
73. Elemchukwu Q, Obeagu EI, Ochei KC. Prevalence of Anaemia among Pregnant Women in Braithwaite Memorial Specialist Hospital (BMSH) Port Harcourt. *IOSR Journal of Pharmacy and Biological Sciences*. 2014;9(5):59-64.
74. Natama HM, Moncunill G, Rovira-Vallbona E, Sanz H, Sorgho H, Aguilar R, Coulibaly-Traoré M, Somé MA, Scott S, Valéa I, Mens PF. Modulation of innate immune responses at birth by prenatal malaria exposure and association with malaria risk during the first year of life. *BMC medicine*. 2018; 16:1-5.
75. Akandinda M, Obeagu EI, Katonera MT. Non Governmental Organizations and Women's Health Empowerment in Uganda: A Review. *Asian Research Journal of Gynaecology and Obstetrics*. 2022;8(3):12-26.
76. Gamde MS, Obeagu EI. IRON DEFICIENCY ANAEMIA: ENEMICAL TO PREGNANCY. *European Journal of Biomedical*. 2023;10(9):272-275.  
[links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf](https://doi.org/10.15406/ejbr.2023.10.00272).
77. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclampsic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
78. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Studies of Some Haemostatic Variables in Preeclampsic Women in Owerri, Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):39-48.
79. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8):22-26.
80. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023;6(2):10-13.
81. Mutabingwa TK, Bolla MC, Li JL, Domingo GJ, Li X, Fried M, Duffy PE. Maternal malaria and gravidity interact to modify infant susceptibility to malaria. *PLoS medicine*. 2005;2(12):e407.
82. Gamble C, Ekwaru PJ, Garner P, Ter Kuile FO. Insecticide-treated nets for the prevention of malaria in pregnancy: a systematic review of randomised controlled trials. *PLoS medicine*. 2007;4(3):e107.
83. Okoko BJ, Enwere G, Ota MO. The epidemiology and consequences of maternal malaria: a review of immunological basis. *Acta tropica*. 2003;87(2):193-205.
84. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.

**Citation:** Obeagu EI, Obeagu GU. Malaria, Anemia, and Iron Deficiency in Pregnancy: An Integrated Review. *Elite Journal of Scientific Research and Review*, 2024; 2(4): 61-76

85. Umbers AJ, Stanisić DI, Ome M, Wangnapi R, Hanieh S, Unger HW, Robinson LJ, Lufele E, Baiwog F, Siba PM, King CL. Does malaria affect placental development? Evidence from in vitro models. *PLoS One*. 2013;8(1):e55269.
86. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
87. Gbedande K, Carpio VH, Stephens R. Using two phases of the CD 4 T cell response to blood-stage murine malaria to understand regulation of systemic immunity and placental pathology in *Plasmodium falciparum* infection. *Immunological reviews*. 2020;293(1):88-114.
88. Lindsay SW, Thomas MB, Kleinschmidt I. Threats to the effectiveness of insecticide-treated bednets for malaria control: thinking beyond insecticide resistance. *The Lancet Global Health*. 2021;9(9):e1325-1331.
89. Akinleye SO, Falade CO, Ajayi IO. Knowledge and utilization of intermittent preventive treatment for malaria among pregnant women attending antenatal clinics in primary health care centers in rural southwest, Nigeria: a cross-sectional study. *BMC pregnancy and childbirth*. 2009; 9:1-9.