

The Role of Placental Malaria in the Development of Childhood Malaria: A Review

*Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²

¹Department of Medical Laboratory Science, Kampala International University, Ishaka, Uganda.

²School of Nursing Science, Kampala International University, Ishaka, Uganda.

*Corresponding author: Emmanuel Ifeanyi Obeagu, [Department of Medical Laboratory Science, Kampala International University, Uganda, emmanuelobeagu@yahoo.com, ORCID: 0000-0002-4538-0161](#)

Abstract

Placental malaria, characterized by the sequestration of Plasmodium parasites in the placenta, significantly impacts maternal and neonatal health. This condition is associated with increased risk of adverse pregnancy outcomes, including low birth weight and preterm birth, which can have lasting effects on the child's health. This review explores the role of placental malaria in the development of childhood malaria, focusing on how intrauterine exposure to malaria influences the infant's immune system and susceptibility to malaria in early childhood. The pathophysiological mechanisms linking placental malaria to childhood malaria include the impact of placental inflammation on fetal immune development and the potential for continued parasite transmission from mother to infant. Epidemiological evidence indicates that infants born to mothers with placental malaria are at higher risk for recurrent malaria and severe disease. These findings highlight the importance of understanding the long-term effects of placental malaria on child health. Effective public health interventions are crucial for mitigating the impact of placental malaria on childhood malaria. Integrated strategies combining malaria prevention, treatment, and maternal health measures, such as the use of insecticide-treated bed nets (ITNs) and intermittent preventive treatment in pregnancy (IPTp), are essential. Community-based approaches and enhanced surveillance can further support efforts to reduce placental malaria and improve health outcomes for both mothers and their children.

Keywords: *Placental Malaria, Childhood Malaria, Maternal-Infant Health, Malaria Immunity, Plasmodium falciparum, Intrauterine Infection, Epidemiological Impact*

Introduction

Placental malaria is a significant complication of malaria in pregnancy, characterized by the sequestration of Plasmodium parasites within the placenta. This condition is associated with a range of adverse pregnancy outcomes, including preterm delivery, low birth weight, and increased maternal morbidity. However, the impact of placental malaria extends beyond the immediate

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

effects on pregnancy, influencing the health and development of the infant. **Malaria in pregnancy** is primarily caused by *Plasmodium falciparum*, although other *Plasmodium* species can also contribute. The parasites adhere to the placental tissue, causing inflammation and compromising placental function. This results in reduced nutrient and oxygen supply to the fetus, which can affect fetal development and lead to complications. The disruption in placental function not only impacts the immediate health of the newborn but can also have long-term effects on the child's susceptibility to malaria and other infections.¹⁻⁵ **Fetal immune system development** is significantly influenced by the intrauterine environment. Maternal malaria-induced inflammation and nutrient deficiencies can alter the development of the fetal immune system. Studies suggest that infants exposed to placental malaria may have altered immune cell populations and cytokine profiles, which can affect their ability to respond to infections and vaccines. These changes can predispose infants to higher rates of malaria and other infectious diseases. **Intrauterine transmission** of malaria is another crucial factor linking placental malaria to childhood malaria. The presence of malaria parasites in the placenta can facilitate the transmission of parasites to the fetus, leading to congenital malaria. Although congenital malaria is relatively rare, it can contribute to the infant's early exposure to malaria and influence their subsequent health and immunity. **Epidemiological evidence** highlights the correlation between placental malaria and increased risk of malaria in early childhood. Research has shown that infants born to mothers with placental malaria are more likely to experience higher rates of malaria episodes and severe disease. These findings underscore the need for targeted interventions to reduce the prevalence of placental malaria and its impact on childhood malaria.⁶⁻¹⁵

Pathophysiological mechanisms underlying the link between placental malaria and childhood malaria include the effects of placental inflammation on fetal immune development and the potential for continued parasite exposure. Chronic inflammation in the placenta can disrupt normal immune system maturation, leading to altered immune responses in the infant. Additionally, persistent parasite presence in the placenta can increase the risk of transmission to the newborn. **Public health interventions** aimed at reducing the impact of placental malaria on childhood malaria are essential. Integrated malaria control strategies, such as the use of insecticide-treated bed nets (ITNs) and intermittent preventive treatment in pregnancy (IPTp), can help prevent placental malaria and improve maternal and infant health. Enhanced monitoring and surveillance of malaria prevalence and its impact on maternal and child health are also crucial for effective intervention. **Community-based approaches** play a vital role in addressing the challenges associated with placental malaria. Educating communities about malaria prevention, timely treatment, and the importance of antenatal care can help reduce the prevalence of placental malaria and improve health outcomes. Community health workers and local health programs are key in disseminating information and supporting malaria control efforts.¹⁶⁻²⁰

Pathophysiological Mechanisms

The pathophysiological mechanisms through which placental malaria influences childhood malaria are multifaceted, involving both direct and indirect effects on maternal and fetal health. *Plasmodium falciparum* parasites adhere to the placenta via a process called sequestration. The parasites bind to specific receptors on the placental syncytiotrophoblast cells, leading to the

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

formation of infected red blood cells in the placenta. This sequestration disrupts normal placental function and contributes to chronic inflammation. The inflammatory response in the placenta can compromise nutrient and oxygen transfer to the fetus, leading to adverse pregnancy outcomes such as low birth weight and preterm birth. The presence of malaria parasites in the placenta triggers a local inflammatory response characterized by the infiltration of immune cells, including macrophages and lymphocytes. This inflammation can result in the release of pro-inflammatory cytokines and chemokines, which further exacerbate tissue damage and impair placental function. Chronic inflammation in the placenta can affect fetal development and the maturation of the fetal immune system, potentially increasing the risk of malaria in early childhood.²¹⁻²⁵ The inflammatory environment created by placental malaria can significantly impact the development of the fetal immune system. Research has shown that exposure to maternal malaria can alter the composition of fetal immune cell populations and cytokine profiles. For instance, elevated levels of inflammatory cytokines such as tumor necrosis factor-alpha (TNF-alpha) and interleukin-6 (IL-6) in the placenta can affect the differentiation and function of fetal T cells and B cells. Infants born to mothers with placental malaria may exhibit impaired immune responses to infections and vaccines. The altered immune development resulting from placental malaria exposure can lead to reduced production of antibodies and ineffective cellular immune responses. This compromised immune function can increase the susceptibility of infants to malaria and other infectious diseases, impacting their overall health and development.²⁶⁻³⁰

Placental malaria can facilitate the intrauterine transmission of Plasmodium parasites to the fetus, leading to congenital malaria. Although congenital malaria is relatively rare, it can occur when parasites cross the placental barrier and infect the fetal blood supply. Infants with congenital malaria may present with symptoms shortly after birth, including fever, anemia, and low birth weight. The presence of malaria parasites in the newborn can contribute to early-life malaria susceptibility. Infants who acquire malaria in utero may experience a higher risk of developing malaria in early childhood. The initial exposure to malaria parasites can influence the infant's immune system and increase their vulnerability to subsequent malaria infections. Addressing congenital malaria and its effects on early childhood health is crucial for reducing the long-term impact of placental malaria.³¹⁻³⁵ The impact of placental malaria on maternal and fetal nutrition is another important aspect of the pathophysiological mechanisms. The inflammatory response in the placenta can affect the transfer of essential nutrients from the mother to the fetus, leading to deficiencies in critical vitamins and minerals. Nutritional deficiencies, such as iron and folate, can further compromise fetal development and immune function, increasing the risk of malaria and other health complications. Placental malaria can also disrupt maternal and fetal metabolism, leading to altered metabolic profiles. For example, chronic inflammation and nutrient deficiencies can affect glucose metabolism and increase the risk of metabolic disorders in both the mother and the infant. These metabolic disturbances can contribute to adverse health outcomes and exacerbate the impact of malaria. Strategies that combine malaria prevention and treatment with maternal and infant health measures are essential for mitigating the effects of placental malaria. Interventions such as insecticide-treated bed nets (ITNs), intermittent preventive treatment in pregnancy (IPTp), and enhanced nutritional support can help reduce the prevalence of placental malaria and its impact on childhood health.³⁶⁻⁴⁰

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

Epidemiological Evidence

Epidemiological studies have provided critical insights into the relationship between placental malaria and childhood malaria, highlighting how maternal malaria can influence the health of infants in the early years of life. Epidemiological data reveal that placental malaria is a significant concern in malaria-endemic regions, particularly in sub-Saharan Africa. Studies have shown varying prevalence rates depending on geographic location, with higher rates in areas with high malaria transmission. For example, in some regions of Africa, the prevalence of placental malaria can exceed 30% among pregnant women. The prevalence of placental malaria is closely linked to maternal health and malaria control measures. Factors such as inadequate access to antenatal care, low utilization of insecticide-treated bed nets (ITNs), and insufficient coverage of intermittent preventive treatment in pregnancy (IPTp) can contribute to higher rates of placental malaria. Epidemiological studies highlight the need for comprehensive malaria control strategies that address both prevention and treatment during pregnancy. Epidemiological studies consistently demonstrate that infants born to mothers with placental malaria have a higher risk of developing malaria in early childhood. For instance, research has shown that these infants experience more frequent malaria episodes compared to those born to uninfected mothers. The increased risk is attributed to both direct effects of placental malaria on immune system development and potential continued parasite exposure from the mother. The impact of placental malaria extends beyond the immediate postnatal period. Longitudinal studies have shown that children exposed to placental malaria are at increased risk of recurrent malaria and severe disease later in childhood. These findings underscore the long-term effects of placental malaria on health and highlight the importance of early and continued malaria prevention and treatment.⁴¹⁻⁵⁰

Epidemiological evidence also suggests that placental malaria can affect the efficacy of childhood vaccinations. Studies have shown that infants exposed to placental malaria may exhibit altered immune responses to vaccines, potentially leading to reduced vaccine effectiveness. This is of particular concern for vaccines against malaria and other infectious diseases, as compromised vaccine responses can affect overall health outcomes. The altered immune responses observed in infants exposed to placental malaria are linked to the inflammatory environment created by maternal malaria. Research indicates that changes in immune cell populations and cytokine profiles can affect the development of effective immune responses to vaccines, emphasizing the need for tailored vaccination strategies for this population. The epidemiological evidence highlights the importance of integrating malaria control strategies with maternal and child health programs. Effective interventions, such as ITNs and IPTp, are crucial for reducing the prevalence of placental malaria and improving childhood health outcomes. Public health programs should focus on enhancing access to these interventions and ensuring adherence to recommended practices. Ongoing surveillance and monitoring are essential for assessing the effectiveness of malaria control measures and identifying emerging issues. Epidemiological data on placental malaria prevalence, childhood malaria incidence, and vaccine efficacy can inform public health strategies and guide the implementation of targeted interventions. Engaging communities in malaria prevention and control efforts is crucial for reducing the impact of placental malaria. Community health education, outreach programs, and support for maternal and child health

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

services can enhance the effectiveness of malaria control measures and improve health outcomes for both mothers and their children.⁵¹⁻⁶⁶

Public Health Interventions

Effective public health interventions are crucial for addressing the impact of placental malaria on maternal and childhood health. By integrating malaria prevention and treatment strategies with maternal and child health programs, public health initiatives can mitigate the adverse effects of placental malaria and improve overall health outcomes. **Insecticide-Treated Bed Nets (ITNs)** are a cornerstone of malaria prevention, providing a barrier against mosquito bites and reducing the transmission of malaria. The widespread use of ITNs during pregnancy is essential for preventing placental malaria. Public health campaigns should focus on increasing the distribution and use of ITNs among pregnant women, ensuring that they receive adequate protection from malaria. **Intermittent Preventive Treatment in Pregnancy (IPTp)** involves administering antimalarial drugs to pregnant women at scheduled intervals to prevent malaria infection and its complications. The World Health Organization (WHO) recommends IPTp with sulfadoxine-pyrimethamine (SP) for pregnant women in malaria-endemic areas. Ensuring high coverage of IPTp through routine antenatal care visits is crucial for reducing the incidence of placental malaria and improving maternal and infant health outcomes.⁶⁷⁻⁷⁷

Timely diagnosis and treatment of malaria in pregnant women are essential for preventing placental malaria and its associated complications. Public health programs should emphasize the importance of early diagnosis using rapid diagnostic tests (RDTs) and prompt treatment with appropriate antimalarial medications. Access to quality healthcare services and effective treatment options should be prioritized to reduce the burden of malaria in pregnancy. Pregnant women with severe malaria or complications associated with placental malaria require specialized care. Public health interventions should include the development of guidelines and protocols for managing severe cases of malaria during pregnancy, ensuring that healthcare providers are equipped to handle complex cases and provide appropriate care. Routine antenatal care visits provide an opportunity to monitor for signs of malaria and other complications. Public health programs should promote regular antenatal check-ups, including screening for malaria and other infectious diseases. Early identification of placental malaria and other issues can lead to timely intervention and improved health outcomes for both mother and child. Educating pregnant women about malaria prevention, symptoms, and the importance of antenatal care is essential for improving health outcomes. Public health campaigns should focus on raising awareness about the risks of malaria during pregnancy, the benefits of ITNs and IPTp, and the importance of seeking timely medical care. Engaging communities in malaria prevention efforts is crucial for reducing the prevalence of placental malaria. Community health education programs can provide information on malaria prevention, symptoms, and treatment, empowering individuals to take proactive measures to protect themselves and their families. Strengthening maternal and child health services at the community level is essential for improving health outcomes. Public health interventions should include support for local health facilities, training for healthcare providers, and efforts to increase access to antenatal care and malaria prevention services.⁷⁸⁻⁸⁹

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

Conclusion

Placental malaria remains a significant public health challenge with profound implications for both maternal and child health. The sequestration of Plasmodium parasites in the placenta leads to inflammation, impaired placental function, and increased risk of adverse pregnancy outcomes such as low birth weight and preterm birth. These complications extend beyond pregnancy, affecting the infant's immune system development and increasing susceptibility to malaria and other infections in early childhood. Epidemiological evidence underscores the strong association between placental malaria and increased risk of childhood malaria, highlighting the need for effective public health interventions. The impact of placental malaria on childhood health is multifaceted, involving altered immune responses, potential congenital malaria, and long-term health consequences. Addressing these issues requires a comprehensive approach that integrates malaria prevention, early diagnosis, and effective treatment. Key public health interventions include the widespread use of insecticide-treated bed nets (ITNs), intermittent preventive treatment in pregnancy (IPTp), and prompt treatment of malaria in pregnant women. Enhanced antenatal care, community-based education, and support for maternal and child health services are also crucial. These strategies not only help prevent placental malaria but also mitigate its impact on childhood health.

References

1. Uneke CJ. Impact of placental Plasmodium falciparum malaria on pregnancy and perinatal outcome in sub-Saharan Africa: part III: placental malaria, maternal health, and public health. The Yale journal of biology and medicine. 2008;81(1):1.
2. Gontie GB, Wolde HF, Baraki AG. Prevalence and associated factors of malaria among pregnant women in Sherkole district, Benishangul Gumuz regional state, West Ethiopia. BMC Infectious Diseases. 2020; 20:1-8.
3. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. J Pub Health Nutri. 2023; 6 (1). 2023;138. [links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf](https://epjournals.com/links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf).
4. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. Int J Curr Res Chem Pharm Sci. 2016;3(3):10-8. [links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf](https://epjournals.com/links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf).
5. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. J Res Med Dent Sci. 2021;9(4):145-8. [links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf](https://epjournals.com/links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf).
6. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. International Research in Medical and Health Sciences. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
7. Jakheng SP, Obeagu EI. Seroprevalence of human immunodeficiency virus based on demographic and risk factors among pregnant women attending clinics in Zaria Metropolis, Nigeria. J Pub Health Nutri. 2022; 5 (8). 2022;137.

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

[links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf](https://epjournals.com/journals/EJSRR/links/6317a6b1acd814437f0ad268/Seroprevalence-of-human-immunodeficiency-virus-based-on-demographic-and-risk-factors-among-pregnant-women-attending-clinics-in-Zaria-Metropolis-Nigeria.pdf).

8. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. *Madonna University journal of Medicine and Health Sciences*. 2022;2(2):1-9.
9. Bonilla FA, Oettgen HC. Adaptive immunity. *Journal of Allergy and Clinical Immunology*. 2010;125(2): S33-40.
10. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(2):1-9.
11. Obeagu EI, Ibeh NC, Nwobodo HA, Ochei KC, Iwegbulam CP. Haematological indices of malaria patients coinfectd with HIV in Umuahia. *Int. J. Curr. Res. Med. Sci*. 2017;3(5):100-104.
12. Feeney ME. The immune response to malaria in utero. *Immunological reviews*. 2020 ;293(1):216-229.
13. Opeyemi AA, Obeagu EI. Regulations of malaria in children with human immunodeficiency virus infection: A review. *Medicine*. 2023;102(46): e36166.
14. Obeagu EI, Chijioke UO, Ekelozie IS. Malaria rapid diagnostic test (RDTs). *Ann Clin Lab Res*. 2018;6(4):275.
15. Ogomaka IA, Obeagu EI. Methods of Breast Feeding as Determinants of Malaria Infections among Babies in IMO State, Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(01):17-24.
16. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(1):292-8. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
17. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. *J. Bio. Innov*11 (3). 2022:936-941.
18. Obeagu EI, Obeagu GU, Egba SI, Emeka-Obi OR. Combatting Anemia in Pediatric Malaria: Effective Management Strategies. *Int. J. Curr. Res. Med. Sci*. 2023;9(11):1-7.
19. Hassan AO, Oso OV, Obeagu EI, Adeyemo AT. Malaria Vaccine: Prospects and Challenges. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022;2(2):22-40.
20. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-9.
21. Moya-Alvarez V, Abellana R, Cot M. Pregnancy-associated malaria and malaria in infants: an old problem with present consequences. *Malaria journal*. 2014; 13:1-10.
22. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2022;9(9): 21-6.DOI: 10.22192/ijcrps.2022.09.09.003

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

23. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. *Madonna University Journal of Medicine and Health Sciences*. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
24. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State. Nigeria. *Journal of Research in Medical and Dental Science*. 2021;9(12):462-. [links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://www.researchgate.net/publication/354117867/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf).
25. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022 Jun 21;5(2):113-121.
26. Obeagu EI. Gestational Thrombocytopaenia. *J Gynecol Women's Health*. 2023;25(3):556163. [links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf](https://www.researchgate.net/publication/368888747/Gestational-Thrombocytopaenia.pdf).
27. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(4):10-19.
28. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
29. Obeagu EI, Busari AI, Uduchi IO, Ogomaka IA, Ibekwe AM, Vincent CC, Chijioke UO, Okafor CJ, Okoroiwu HU, Adike CN. Age-Related Haematological Variations in Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):218-24.
30. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
31. Ogbonna CO, Obeagu EI, Ufelle SA, Ogbonna LN. Evaluation of haematological alterations in children infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(1):38-45.
32. Appay V. The physiological role of cytotoxic CD4+ T-cells: the holy grail? *Clinical & Experimental Immunology*. 2004;138(1):10-13.
33. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(1):25-36.
34. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med*. 2018;3(9):1-4.
35. Nwosu DC, Obeagu EI, Ezenwuba C, Agu GC, Amah H, Ozims SJ, Nwanjo HU, Edward A, Izuchukwu IF, Amadike JN, Nwagwu AJ. Antioxidant status of children with

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

- Plasmodium falciparum malaria in Owerri municipal council of Imo state. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2016;3(8):40-46.
36. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology.* 2019;41(3): e12573.
 37. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research.* 2017;5(4):206.
 38. Anyiam AF, Arinze-Anyiam OC, Omosigho PO, Ibrahim M, Ironi EA, Obeagu EI, Obi E. Blood Group, Genotype, Malaria, Blood Pressure and Blood Glucose Screening Among Selected Adults of a Community in Kwara State: Implications to Public Health. *Asian Hematology Research Journal.* 2022;6(3):9-17.
 39. Madekwe CC, Madekwe CC, Obeagu EI. Inequality of monitoring in Human Immunodeficiency Virus, Tuberculosis and Malaria: A Review. *Madonna University journal of Medicine and Health Sciences.* 2022;2(3):6-15.
 40. Offie DC, Ibekwe AM, Agu CC, Esimai BN, Okpala PU, Obeagu EI, Ufelle SA, Ogbonna LN. Fibrinogen and C-Reactive Protein Significance in Children Infected by Plasmodium falciparum Species in Enugu, Enugu State, Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(15):1-8.
 41. Obeagu EI, Ogunnaya FU. PREGNANCYINDUCED HAEMATOLOGICAL CHANGES: A KEY TO MATERNAL AND CHILD HEALTH. *European Journal of Biomedical.* 2023;10(8):42-43. [links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MATERNAL-AND-CHILD-HEALTH.pdf](https://doi.org/10.22192/ijcrms.2023.09.02.003)
 42. Obeagu EI, Ofodile AC, Okwuanaso CB. A review of urinary tract infections in pregnant women: Risks factors. *J Pub Health Nutri.* 2023; 6 (1). 2023; 137:26-35. [links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf](https://doi.org/10.22192/ijcrms.2023.09.02.003).
 43. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. *Int. J. Curr. Res. Med. Sci.* 2023;9(2): 14-17.DOI: 10.22192/ijcrms.2023.09.02.003
 44. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 14-21.DOI: 10.22192/ijcrps.2023.10.08.002
 45. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology.* 2018; 9:1733.
 46. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9): 1-4.DOI: 10.22192/ijcrbm.2018.03.09.001
 47. Onyenweaku FC, Amah HC, Obeagu EI, Nwandikor UU, Onwuasoanya UF. Prevalence of asymptomatic bacteriuria and its antibiotic susceptibility pattern in pregnant women attending private ante natal clinics in Umuahia Metropolitan. *Int J Curr Res Biol Med.* 2017;2(2): 13-23.DOI: 10.22192/ijcrbm.2017.02.02.003

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

48. Okoroiwu IL, Chinedu-Madu JU, Obeagu EI, Vincent CC, Ochiabuto OM, Ibekwe AM, Amaechi CO, Agu CC, Anoh NV, Amadi NM. Evaluation of Iron Status, Haemoglobin and Protein Levels of Pregnant Women in Owerri Metropolis. *Journal of Pharmaceutical Research International*. 2021;33(27A):36-43.
49. Obeagu EI, Njar VE, Obeagu GU. Infertility: Prevalence and Consequences. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(7):43-50.
50. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
51. Broen K, Brustoski K, Engelmann I, Luty AJ. Placental Plasmodium falciparum infection: causes and consequences of in utero sensitization to parasite antigens. *Molecular and biochemical parasitology*. 2007;151(1):1-8.
52. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9):41-49.
53. Ogbonna LN, Ezeoru VC, Ofodile AC, Ochiabuto OM, Obi-Ezeani CN, Okpala PU, Okafor CJ, Obeagu GU, Busari AI, Obeagu EI. Gender Based Variations of Haematological Parameters of Patients with Asymptomatic Malaria in Akure, Ondo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(8):75-80.
54. Eberendu IF, Ozims SJ, Agu GC, Amah HC, Obasi CC, Obioma-Elemba JE, Ihekaire DE, Ibanga IE, Amah CC, Obeagu EI, Nwosu DC. Impact of human activities on the breeding of mosquitoes of human disease in Owerri metropolis, Imo state. *Int J Adv Res Biol Sci IJARBS*. 2017;4(12):98-106.
55. Obeagu EI, Ofodile AC, Okwuanaso CB. A review on socio economic and behavioral aspects of malaria and its control among children under 5 years of age in Africa. *J Pub Health Nutri*. 2023; 6 (1): 136.
56. Djontu JC, Siewe Siewe S, Mpeke Edene YD, Nana BC, Chomga Foko EV, Bigoga JD, Leke RF, Megnekou R. Impact of placental Plasmodium falciparum malaria infection on the Cameroonian maternal and neonate's plasma levels of some cytokines known to regulate T cells differentiation and function. *Malaria journal*. 2016; 15:1-1.
57. Obeagu EI, Faduma MH, Uzoma G. Ectopic Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(4): 40-4.DOI: [10.22192/ijcrps.2023.10.04.004](https://doi.org/10.22192/ijcrps.2023.10.04.004)
58. Obeagu EI, Gamade SM, Obeagu GU. The roles of Neutrophils in pregnancy. *Int. J. Curr. Res. Med. Sci*. 2023;9(5): 31-35.DOI: [10.22192/ijcrms.2023.09.05.005](https://doi.org/10.22192/ijcrms.2023.09.05.005)
59. Obeagu EI, Obeagu GU. Molar Pregnancy: Update of prevalence and risk factors. *Int. J. Curr. Res. Med. Sci*. 2023;9(7): 25-28.DOI: [10.22192/ijcrms.2023.09.07.005](https://doi.org/10.22192/ijcrms.2023.09.07.005)
60. Kabyemela E, Gonçalves BP, Prevots DR, Morrison R, Harrington W, Gwamaka M, Kurtis JD, Fried M, Duffy PE. Cytokine profiles at birth predict malaria severity during infancy. *PloS one*. 2013;8(10):e77214.
61. Ibebuikwe JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Barriers to utilization of maternal health services in southern senatorial district of Cross Rivers state, Nigeria. *International Journal of Advanced Multidisciplinary Research*. 2017;4(8): 1-9.DOI: [10.22192/ijamr.2017.04.08.001](https://doi.org/10.22192/ijamr.2017.04.08.001)

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

62. Emmanuel G, Martin O, Peter OS, Obeagu EI, Daniel K. Factors Influencing Early Neonatal Adverse Outcomes among Women with HIV with Post Dated Pregnancies Delivering at Kampala International University Teaching Hospital, Uganda. *Asian Journal of Pregnancy and Childbirth*. 2023;6(1):203-211.
<http://research.sdpublishers.net/id/eprint/2819/>.
63. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9): 41-9.DOI: [10.22192/ijcrms.2018.04.09.006](https://doi.org/10.22192/ijcrms.2018.04.09.006)
64. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8): 22-6.DOI: [10.22192/ijcrmps.2023.10.08.003](https://doi.org/10.22192/ijcrmps.2023.10.08.003)
65. Obeagu E, Eze RI, Obeagu EI, Nnatuanya IN, Dara EC. ZINC LEVEL IN APPARENTLY PREGNANT WOMEN IN URBAN AREA. *Madonna University journal of Medicine and Health Sciences*. 2022;2(1):134-48.
<https://www.journal.madonnauniversity.edu.ng/index.php/medicine/article/view/40>.
66. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
67. Obeagu EI, Ogunnaya FU, Obeagu GU, Ndidi AC. SICKLE CELL ANAEMIA: A GESTATIONAL ENIGMA. *migration*. 2023; 17:18.
68. Harrington WE, Kakuru A, Jagannathan P. Malaria in pregnancy shapes the development of foetal and infant immunity. *Parasite immunology*. 2019;41(3):e12573.
69. Ifeanyi OE, Uzoma OG. A review on erythropietin in pregnancy. *J. Gynecol. Womens Health*. 2018;8(3):1-4.
https://www.academia.edu/download/56538560/A_Review_on_Erythropietin_in_Pregnancy.pdf.
70. Ifeanyi OE. A review on pregnancy and haematology. *Int. J. Curr. Res. Biol. Med*. 2018;3(5): 26-8.DOI: [10.22192/ijcrbm.2018.03.05.006](https://doi.org/10.22192/ijcrbm.2018.03.05.006)
71. Nwosu DC, Nwanjo HU, Obeagu EI, Ibebuike JE, Ezeama MC. Ihekireh. Changes in liver enzymes and lipid profile of pregnant women with malaria in Owerri, Nigeria. *International Journal of Current Research and Academic Review*. 2015;3(5):376-383.
72. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Factors that influence women's utilization of primary health care services in Calabar Cross river state, Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2017;4(7):28-33.
73. Elemchukwu Q, Obeagu EI, Ochei KC. Prevalence of Anaemia among Pregnant Women in Braithwaite Memorial Specialist Hospital (BMSH) Port Harcourt. *IOSR Journal of Pharmacy and Biological Sciences*. 2014;9(5):59-64.
74. Natama HM, Moncunill G, Rovira-Vallbona E, Sanz H, Sorgho H, Aguilar R, Coulibaly-Traoré M, Somé MA, Scott S, Valéa I, Mens PF. Modulation of innate immune responses at birth by prenatal malaria exposure and association with malaria risk during the first year of life. *BMC medicine*. 2018; 16:1-5.

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

75. Akandinda M, Obeagu EI, Katonera MT. Non Governmental Organizations and Women's Health Empowerment in Uganda: A Review. *Asian Research Journal of Gynaecology and Obstetrics*. 2022;8(3):12-26.
76. Gamde MS, Obeagu EI. IRON DEFICIENCY ANAEMIA: ENEMICAL TO PREGNANCY. *European Journal of Biomedical*. 2023;10(9):272-275. links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf.
77. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
78. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Studies of Some Haemostatic Variables in Preeclamptic Women in Owerri, Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(42B):39-48.
79. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2023;10(8):22-26.
80. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. *International Research in Medical and Health Sciences*. 2023;6(2):10-13.
81. Mutabingwa TK, Bolla MC, Li JL, Domingo GJ, Li X, Fried M, Duffy PE. Maternal malaria and gravidity interact to modify infant susceptibility to malaria. *PLoS medicine*. 2005;2(12):e407.
82. Gamble C, Ekwaru PJ, Garner P, Ter Kuile FO. Insecticide-treated nets for the prevention of malaria in pregnancy: a systematic review of randomised controlled trials. *PLoS medicine*. 2007;4(3):e107.
83. Okoko BJ, Enwere G, Ota MO. The epidemiology and consequences of maternal malaria: a review of immunological basis. *Acta tropica*. 2003;87(2):193-205.
84. Dobaño C, Berthoud T, Manaca MN, Nhabomba A, Guinovart C, Aguilar R, Barbosa A, Groves P, Rodríguez MH, Jimenez A, Quimice LM. High production of pro-inflammatory cytokines by maternal blood mononuclear cells is associated with reduced maternal malaria but increased cord blood infection. *Malaria Journal*. 2018; 17:1-3.
85. Umbers AJ, Stanisic DI, Ome M, Wangnapi R, Hanieh S, Unger HW, Robinson LJ, Lufele E, Baiwog F, Siba PM, King CL. Does malaria affect placental development? Evidence from in vitro models. *PLoS One*. 2013;8(1):e55269.
86. Arama C, Quin JE, Kouriba B, Östlund Farrants AK, Troye-Blomberg M, Doumbo OK. Epigenetics and malaria susceptibility/protection: A missing piece of the puzzle. *Frontiers in Immunology*. 2018; 9:1733.
87. Gbedande K, Carpio VH, Stephens R. Using two phases of the CD 4 T cell response to blood-stage murine malaria to understand regulation of systemic immunity and placental pathology in *Plasmodium falciparum* infection. *Immunological reviews*. 2020;293(1):88-114.
88. Lindsay SW, Thomas MB, Kleinschmidt I. Threats to the effectiveness of insecticide-treated bednets for malaria control: thinking beyond insecticide resistance. *The Lancet Global Health*. 2021;9(9):e1325-1331.

Citation: Obeagu EI, Obeagu GU. The Role of Placental Malaria in the Development of Childhood Malaria: A Review. *Elite Journal of Scientific Research and Review*, 2024; 2(5): 1-13

89. Akinleye SO, Falade CO, Ajayi IO. Knowledge and utilization of intermittent preventive treatment for malaria among pregnant women attending antenatal clinics in primary health care centers in rural southwest, Nigeria: a cross-sectional study. *BMC pregnancy and childbirth*. 2009; 9:1-9.