

Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes

*Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²

¹Department of Medical Laboratory Science, Kampala International University, Ishaka, Uganda.

²School of Nursing Science, Kampala International University, Ishaka, Uganda.

*Corresponding authour: Emmanuel Ifeanyi Obeagu, Department of Medical Laboratory Science, Kampala International University, Uganda.

Abstract

Blood transfusion protocols in obstetrics are pivotal in managing critical conditions like hemorrhage and severe anemia, which pose significant risks to maternal and neonatal health. This review explores current practices, challenges, and innovations in blood transfusion protocols aimed at enhancing maternal care and improving neonatal outcomes. Massive transfusion protocols (MTPs) are essential in obstetrics for managing life-threatening hemorrhage by ensuring rapid and coordinated delivery of blood products. These protocols involve predefined ratios of packed red blood cells (PRBCs), fresh frozen plasma (FFP), platelets, and clotting factors to maintain hemostasis and prevent coagulopathy. The implementation of MTPs requires multidisciplinary collaboration among healthcare providers and efficient communication with blood bank services to promptly respond to obstetric emergencies and minimize maternal morbidity and mortality. Advancements in blood product management have significantly improved the safety and efficacy of transfusions in obstetrics. Innovations include the use of leukoreduced and pathogen-reduced blood products, which reduce the risk of transfusion-related complications. Moreover, technologies such as temperature-controlled storage systems and point-of-care testing devices facilitate the timely availability of blood products, critical for managing acute hemorrhage in obstetric settings. Optimizing transfusion practices in obstetrics involves evidence-based strategies to balance the need for transfusion with minimizing risks. This includes preoperative assessment and optimization of maternal hemoglobin levels, judicious use of intraoperative cell salvage techniques, and personalized transfusion approaches based on individual patient factors.

Keywords: *Blood transfusion, obstetrics, maternal care, neonatal outcomes, massive transfusion protocols, anemia, hemorrhage*

Introduction

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

Blood transfusion is a cornerstone of obstetric care, essential for managing complications such as hemorrhage and severe anemia that threaten maternal and neonatal health. Pregnancy-related hemorrhage, particularly postpartum hemorrhage (PPH), remains a leading cause of maternal mortality worldwide, highlighting the critical importance of effective transfusion protocols. The management of obstetric hemorrhage requires prompt intervention and coordinated efforts among healthcare providers to stabilize maternal hemodynamics and prevent life-threatening complications. Blood transfusion protocols, including massive transfusion protocols (MTPs), play a pivotal role in ensuring timely access to blood products and optimizing patient outcomes.¹⁻¹⁰ Historically, obstetric hemorrhage has posed significant challenges due to its unpredictable nature and rapid onset during labor and delivery. While advances in obstetric care have reduced maternal mortality rates in many regions, disparities persist, particularly in low-resource settings where access to emergency obstetric services and blood transfusion facilities may be limited. Effective transfusion protocols are critical in these settings to mitigate the impact of hemorrhage and improve maternal survival rates. Therefore, understanding the evolution, current practices, and challenges in blood transfusion protocols in obstetrics is essential for advancing maternal healthcare globally.¹¹⁻¹² The implementation of MTPs in obstetrics represents a structured approach to managing massive blood loss during childbirth. These protocols involve the rapid administration of blood products in predefined ratios of packed red blood cells (PRBCs), fresh frozen plasma (FFP), platelets, and clotting factors to maintain hemostasis and prevent coagulopathy. Multidisciplinary collaboration among obstetricians, hematologists, anesthesiologists, and blood bank services is crucial for effective protocol implementation. This collaborative effort ensures timely access to blood products and adherence to transfusion guidelines, minimizing delays in treatment and optimizing maternal outcomes.¹³⁻¹⁷

Advancements in blood product management have significantly enhanced the safety and efficacy of transfusions in obstetrics. Leukoreduced and pathogen-reduced blood products have reduced the risk of transfusion-related complications, including febrile non-hemolytic reactions and infections. Moreover, innovations in blood storage and transport, such as temperature-controlled systems and rapid point-of-care testing devices, have improved the availability and quality of blood products in critical care settings. These advancements are particularly beneficial in managing acute hemorrhage scenarios, where rapid response is crucial for maternal survival.¹⁸⁻²⁰ Optimizing transfusion practices involves adopting evidence-based strategies to balance the benefits of transfusion with minimizing potential risks. Preoperative assessment and optimization of maternal hemoglobin levels, judicious use of intraoperative cell salvage techniques, and personalized transfusion approaches based on individual patient factors are integral components of optimal transfusion practices. These strategies not only enhance clinical outcomes but also contribute to the efficient use of blood products and resource allocation in healthcare settings.²¹⁻²³ Despite these advancements, challenges remain in the effective implementation of transfusion protocols in obstetrics, particularly in resource-limited settings. Access to adequate blood supply, infrastructure for safe blood storage and transport, and trained personnel capable of managing obstetric emergencies are critical barriers that impact maternal health outcomes. Addressing these challenges requires continued investment in healthcare infrastructure, provider training, and community-based interventions to improve access to timely and lifesaving transfusion services for pregnant women worldwide.²⁴⁻²⁷ The global burden of maternal mortality underscores the urgent

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

need for comprehensive strategies to enhance obstetric care, including effective blood transfusion protocols. Innovations in obstetric transfusion practices, such as the development of heat-stable oxytocin formulations and low-cost balloon tamponade devices, have shown promise in improving outcomes for women at risk of hemorrhage. Moreover, ongoing research in pharmacogenomics and personalized medicine holds potential for tailoring transfusion therapies to individual patient needs, further optimizing maternal healthcare delivery.²⁸⁻³¹

Massive Transfusion Protocols (MTPs) in Obstetrics

Massive transfusion protocols (MTPs) are structured guidelines designed to manage acute, life-threatening hemorrhage by rapidly delivering large volumes of blood products to stabilize hemodynamics and prevent coagulopathy. In obstetrics, where hemorrhage remains a leading cause of maternal mortality, MTPs play a crucial role in ensuring timely and coordinated interventions to optimize maternal outcomes. This article explores the principles, implementation, challenges, and advancements related to MTPs specifically in obstetric settings.³² MTPs are activated in response to severe hemorrhage scenarios where the estimated blood loss exceeds 1500 mL or when the patient shows signs of hemodynamic instability despite initial resuscitative efforts. The primary goal of MTPs is to maintain adequate tissue perfusion, hemostasis, and oxygenation by providing a balanced ratio of blood products. This typically includes packed red blood cells (PRBCs), fresh frozen plasma (FFP), platelets, and occasionally, clotting factors, administered in predefined ratios (e.g., 1:1:1 for PRBCs:FFP) to approximate the composition of whole blood and restore coagulation factors.³³ Successful implementation of MTPs in obstetrics requires a multidisciplinary approach involving obstetricians, anesthesiologists, hematologists, blood bank personnel, and nursing staff. Clear activation criteria and rapid communication protocols are essential to ensure prompt initiation of MTPs during obstetric emergencies, such as postpartum hemorrhage or complications during cesarean sections. Hospitals and birthing centers must have protocols in place for immediate availability of blood products and resources to support ongoing monitoring and adjustment of transfusion therapy based on clinical response and laboratory findings.³⁴

Several challenges complicate the implementation of MTPs in obstetrics, particularly in resource-limited settings. Access to adequate blood supply, storage facilities for blood products, and trained personnel capable of managing acute hemorrhage are critical barriers. Furthermore, logistical issues such as transportation and timely delivery of blood products to emergency settings pose significant challenges, especially in rural or underserved areas. Ensuring the safety of transfusions, including screening for infectious diseases and monitoring for transfusion reactions, adds complexity to MTP implementation.³⁶ Advancements in transfusion medicine have enhanced the effectiveness and safety of MTPs in obstetrics. The introduction of leukoreduced and irradiated blood products has reduced the incidence of transfusion-related complications, such as febrile reactions and immune-mediated responses. Pathogen reduction technologies further mitigate the risk of transmitting infections through transfusions. Innovations in blood storage, such as temperature-controlled systems and point-of-care testing devices, support the rapid availability and administration of blood products, critical for managing acute hemorrhage and improving maternal outcomes.³⁶

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

Advancements in Blood Product Management

Blood product management in medical practice, particularly in obstetrics, has undergone significant advancements aimed at enhancing safety, efficacy, and availability. These innovations are crucial in managing complications such as hemorrhage and severe anemia, which pose substantial risks to maternal and neonatal health. One major advancement in blood product management is the widespread implementation of leukoreduction and irradiation protocols. Leukoreduction involves the filtration of donor blood to remove white blood cells, which can reduce the risk of febrile non-hemolytic reactions and alloimmunization in recipients, particularly in pregnant women who may be more susceptible to immune-mediated responses. Irradiation of blood products is used to prevent transfusion-associated graft-versus-host disease (TA-GVHD), a rare but potentially fatal complication in immunocompromised patients and fetuses.³⁷ Pathogen reduction technologies represent another significant innovation aimed at improving blood safety. These methods use chemical agents or ultraviolet light to target and inactivate pathogens, including viruses, bacteria, and parasites, in blood components without compromising their therapeutic efficacy. These technologies help mitigate the risk of transfusion-transmitted infections, which remain a concern in obstetrics and other medical settings where rapid diagnosis and treatment are crucial.³⁸ Advancements in blood storage technology include the development of temperature-controlled systems that maintain the quality and viability of blood products during storage and transportation. Temperature fluctuations can affect the integrity of red blood cells, platelets, and clotting factors, potentially reducing their effectiveness upon transfusion. These systems ensure that blood products remain within optimal temperature ranges, thereby minimizing wastage and improving the availability of viable blood components for obstetric emergencies.³⁹

Point-of-care testing (POCT) devices have revolutionized blood product management by enabling rapid and accurate assessment of blood components at the bedside or in emergency settings. These portable devices measure hemoglobin levels, clotting factors, and other parameters critical for guiding transfusion decisions in real-time. In obstetrics, POCT devices facilitate immediate assessment of maternal hemodynamics and coagulation status during labor, delivery, and postpartum hemorrhage, allowing healthcare providers to tailor transfusion therapies promptly and effectively.⁴⁰ Recent advances in immunohematology and transfusion medicine have improved the accuracy and efficiency of blood compatibility testing. Techniques such as molecular typing of blood group antigens and antibody screening using automated platforms enhance the identification of compatible blood products for obstetric patients with specific immunologic profiles or previous transfusion reactions. These advancements reduce the risk of alloimmunization and hemolytic transfusion reactions, ensuring safer transfusion practices in obstetrics and minimizing adverse outcomes for both mothers and newborns.⁴¹

Optimizing Transfusion Practices in Obstetrics

Optimizing transfusion practices in obstetrics is crucial for effectively managing complications such as hemorrhage and severe anemia, which can significantly impact maternal and neonatal outcomes. A fundamental aspect of optimizing transfusion practices in obstetrics is conducting thorough preoperative assessment and optimization of maternal hemoglobin levels. This includes

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

screening for anemia early in pregnancy and implementing iron supplementation and nutritional interventions as needed to improve hemoglobin concentrations before delivery. Optimizing maternal hemoglobin levels reduces the likelihood of requiring transfusions during childbirth and minimizes the risks associated with perioperative anemia.⁴¹ Transfusion triggers refer to the hemoglobin thresholds at which transfusion is initiated based on clinical indications and patient-specific factors. In obstetrics, transfusion decisions should be guided by evidence-based guidelines and individual patient assessments rather than arbitrary hemoglobin levels alone. Factors such as ongoing bleeding, signs of hemodynamic instability, and symptomatic anemia should prompt consideration for transfusion, while recognizing that liberal transfusion practices may not necessarily improve outcomes and can introduce risks.⁴² Intraoperative techniques play a critical role in minimizing blood loss during obstetric procedures, particularly cesarean sections. Utilizing cell salvage systems, also known as intraoperative autologous blood transfusion, allows for the collection and reinfusion of a woman's own blood lost during surgery. This approach reduces the need for allogeneic transfusions and avoids potential transfusion-related complications, supporting hemostasis while preserving maternal blood volume.⁴³ Patient blood management encompasses a multidisciplinary approach to optimize transfusion practices by focusing on patient-specific strategies to minimize blood loss, maintain hemoglobin levels, and maximize tolerance of anemia. PBM strategies in obstetrics include the use of pharmacologic agents such as tranexamic acid to reduce bleeding during cesarean sections or postpartum hemorrhage, as well as implementing restrictive transfusion thresholds based on clinical evidence and patient safety profiles.⁴⁴

Personalized medicine approaches are increasingly recognized in obstetrics to tailor transfusion strategies based on individual patient factors, including obstetric history, comorbidities, and immunologic profiles. Advances in immunohematology allow for more precise matching of blood products to minimize alloimmunization risks and transfusion reactions. Personalized transfusion strategies ensure that obstetric patients receive the most appropriate blood components while minimizing potential adverse effects associated with transfusion therapy.⁴⁵ Effective implementation of optimized transfusion practices in obstetrics requires ongoing education and training for healthcare providers. Obstetricians, anesthesiologists, midwives, and nursing staff should be well-versed in evidence-based transfusion guidelines, MTP protocols, and the use of transfusion alternatives to ensure consistent and safe practice. Training programs should emphasize the importance of multidisciplinary teamwork, communication strategies during obstetric emergencies, and the proper handling and administration of blood products to enhance patient care and outcomes.⁴⁶ Integration of technological advancements, such as point-of-care testing devices for rapid hemoglobin assessment and blood product compatibility testing, enhances the efficiency and safety of transfusion practices in obstetrics. Quality improvement initiatives, including regular audits of transfusion practices and outcomes, facilitate continuous monitoring and optimization of transfusion protocols. These initiatives promote adherence to best practices, identify areas for improvement, and ultimately contribute to reducing unnecessary transfusions and improving patient safety.⁴⁷⁻⁵⁵

Challenges

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

One of the primary challenges in obstetric transfusion practices is ensuring access to an adequate and safe blood supply, especially in resource-limited settings. Blood shortages, variability in blood product availability, and logistical challenges in transporting blood to remote areas can delay lifesaving interventions during obstetric emergencies.⁵⁶⁻⁵⁸ While blood transfusion is a life-saving intervention, it is not without risks. Transfusion-related complications such as transfusion-transmitted infections, hemolytic reactions, and immunological complications remain concerns, necessitating stringent screening, testing, and monitoring protocols to ensure blood product safety.⁵⁹⁻⁶³ Determining the optimal timing and volume of transfusions in obstetrics can be complex. Factors such as patient-specific risks, hemodynamic stability, and laboratory parameters must be carefully considered to avoid under-transfusion leading to inadequate tissue perfusion or over-transfusion associated with increased risks of complications.⁶⁴⁻⁶⁷ Adequate training and education of healthcare providers in obstetric transfusion practices are essential for ensuring adherence to guidelines, safe administration of blood products, and effective management of transfusion-related complications. Continuous professional development and simulation training can enhance preparedness and improve outcomes during obstetric emergencies.⁶⁸⁻⁷⁰ Ethical dilemmas related to blood transfusion, such as religious or cultural beliefs that may influence acceptance or refusal of transfusion, can complicate decision-making in obstetrics. Respectful communication and collaboration with patients and families are essential to navigating these sensitive issues while prioritizing maternal and fetal health.

Future Directions

Future efforts should focus on improving blood collection, processing, and distribution systems to ensure consistent availability of safe blood products, particularly in underserved regions. This may involve expanding blood donor recruitment programs, implementing innovative storage solutions, and strengthening blood transfusion services infrastructure.⁷¹ Continued research into pathogen reduction technologies and the development of novel blood products with extended shelf life and enhanced safety profiles can further mitigate transfusion-related risks. Innovations in blood banking practices and regulatory frameworks can support the implementation of these advancements in clinical practice.⁷² Advances in molecular and genetic testing allow for personalized transfusion strategies based on individual patient factors, including blood group compatibility, immunologic profiles, and transfusion history. Tailoring transfusion therapies to meet specific patient needs can optimize outcomes while minimizing risks and resource utilization.⁷³ Integration of digital health technologies, such as electronic decision support systems and telemedicine platforms, can enhance real-time monitoring of transfusion practices, facilitate remote consultation during obstetric emergencies, and support data-driven quality improvement initiatives. Collaborative efforts among healthcare organizations, professional societies, and governmental agencies are essential for promoting standardized transfusion guidelines, sharing best practices, and building capacity in obstetric transfusion practices worldwide. Educational initiatives and training programs should prioritize transfusion safety, clinical decision-making, and multidisciplinary teamwork.⁷⁴

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. *Elite Journal of Nursing and Health Science*, 2024; 2(6):14-25

Conclusion

The management of blood transfusion practices in obstetrics is pivotal for ensuring maternal and neonatal safety during pregnancy and childbirth. As highlighted throughout this discussion, optimizing transfusion strategies involves navigating various challenges while embracing advancements and future opportunities in healthcare. Effective transfusion practices in obstetrics require a balanced approach that integrates evidence-based guidelines, multidisciplinary collaboration, and continuous quality improvement initiatives. Key principles include preoperative assessment and optimization of maternal hemoglobin levels, judicious use of transfusion triggers based on clinical indications, and the implementation of massive transfusion protocols (MTPs) for severe hemorrhage scenarios. These practices not only aim to stabilize maternal hemodynamics and prevent complications but also emphasize the importance of personalized medicine approaches to tailor transfusion therapies to individual patient needs.

References

1. Smith C, Teng F, Branch E, Chu S, Joseph KS. Maternal and perinatal morbidity and mortality associated with anemia in pregnancy. *Obstetrics & Gynecology*. 2019;134(6):1234-1244.
2. Shi H, Chen L, Wang Y, Sun M, Guo Y, Ma S, Wang X, Jiang H, Wang X, Lu J, Ge L. Severity of anemia during pregnancy and adverse maternal and fetal outcomes. *JAMA network open*. 2022;5(2):e2147046-.
3. Harrison RK, Lauhon SR, Colvin ZA, McIntosh JJ. Maternal anemia and severe maternal morbidity in a US cohort. *American journal of obstetrics & gynecology MFM*. 2021;3(5):100395.
4. Beckert RH, Baer RJ, Anderson JG, Jelliffe-Pawlowski LL, Rogers EE. Maternal anemia and pregnancy outcomes: a population-based study. *Journal of Perinatology*. 2019;39(7):911-919.
5. Daru J, Zamora J, Fernández-Félix BM, Vogel J, Oladapo OT, Morisaki N, Tunçalp Ö, Torloni MR, Mittal S, Jayaratne K, Lumbiganon P. Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: a multilevel analysis. *The Lancet Global Health*. 2018;6(5):e548-554.
6. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC, Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria, *J Res Med Dent Sci*, 2021, 9 (4):145-148
7. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri*. 2023; 6 (1). 2023;138. [links/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf](https://doi.org/10.3390/63da799664fc860638054562/Anaemia-among-pregnant-women-A-review-of-African-pregnant-teenagers.pdf).
8. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci*. 2016;3(3):10-8. [links/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf](https://doi.org/10.3390/5710fae108ae846f4ef05afb/ERYTHROPOIETIN-IN-THE-ANAEMIAS-OF-PREGNANCY-A-REVIEW.pdf).

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. *Elite Journal of Nursing and Health Science*, 2024; 2(6):14-25

9. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. J Res Med Dent Sci. 2021;9(4):145-8. [links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf](https://epjournals.com/journals/EJNHS/links/608a6728a6fdccaebdf52d94/Assessment-of-Haematological-Changes-in-Pregnant-Women-of-Ido-Ondo.pdf).
10. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. International Research in Medical and Health Sciences. 2023 ;6(2):10-3. <http://irmhs.com/index.php/irmhs/article/view/111>.
11. Georgieff MK. Iron deficiency in pregnancy. American journal of obstetrics and gynecology. 2020;223(4):516-524.
12. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of Protein C, Protein S and Fibrinogen of Pregnant Women with Malaria in Owerri Metropolis. Madonna University journal of Medicine and Health Sciences. 2022;2(2):1-9.
13. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. Madonna University Journal of Medicine and Health Sciences. 2022;2(1):292-8. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/57>.
14. Obeagu EI, Obeagu GU, Adepoju OJ. Evaluation of haematological parameters of pregnant women based on age groups in Olorunsogo road area of Ido, Ondo state. J. Bio. Innov11 (3). 2022:936-941.
15. Cantor AG, Bougatsos C, Dana T, Blazina I, McDonagh M. Routine iron supplementation and screening for iron deficiency anemia in pregnancy: a systematic review for the US Preventive Services Task Force. Annals of internal medicine. 2015;162(8):566-576.
16. Siu AL, US Preventive Services Task Force*. Screening for iron deficiency anemia and iron supplementation in pregnant women to improve maternal health and birth outcomes: US Preventive Services Task Force recommendation statement. Annals of Internal Medicine. 2015;163(7):529-536.
17. Obeagu EI. An update on utilization of antenatal care among pregnant Women in Nigeria. Int. J. Curr. Res. Chem. Pharm. Sci. 2022;9(9): 21-6.DOI: [10.22192/ijcrps.2022.09.09.003](https://doi.org/10.22192/ijcrps.2022.09.09.003)
18. Okoroiwu IL, Obeagu EI, Obeagu GU. Determination of clot retraction in pregnant women attending antenatal clinic in federal medical centre Owerri, Nigeria. Madonna University Journal of Medicine and Health Sciences. 2022;2(2):91-97. <https://madonnauniversity.edu.ng/journals/index.php/medicine/article/view/67>.
19. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State, Nigeria. Journal of Research in Medical and Dental Science. 2021;9(12):462-.[links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf](https://epjournals.com/journals/EJNHS/links/61b1e32f0c4bfb675178bfa7/Evaluation-of-Changes-in-Haematological-Parameters-of-Pregnant-Women-Based-on-Gestational-Age-at-Olorunsogo-Road-Area-of-Ido-Ondo-State-Nigeria.pdf).
20. Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Ironi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. International Journal of Research and Reports in Hematology. 2022 Jun 21;5(2):113-121.

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

21. Obeagu EI. Gestational Thrombocytopaenia. J Gynecol Women's Health. 2023;25(3):556163. <links/64b01aa88de7ed28ba95fccb/Gestational-Thrombocytopaenia.pdf>.
22. Liu CN, Yu FB, Xu YZ, Li JS, Guan ZH, Sun MN, Liu CA, He F, Chen DJ. Prevalence and risk factors of severe postpartum hemorrhage: a retrospective cohort study. BMC pregnancy and childbirth. 2021;21(1):332.
23. Obeagu EI, Ogbonna US, Nwachukwu AC, Ochiabuto O, Enweani IB, Ezeoru VC. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. Journal of Pharmaceutical Research International. 2021;33(4):10-9.
24. Obeagu EI, Abdirahman BF, Bunu UO, Obeagu GU. Obstetrics characteristics that effect the newborn outcomes. Int. J. Adv. Res. Biol. Sci. 2023;10(3): 134-43.DOI: <10.22192/ijarbs.2023.10.03.016>
25. Obeagu EI, Ogunnaya FU. PREGNANCYINDUCED HAEMATOLOGICAL CHANGES: A KEY TO MARTERNAL AND CHILD HEALTH. European Journal of Biomedical. 2023;10(8):42-43. <links/64c890bddb38b20d6dad2c5c/PREGNANCY-INDUCED-HAEMATOLOGICAL-CHANGES-A-KEY-TO-MARTERNAL-AND-CHILD-HEALTH.pdf>.
26. Ezeoru VC, Enweani IB, Ochiabuto O, Nwachukwu AC, Ogbonna US, Obeagu EI. Prevalence of Malaria with Anaemia and HIV status in women of reproductive age in Onitsha, Nigeria. Journal of Pharmaceutical Research International. 2021;33(4):10-19.
27. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC, Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. Annals of Clinical and Laboratory Research. 2017;5(4):206. <links/5ea97df145851592d6a8acf2/Iron-Status-of-Pregnant-and-Post-Partum-Women-with-Malaria-Parasitaemia-in-Aba-Abia-State-Nigeria.pdf>.
28. Eze RI, Obeagu EI, Edet FN. Frequency of Rh Antigen C And c among pregnant women in Sub-Urban area in Eastern Nigeria. Madonna Uni J Med Health Sci. 2021;1(1):19-30.
29. Obeagu EI, Ofodile AC, Okwuanaso CB. A review of urinary tract infections in pregnant women: Risks factors. J Pub Health Nutri. 2023; 6 (1). 2023; 137:26-35. <links/63c3a9116fe15d6a571e8bba/A-review-of-urinary-tract-infections-in-pregnant-women-Risks-factors.pdf>.
30. Obeagu EI, Obeagu GU, Musiimenta E. Post partum haemorrhage among pregnant women: Update on risks factors. Int. J. Curr. Res. Med. Sci. 2023;9(2): 14-17.DOI: <10.22192/ijcrms.2023.09.02.003>
31. Obeagu EI, Obeagu GU, Ogunnaya FU. Deep vein thrombosis in pregnancy: A review of prevalence and risk factors. Int. J. Curr. Res. Chem. Pharm. Sci. 2023;10(8): 14-21.DOI: <10.22192/ijcrmps.2023.10.08.002>
32. Pacheco LD, Saade GR, Costantine MM, Clark SL, Hankins GD. An update on the use of massive transfusion protocols in obstetrics. American journal of obstetrics and gynecology. 2016;214(3):340-344.
33. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. Int. J. Curr. Res. Biol. Med. 2018;3(9): 1-4.DOI: <10.22192/ijcrbm.2018.03.09.001>

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. Elite Journal of Nursing and Health Science, 2024; 2(6):14-25

34. Onyenweaku FC, Amah HC, Obeagu EI, Nwandikor UU, Onwuasoanya UF. Prevalence of asymptomatic bacteriuria and its antibiotic susceptibility pattern in pregnant women attending private ante natal clinics in Umuahia Metropolitan. *Int J Curr Res Biol Med.* 2017;2(2): 13-23.DOI: [10.22192/ijcrbm.2017.02.02.003](https://doi.org/10.22192/ijcrbm.2017.02.02.003)
35. Okoroiwu IL, Chinedu-Madu JU, Obeagu EI, Vincent CC, Ochiabuto OM, Ibekwe AM, Amaechi CO, Agu CC, Anoh NV, Amadi NM. Evaluation of Iron Status, Haemoglobin and Protein Levels of Pregnant Women in Owerri Metropolis. *Journal of Pharmaceutical Research International.* 2021;33(27A):36-43.
36. Obeagu EI, Njar VE, Obeagu GU. Infertility: Prevalence and Consequences. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(7):43-50.
37. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International.* 2021;33(42A):53-65.
38. Carson JL, Guyatt G, Heddle NM, Grossman BJ, Cohn CS, Fung MK, Gernsheimer T, Holcomb JB, Kaplan LJ, Katz LM, Peterson N. Clinical practice guidelines from the AABB: red blood cell transfusion thresholds and storage. *Jama.* 2016;316(19):2025-2035.
39. Obeagu EI, Faduma MH, Uzoma G. Ectopic Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(4): 40-4.DOI: [10.22192/ijcrmps.2023.10.04.004](https://doi.org/10.22192/ijcrmps.2023.10.04.004)
40. Obeagu EI, Gamade SM, Obeagu GU. The roles of Neutrophils in pregnancy. *Int. J. Curr. Res. Med. Sci.* 2023;9(5): 31-35.DOI: [10.22192/ijcrms.2023.09.05.005](https://doi.org/10.22192/ijcrms.2023.09.05.005)
41. Eze R, Obeagu EI, Nwakulite A, Okoroiwu IL, Vincent CC, Okafor CJ, Chukwurah EF, Chijioke UO, Amaechi CO. Evaluation of Copper Status and Some Red Cell Parameters of Pregnant Women in Enugu State, South Eastern Nigeria. *Journal of Pharmaceutical Research International.* 2021;33(30A):67-71.
42. Obeagu EI, Obeagu GU. Molar Pregnancy: Update of prevalence and risk factors. *Int. J. Curr. Res. Med. Sci.* 2023;9(7): 25-28.DOI: [10.22192/ijcrms.2023.09.07.005](https://doi.org/10.22192/ijcrms.2023.09.07.005)
43. Obeagu EI, Bunu UO. Factors that influence unmet need for family planning. *International Journal of Current Research in Biology and Medicine.* 2023;8(1):23-27.
44. Ibebuike JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Barriers to utilization of maternal health services in southern senatorial district of Cross Rivers state, Nigeria. *International Journal of Advanced Multidisciplinary Research.* 2017;4(8): 1-9.DOI: [10.22192/ijamr.2017.04.08.001](https://doi.org/10.22192/ijamr.2017.04.08.001)
45. Emmanuel G, Martin O, Peter OS, Obeagu EI, Daniel K. Factors Influencing Early Neonatal Adverse Outcomes among Women with HIV with Post Dated Pregnancies Delivering at Kampala International University Teaching Hospital, Uganda. *Asian Journal of Pregnancy and Childbirth.* 2023;6(1):203-211. <http://research.sdpublishers.net/id/eprint/2819/>.
46. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences.* 2018;4(9): 41-9.DOI: [10.22192/ijcrms.2018.04.09.006](https://doi.org/10.22192/ijcrms.2018.04.09.006)
47. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. *Int. J. Curr. Res. Chem. Pharm. Sci.* 2023;10(8): 22-6.DOI: [10.22192/ijcrmps.2023.10.08.003](https://doi.org/10.22192/ijcrmps.2023.10.08.003)

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. *Elite Journal of Nursing and Health Science*, 2024; 2(6):14-25

48. Obeagu E, Eze RI, Obeagu EI, Nnatuanya IN, Dara EC. ZINC LEVEL IN APPARENTLY PREGNANT WOMEN IN URBAN AREA. *Madonna University journal of Medicine and Health Sciences*. 2022;2(1):134-48.
<https://www.journal.madonnauniversity.edu.ng/index.php/medicine/article/view/40>.
49. Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, Ota E, Gilmour S, Shibuya K. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis. *The American journal of clinical nutrition*. 2016;103(2):495-504.
50. Ogomaka IA, Obeagu EI. Malaria in Pregnancy Amidst Possession of Insecticide Treated Bed Nets (ITNs) in Orlu LGA of Imo State, Nigeria. *Journal of Pharmaceutical Research International*. 2021;33(41B):380-386.
51. Obeagu EI, Ogunnaya FU, Obeagu GU, Ndidi AC. SICKLE CELL ANAEMIA: A GESTATIONAL ENIGMA. *migration*. 2023; 17:18.
52. Ifeanyi OE, Uzoma OG. A review on erythropietin in pregnancy. *J. Gynecol. Womens Health*. 2018;8(3):1-4.
https://www.academia.edu/download/56538560/A_Review_on_Erythropietin_in_Pregnancy.pdf.
53. Ifeanyi OE. A review on pregnancy and haematology. *Int. J. Curr. Res. Biol. Med*. 2018;3(5): 26-8.DOI: [10.22192/ijcrbm.2018.03.05.006](https://doi.org/10.22192/ijcrbm.2018.03.05.006)
54. Nwosu DC, Nwanjo HU, Obeagu EI, Ibebuikie JE, Ezeama MC. Ihekireh. Changes in liver enzymes and lipid profile of pregnant women with malaria in Owerri, Nigeria. *International Journal of Current Research and Academic Review*. 2015;3(5):376-383.
55. Ibebuikie JE, Ojie CA, Nwokike GI, Obeagu EI, Nwosu DC, Nwanjo HU, Agu GC, Ezenwuba CO, Nwagu SA, Akujuobi AU. Factors that influence women's utilization of primary health care services in Calabar Cross river state, Nigeria. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2017;4(7):28-33.
56. Eze R, Ezeah GA, Obeagu EI, Omeje C, Nwakulite A. Evaluation of iron status and some haematological parameters of pregnant women in Enugu, South Eastern Nigeria. *World Journal of Pharmaceutical and Medical Research*. 2021;7(5):251-4.
57. Elemchukwu Q, Obeagu EI, Ochei KC. Prevalence of Anaemia among Pregnant Women in Braithwaite Memorial Specialist Hospital (BMSH) Port Harcourt. *IOSR Journal of Pharmacy and Biological Sciences*. 2014;9(5):59-64.
58. Akandinda M, Obeagu EI, Katonera MT. Non Governmental Organizations and Women's Health Empowerment in Uganda: A Review. *Asian Research Journal of Gynaecology and Obstetrics*. 2022;8(3):12-26.
59. Tandon R, Jain A, Malhotra P. Management of iron deficiency anemia in pregnancy in India. *Indian Journal of Hematology and Blood Transfusion*. 2018; 34:204-215.
60. Montoro M, Cucala M, Lanas Á, Villanueva C, Hervás AJ, Alcedo J, Gisbert JP, Aisa AP, Bujanda L, Calvet X, Mearin F. Indications and hemoglobin thresholds for red blood cell transfusion and iron replacement in adults with gastrointestinal bleeding: An algorithm proposed by gastroenterologists and patient blood management experts. *Frontiers in Medicine*. 2022; 9:903739.
61. Kapur D, Nath Agarwal K, Kumari Agarwal D. Nutritional anemia and its control. *The Indian Journal of Pediatrics*. 2002; 69:607-616.

Citation: Obeagu EI, Obeagu GU. Blood Transfusion Protocols in Obstetrics: Enhancing Maternal Care and Neonatal Outcomes. *Elite Journal of Nursing and Health Science*, 2024; 2(6):14-25

62. Gamde MS, Obeagu EI. IRON DEFICIENCY ANAEMIA: ENEMICAL TO PREGNANCY. European Journal of Biomedical. 2023;10(9):272-275. [links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf](https://epjournals.com/journals/EJNHS/links/64f63358827074313ffaae7b/IRON-DEFICIENCY-ANAEMIA-ENEMICAL-TO-PREGNANCY.pdf).
63. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. Journal of Pharmaceutical Research International. 2021;33(42A):53-65.
64. Hasan MM, Magalhaes RJ, Garnett SP, Fatima Y, Tariqujjaman M, Pervin S, Ahmed S, Mamun AA. Anaemia in women of reproductive age in low-and middle-income countries: progress towards the 2025 global nutrition target. Bulletin of the World Health Organization. 2022;100(3):196.
65. Paulman PM, Paulman A, Harrison JD, Nasir LS, Bryan SK, Collier DS. Signs and Symptoms in Family Medicine E-Book: A Literature-Based Approach. Elsevier Health Sciences; 2011.
66. Stoffel NU, von Siebenthal HK, Moretti D, Zimmermann MB. Oral iron supplementation in iron-deficient women: How much and how often? Molecular aspects of medicine. 2020; 75:100865.
67. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Studies of Some Haemostatic Variables in Preeclamptic Women in Owerri, Imo State, Nigeria. Journal of Pharmaceutical Research International. 2021;33(42B):39-48.
68. Obeagu EI, Obeagu GU. Postpartum haemorrhage among women delivering through spontaneous vaginal delivery: Prevalence and risk factors. Int. J. Curr. Res. Chem. Pharm. Sci. 2023;10(8):22-26.
69. Obeagu EI, Obeagu GU. Sick Cell Anaemia in Pregnancy: A Review. International Research in Medical and Health Sciences. 2023;6(2):10-13.
70. Al RA, Unlubilgin E, Kandemir O, Yalvac S, Cakir L, Haberal A. Intravenous versus oral iron for treatment of anemia in pregnancy: a randomized trial. Obstetrics & Gynecology. 2005;106(6):1335-1340.
71. Ngwenya S. Postpartum hemorrhage: incidence, risk factors, and outcomes in a low-resource setting. International journal of women's health. 2016:647-650.
72. Obeagu EI, Ubosi NI, Uzoma G. Maternal Hemorrhage and Blood Transfusions: Safeguarding Pregnancy Health. Int. J. Curr. Res. Chem. Pharm. Sci. 2023;10(11):26-35.
73. Obeagu EI, Obeagu GU. The Vital Role of Blood Transfusions during Pregnancy: A Comprehensive Review. Asian Journal of Dental and Health Sciences. 2024;4(1):26-31.
74. Obeagu EI, Obeagu GU. Addressing Anemia and Hemorrhage in Pregnancy: Blood Transfusion Strategies for Improved Maternal Health. *Elite Journal of Haematology*, 2024; 2(6): 46-59