Homework 1.1: Can an AI Rediscover Kepler's Orbit of Mars?

Mathematical Foundations of Generative AI

Project Goal

The goal of this project was to use a large language model (LLM) as a scientific collaborator rather than as an answer generator. The project built a synthetic version of the observational problem faced by Kepler: generate noisy measurements of Mars from Uraniborg, reconstruct heliocentric Mars positions without using the hidden Mars orbit parameters, fit competing geometric models, and then judge whether the data support a circle or an ellipse.

The workflow was:

1. Generate synthetic Tycho-like observations of Mars from 1582 to 1600.
2. Independently verify the noiseless orbital model using conservation of energy and angular momentum.
3. Plot Mars in the apparent sky using right ascension/declination and altitude/azimuth.
4. Convert angular measurements to geometric lines of sight and reconstruct heliocentric Mars positions.
5. Plot the reconstructed Mars positions together with the Sun and Earth's orbit.
6. Fit a Sun-centered circle, an arbitrary-center circle, and a Kepler ellipse.
7. Repeat the analysis for 1, 2, and 5 arcminutes of observational noise.
8. Ask the LLM to interpret the numerical evidence and critique its reasoning.
9. Discuss the connection between Kepler's work and modern model fitting from data.

---

1. Synthetic Observation Table

Three synthetic observation tables were generated, one for each noise level:

- 1 arcminute observations
- 2 arcminute observations
- 5 arcminute observations

Each file contains 166 accepted observations. The columns are:

- `datetime_utc` : observation date and time
- `ra_deg` : right ascension in degrees
- `dec_deg` : declination in degrees

- `alt_deg` : altitude above the Uraniborg horizon in degrees
- `az_deg` : azimuth measured clockwise from north in degrees
- `noise_arcmin` : angular noise level used for that table

The observations were generated every 10 days at 22:00 UTC between 1582 and 1600, but an observation was kept only when Mars was at least 10 degrees above the horizon and the Sun was at least 6 degrees below the horizon. This prevents the synthetic observer from recording Mars when it would be below the horizon or in bright daylight.

A sample from the 1 arcminute table is shown below. The complete table is included in the submitted CSV file.

Date (UTC)	RA (deg)	Dec (deg)	Altitude (deg)	Azimuth (deg)	Noise
1582-10-18 22:00	113.1434	21.7463	10.6741	66.7871	1'
1582-10-28 22:00	116.7729	21.1738	13.4672	72.0128	1'
1582-11-07 22:00	119.5004	20.6809	16.9230	77.9334	1'
1582-11-17 22:00	121.1073	20.3617	21.2341	84.7275	1'
1582-11-27 22:00	121.4645	20.2909	26.4962	92.5909	1'
1582-12-07 22:00	120.3862	20.5386	32.7699	101.9267	1'
1582-12-17 22:00	117.8709	20.9645	39.7199	113.4277	1'
1582-12-27 22:00	114.1580	21.5882	46.7929	127.8897	1'
1583-01-06 22:00	109.9395	22.1667	52.7118	146.0109	1'
1583-01-16 22:00	106.0045	22.6162	56.2296	167.2120	1'
1583-01-26 22:00	103.1156	22.9017	56.7807	188.5722	1'
1583-02-05 22:00	101.5174	22.9913	54.8962	207.1125	1'

The full 1 arcminute data range was approximately:

- RA: 0.89 to 355.91 degrees
- Declination: -23.14 to 23.44 degrees
- Altitude: 10.25 to 57.48 degrees
- Azimuth: 66.03 to 294.84 degrees

The 2 and 5 arcminute tables contain the same underlying observations with independently perturbed angular directions.

---

2. Hidden Orbital Model Used to Generate the Data

The synthetic model is intentionally simplified. The Sun is fixed at the origin, and Earth and Mars move on fixed Kepler ellipses in a common ecliptic plane. Planetary perturbations, precession, and a full historical ephemeris are not included. The purpose is to create a controlled data-fitting problem rather than a high-precision astronomical simulator.

The observer is located at Uraniborg using the coordinates

- latitude: 55 degrees 54 minutes 28 seconds N
- longitude: 12 degrees 41 minutes 48 seconds E

The hidden orbital parameters used only in data generation were:

Planet	Semi-major axis a (AU)	Eccentricity e	Initial mean anomaly M0	Perihelion direction
Earth	1.00000011	0.01671022	357.51716 deg	102.93735 deg
Mars	1.52366231	0.09341233	19.41248 deg	336.04084 deg

The solar gravitational parameter was expressed in AU and days as

$$\mu = \frac{4\pi^2}{(365.256363004)^2}$$

For a planet with semi-major axis a, the mean angular rate is

$$n = \sqrt{\frac{\mu}{a^3}}$$

and the mean anomaly is

$$M(t) = M_0 + nt$$

The eccentric anomaly E is obtained by solving Kepler's equation

$$E - e\sin(E) = M$$

Newton's method was used to solve this nonlinear equation. Once E is known, the position in the orbital plane is

$$x = a(\cos E - e)$$

$$y = a\sqrt{1 - e^2}\sin E$$

The velocity is obtained by differentiating Kepler's equation. Since

$$\frac{dE}{dt} = \frac{n}{1 - e\cos E}$$

the velocity components are

$$v_x = -a \sin E \frac{dE}{dt}$$

$$v_y = a\sqrt{1-e^2} \cos E \frac{dE}{dt}$$

The orbital plane is rotated by the chosen perihelion direction.

Observer geometry

Earth's axial tilt was taken to be 23.43928 degrees. The heliocentric planet positions were converted from ecliptic coordinates to equatorial coordinates. Earth's daily rotation was represented using an approximate Greenwich mean sidereal time calculation. Uraniborg's longitude shifts the sidereal angle to the local angle at the observatory, while its latitude determines the local east, north, and up basis.

The observer's heliocentric position is

$$\mathbf{r}_{\text{obs}} = \mathbf{r}_E + \mathbf{r}_{\text{site}}$$

The direction from Uraniborg to Mars is

$$\boldsymbol{\rho} = \mathbf{r}_M - \mathbf{r}_{\text{obs}}$$

After normalization, this unit vector is converted to right ascension/declination and altitude/azimuth.

Observational noise

Noise was added directly to the unit line-of-sight vector in its local tangent plane. This is more physically meaningful than independently adding random values to RA and declination because it represents a small random pointing error on the celestial sphere. Three Gaussian angular noise scales were tested:

- 1 arcminute
- 2 arcminutes
- 5 arcminutes

The synthetic-data generation code is contained in synthetic_data_generation(step_1).ipynb.

---

3. Verification of the Noiseless Orbital Simulation

The orbital code was checked independently rather than accepted because it was generated

with LLM assistance. Two conserved quantities were tested: specific orbital energy and specific angular momentum.

For a Kepler orbit, the specific orbital energy is

$$\mathcal{E} = \frac{1}{2}\|\mathbf{v}\|^2 - \frac{\mu}{\|\mathbf{r}\|}$$

and the exact theoretical value is

$$\mathcal{E}_{\text{theory}} = -\frac{\mu}{2a}$$

The specific angular momentum is

$$h = \|\mathbf{r} \times \mathbf{v}\|$$

with theoretical value

$$h_{\text{theory}} = \sqrt{\mu a(1 - e^2)}$$

The independent verification script evaluated the Earth and Mars orbits from 1582 through 1600 and compared the computed invariants with these closed-form values.


Verification summary

Planet	Period (days)	Perihelion (AU)	Aphelion (AU)	Max relative energy error	Energy relative variation	Max relative angular momentum error	Angular momentum relative variation
Earth	365.2564	0.983290	1.016710	1.28e-15	2.38e-15	6.05e-16	1.01e-15
Mars	686.9591	1.381333	1.665991	1.40e-15	2.51e-15	8.21e-16	1.15e-15


These errors are at the scale of floating-point roundoff. Therefore the position and velocity formulas are internally consistent with the theoretical Kepler invariants.

One limitation of this check is important: the orbit is generated analytically from Kepler's equation rather than integrated forward with a numerical ODE solver. Therefore near-perfect conservation is expected. The verification confirms that the formulas used for position and velocity are mutually consistent; it is not a test of long-time numerical integration stability.


Earth energy conservation




Earth angular momentum conservation



Mars energy conservation



Mars angular momentum conservation



The complete numerical verification table is stored in orbit_verification_summary.csv.

---


4. Apparent Position of Mars in Tycho's Sky

The synthetic observations were plotted in two coordinate systems.


Right ascension and declination describe the direction of Mars on the celestial sphere. Altitude and azimuth describe what an observer at Uraniborg would see relative to the local horizon. The latter coordinates make the effect of Earth's rotation and Uraniborg's geographic location especially clear.

Right ascension and declination


1 arcminute noise



2 arcminutes noise




5 arcminutes noise




The apparent path is not expected to look like a simple ellipse because the observer is moving with Earth. The plotted direction depends on the relative vector between Mars and the observer, not simply on Mars's heliocentric coordinates. Apparent reversals and loops are therefore natural in the sky coordinates.

Altitude and azimuth


1 arcminute noise



2 arcminutes noise



5 arcminutes noise



The lower boundary near 10 degrees altitude is a direct consequence of the visibility condition used during data generation.

The three noise levels look visually similar because even 5 arcminutes is only about 0.083 degrees, much smaller than the overall angular scale of the sky plots.

---

5. Reconstruction of Heliocentric Mars Positions

A single RA/declination observation provides a direction but not a distance. If $\mathbf{p}_i$ is the heliocentric observer position and $\mathbf{u}_i$ is the observed unit direction, then Mars must lie somewhere on the ray

$$\mathbf{r} = \mathbf{p}_i + \rho_i \mathbf{u}_i$$

where the range ρ_i is unknown.

Corrected multi-ray reconstruction

The final reconstruction did not use the hidden Mars orbital parameters. It used only:

- the noisy RA/declination observations,
- Earth's known orbit,
- Uraniborg's location,
- Earth's rotation.

The method searched candidate repetition periods between 500 and 900 days. Observations separated by approximately one or more candidate periods were grouped under the assumption that Mars should return near the same orbital phase. At least three rays were required for each reconstructed point.

For a group of rays $\mathbf{p}_i + \rho_i \mathbf{u}_i$, the common point $\mathbf{x}$ was found by minimizing the total squared perpendicular distance to the rays. In matrix form, each ray contributes the projection matrix

$$I - \mathbf{u}_i \mathbf{u}_i^T$$

The least-squares system is

$$A\mathbf{x} = \mathbf{b}$$

where

$$A = \sum_i \left(I - \mathbf{u}_i \mathbf{u}_i^T \right)$$

$$\mathbf{b} = \sum_i \left(I - \mathbf{u}_i \mathbf{u}_i^T \right) \mathbf{p}_i$$

To avoid poorly conditioned triangulation, the observer positions were required to have a heliocentric baseline of at least 0.30 AU.


The best inferred repetition period was 689.25 days for all three noise levels. The hidden Mars period used to generate the data was 686.959 days, so the inferred period differs by only about 2.29 days, or approximately 0.33 percent.

Reconstruction statistics


Noise	Reconstructed points	Inferred period (days)	Minimum radius (AU)	Median radius (AU)	Maximum radius (AU)	Median ray RMS (AU)	Median absolute z (AU)
1'	26	689.25	1.4268	1.5387	1.6072	0.01235	0.000134
2'	26	689.25	1.4262	1.5380	1.6071	0.01236	0.000267
5'	26	689.25	1.4245	1.5359	1.6070	0.01242	0.000667

The hidden model places Mars between approximately 1.381 and 1.666 AU from the Sun. The reconstructed radii are therefore physically reasonable. The reconstructed z coordinates also remain very close to zero, as expected because the hidden simplified model places the planets in a common ecliptic plane.


Period search, 1 arcminute




Reconstructed Mars positions, 1 arcminute



Reconstructed Mars positions, 2 arcminutes



Reconstructed Mars positions, 5 arcminutes



Post-hoc verification against the hidden orbit

The hidden Mars parameters were not used during reconstruction. After the reconstruction was finished, they were used only as a verification check. The reconstructed 1 arcminute points follow the same orbital scale, orientation, and curvature as the hidden generating ellipse.

This check is important because an earlier LLM-suggested two-ray method produced an unphysical reconstruction near 1 AU, almost directly on Earth's orbit. That failure was detected by comparing the reconstructed heliocentric distance with the physically expected Mars scale. The method was then replaced with the multi-ray least-squares reconstruction above.

---

6. Circle and Ellipse Fits with Residual Errors

Three models were fitted to each reconstructed data set.

Model 1: Sun-centered circle

The model is

$$x^2 + y^2 = R^2$$

Only the radius R is fitted.

Model 2: Free-center circle

The model is

$$(x - h)^2 + (y - k)^2 = R^2$$

The center (h, k) and radius R are fitted.

Model 3: Kepler ellipse with the Sun at one focus

The polar form is

$$r(\theta) = \frac{p}{1 + e\cos(\theta - \omega)}$$

The fitted parameters are the semi-latus rectum p, eccentricity e, and orientation ω. The semi-major axis is then

$$a = \frac{p}{1 - e^2}$$

To compare all three models fairly, the final residual for each reconstructed point was the Euclidean distance to the nearest point on the fitted curve:

$$d_i = \min_{\mathbf{q} \in C} \|\mathbf{x}_i - \mathbf{q}\|$$

The reported RMSE values therefore use the same geometric error definition for all models:

$$\mathrm{RMSE} = \sqrt{\frac{1}{N}\sum_{i=1}^{N} d_i^2}$$

Residual comparison


Noise	Model	RMSE (AU)	MAE (AU)	Median error (AU)	Maximum error (AU)
1'	Sun-centered circle	0.051504	0.046848	0.042350	0.099385
1'	Free-center circle	0.021718	0.017068	0.012948	0.049116
1'	Kepler ellipse	0.021759	0.017124	0.013161	0.048913
2'	Sun-centered circle	0.051629	0.046963	0.042165	0.099916
2'	Free-center circle	0.021879	0.017224	0.012521	0.048463
2'	Kepler ellipse	0.021920	0.017268	0.012740	0.048260
5'	Sun-centered circle	0.052027	0.047300	0.043520	0.101501
5'	Free-center circle	0.022431	0.017721	0.012080	0.046529
5'	Kepler ellipse	0.022475	0.017760	0.011934	0.046580

For the 1 arcminute data, the fitted parameters were:


- Sun-centered circle radius: 1.52619 AU
- Free-circle center: (-0.09834, 0.01994) AU
- Free-circle radius: 1.48445 AU
- Fitted ellipse semi-major axis: 1.48547 AU
- Fitted ellipse eccentricity: 0.06689
- Fitted ellipse orientation: -12.63 degrees

The hidden generating values were $a = 1.52366\,\mathrm{AU}$ and $e = 0.09341$, so the ellipse fit recovered the correct general scale and a nonzero eccentricity, although the reconstruction underestimates the eccentricity.


1 arcminute model fits



2 arcminute model fits




5 arcminute model fits



The Sun-centered circular model is clearly worse. At 1 arcminute, changing from a Sun-centered circle to the ellipse lowers RMSE by about 57.75 percent.

However, the free-center circle and ellipse are almost indistinguishable for the reconstructed arc. The free-center circle is actually slightly better in RMSE by approximately 0.19 percent at every tested noise level. This does not mean the physical orbit is circular. It means that a shifted circle can approximate the limited reconstructed arc extremely well.

Effect of observational noise



Noise	Sun-circle RMSE	Free-circle RMSE	Ellipse RMSE	Ellipse minus free-circle RMSE	Ellipse vs free-circle difference	Ellipse improvement vs Sun circle
1'	0.051504	0.021718	0.021759	0.000041	0.188% worse	57.75% better
2'	0.051629	0.021879	0.021920	0.000041	0.188% worse	57.54% better
5'	0.052027	0.022431	0.022475	0.000043	0.193% worse	56.80% better

Therefore, within the tested noise range, there is no observational accuracy at which the ellipse clearly wins over the free-center circle. Even at 1 arcminute, the two models are effectively tied. The dominant limitation appears to be incomplete orbital coverage and reconstruction error rather than the difference between 1 and 5 arcminutes of measurement noise.

---

7. LLM Prompts and Main Suggestions

The LLM was used throughout the project as a collaborator. The prompts below are representative of the actual interaction; wording has been lightly cleaned for readability while preserving the original requests.

Prompt 1: Build the synthetic observation model

> Start with the code for synthetic Tycho-like observations of Mars. Explain how

> the Uraniborg coordinates are used and what equations are needed to generate the synthetic data. Include Earth motion, Mars motion, Earth rotation, and observational noise.

Main LLM suggestions

The LLM proposed:

- use a simplified heliocentric two-planet model with the Sun fixed at the origin;
- represent Earth and Mars using Kepler ellipses;
- solve Kepler's equation numerically with Newton's method;
- use Uraniborg latitude and longitude together with sidereal time to construct the local observer position and local east/north/up basis;
- convert ecliptic coordinates to equatorial coordinates using Earth's obliquity;
- calculate both RA/declination and altitude/azimuth;
- add angular noise in the tangent plane of the line-of-sight vector rather than directly perturbing Cartesian coordinates;
- reject observations when Mars is too low or the Sun is too high.

These suggestions became the basis of the synthetic data generator.

Prompt 2: Verify the simulation independently

> Verify the simulation code independently. Check energy and angular momentum for the noiseless Earth and Mars orbits and give me a standalone verification program.

Main LLM suggestions

The LLM recommended comparing the numerical values with the exact Kepler formulas

$$\mathcal{E} = -\frac{\mu}{2a}$$

$$h = \sqrt{\mu a(1 - e^2)}$$

It also recommended using relative errors instead of deciding visually that a curve looked flat. This led to an independent verification file rather than reusing the same functions blindly inside the data generator.

Prompt 3: Visualize Mars in Tycho's sky

> Plot the apparent position of Mars in Tycho's sky using the synthetic CSV files.

Main LLM suggestions

The LLM recommended plotting both RA/declination and altitude/azimuth. It also warned against connecting points across the 0/360 degree wrap in RA or azimuth, so scatter plots were used.

Prompt 4: Reconstruct Mars from lines of sight

> Convert the observations into lines of sight and reconstruct heliocentric Mars positions without using the hidden Mars orbital parameters.

Main LLM suggestions

The first suggestion was a repeated-phase triangulation method based on pairs of rays. That initial method was not reliable and is discussed in the critique below. After the failure was detected, the LLM proposed a corrected multi-ray method:

- search a broad range of candidate orbital periods rather than inserting Mars's true period;
- group observations separated by approximately integer multiples of the candidate period;
- use three or more rays for each candidate heliocentric point;
- solve a least-squares closest-point problem for the full ray bundle;
- reject bundles with poor geometric baseline;
- validate the inferred radii and out-of-plane coordinate before accepting the reconstruction.

The corrected method inferred a period of 689.25 days and physically reasonable heliocentric Mars distances.

Prompt 5: Compare circle and ellipse models

> Fit a Sun-centered circle, a circle with arbitrary center, and an ellipse to the reconstructed Mars positions. Compute comparable residual errors.

Main LLM suggestions

The LLM suggested using a common geometric residual for the final comparison: the Euclidean distance from each reconstructed point to the nearest point on each fitted curve. This avoided comparing incompatible residual definitions across different model types.

Prompt 6: Interpret the noise experiment

> Compare the 1, 2, and 5 arcminute results. How much observational accuracy is needed before the ellipse clearly wins?

Main LLM suggestions

The LLM did not force the historical conclusion. It observed that the Sun-centered circle is clearly worse, but the free-center circle and ellipse are almost tied at every tested noise level. It therefore suggested that the main limitation is the reconstructed orbital coverage rather than measurement noise alone.

Prompt 7: Interpret the result as if reasoning from the data

> Interpret the numerical results. Did the AI reason from the observations, or did it merely recall that Mars follows an ellipse?

Main LLM suggestion

The LLM concluded that the data justify rejection of a Sun-centered circle, but do not justify claiming that the ellipse clearly beats every alternative model. This interpretation is important because it follows the computed residuals even though the LLM already knows the historical fact that Mars follows an ellipse.

---

8. Critique of the LLM: What Was Useful and What Was Not

The LLM was most useful as a source of mathematical structure, implementation ideas, and verification strategies. It was less reliable when a plausible geometric method had hidden conditioning problems.

Useful contributions

1. It translated the physical setup into a modular computational model

The LLM separated the problem into orbital motion, coordinate transformations, Earth rotation, observer position, angular observations, noise, reconstruction, and model fitting. This modular organization made the program easier to inspect line by line and easier to verify.

2. It connected the code to explicit equations

Rather than only supplying Python code, the LLM explained Kepler's equation, the ellipse parameterization, velocity formulas, conserved energy, conserved angular momentum, RA/declination conversion, and the observer geometry. This made it possible to check whether individual code blocks had a mathematical interpretation.

3. Its verification strategy was strong

The most useful suggestion was to compare the computed energy and angular momentum with independent closed-form theoretical values. The approximately 1e-15 relative errors provided a much stronger verification than simply looking at orbital plots.

4. It helped identify physical sanity checks

The project benefited from simple physical checks such as:

- Earth should remain near 1 AU;
- Mars should lie roughly between 1.38 and 1.67 AU in the hidden model;
- a successful planar reconstruction should have small z values;
- an inferred orbital period should be near the scale suggested by repeated observations but should not be inserted directly from historical knowledge.

These checks became especially important when the first reconstruction failed.

5. It adapted when the numerical evidence contradicted expectations

The final model comparison did not show the ellipse clearly winning. The LLM correctly interpreted this instead of claiming that the result had confirmed Kepler simply because Mars is historically known to follow an ellipse.

Weak or incorrect contributions

1. The first reconstruction method was wrong in an important way

The initial LLM suggestion used pairs of repeated-phase lines of sight and chose the candidate period that made the rays intersect most closely. That method produced reconstructed radii near 1 AU, almost on Earth's orbit. The result looked geometrically clean but was physically impossible for Mars in the hidden model.

This was the clearest example of why the LLM could not be trusted without verification. A low ray-intersection residual did not imply that the inferred distance was correct. The

method could prefer a false near-Earth intersection when the observer geometry nearly repeated.

The error was detected only after checking the reconstructed physical scale. The solution was to replace the pairwise method with a multi-ray least-squares reconstruction and require a meaningful observer baseline.

2. The LLM initially had a natural bias toward expecting the ellipse to win

Because an LLM already knows Kepler's laws, it is easy for it to frame the experiment around the expected historical answer. The actual fitted residuals did not support a strong ellipse-over-shifted-circle conclusion. The final interpretation had to remain disciplined and follow the computed values rather than the prior historical fact.

3. Some modeling choices were practical rather than uniquely correct

The 10-day sampling interval, fixed 22:00 UTC observation time, 10-degree Mars altitude cutoff, -6-degree Sun altitude cutoff, planar planetary orbits, and simplified sidereal-time calculation are reasonable choices for a synthetic experiment, but they are not unique or fully historical. These choices should be described as assumptions, not as exact features of Tycho Brahe's observing program.

4. The reconstruction does not recover the entire orbit equally well

Only a subset of orbital phases produced sufficiently good repeated multi-ray geometry. The reconstructed points therefore cover a limited arc rather than a uniformly sampled complete ellipse. This helps explain why a shifted circle can imitate the fitted ellipse so closely.

Overall assessment of the LLM

The LLM was useful when treated as a collaborator that generates equations, code structures, diagnostic ideas, and alternative approaches. It was not reliable enough to accept without independent mathematical and physical checks. The failed first reconstruction is a useful example: the code ran, the geometry looked plausible, and the numerical residual was small, but the physical result was wrong. Verification changed the method and materially improved the project.

---

9. Did the AI Reason Like Kepler, or Did It Merely Remember Kepler?

The experiment does not justify claiming that the AI independently rediscovered Kepler's law in the historical sense. The LLM already knows that Mars follows an elliptical orbit, and the synthetic data themselves were generated from a hidden Kepler ellipse. Therefore the AI cannot be treated as an observer with no prior astronomical knowledge.

However, the final analysis did show a meaningful form of data-based reasoning. When the residuals were computed, the LLM did not simply say that the ellipse must be best because Kepler discovered ellipses. The numerical evidence showed that the Sun-centered circle was clearly worse, while the free-center circle and Kepler ellipse had almost identical residual errors. In fact, the free-center circle had a slightly lower RMSE at all three tested noise levels. The LLM therefore concluded only what the data supported: a Sun-centered circle can be rejected, but the available reconstructed arc is not sufficient to distinguish an ellipse clearly from a shifted circle.

The best description is that the AI acted more like a modern scientific collaborator than a true historical Kepler. It supplied hypotheses and computational methods, made at least one important mistake, and then revised the method after the mistake was exposed by physical verification. Its strongest reasoning occurred when it followed the numerical evidence even when that evidence did not reproduce the expected historical conclusion. The project therefore demonstrates both the usefulness and the limitation of LLM-assisted scientific work: the LLM can accelerate exploration, but the student must still decide what the data actually prove.

Discussion: Kepler's Problem as Model Fitting from Data

This problem is an early example of model fitting from data because we begin with observations and ask which mathematical model best explains them. Instead of assuming the correct orbit in advance, we compare candidate models, fit their parameters, and use residual errors to determine how well each model explains the reconstructed observations.

Kepler did something conceptually similar when he analyzed Tycho Brahe's measurements of Mars. He did not simply accept the traditional circular description because it was mathematically attractive. The observational discrepancies forced him to consider a different model. In that sense, the historical process resembles modern approximation: observations constrain a family of candidate mathematical descriptions, and the model is judged by how well it accounts for the data.

The general workflow is:

1. collect observations;
2. choose candidate models;

3. fit model parameters;
4. compute residual errors;
5. compare competing explanations;
6. retain only conclusions justified by the data.

This is conceptually related to modern machine learning and approximation theory. In many modern problems, the true input-output relationship is unknown. A parameterized model is fitted to observations by minimizing a loss or error. Kepler was not using a neural network or gradient descent, but the underlying scientific logic is similar: infer mathematical structure from data rather than imposing the preferred model in advance.

The current experiment also demonstrates an important limitation of model fitting. The observations strongly reject the simplest Sun-centered circle, but they do not clearly distinguish the ellipse from a shifted circle. A scientifically valid conclusion must therefore stop at what the available data support. Knowing the correct historical answer is not a substitute for evidence from the experiment.

Files Submitted with This Report

Code

- `codes/synthetic_data_generation(step_1).ipynb`
- `codes/verify_orbits(step_2).ipynb`
- `codes/plot_mars_sky(step_3).ipynb`
- `codes/reconstruct_mars(step_4).ipynb`
- `codes/plot_reconstructed_mars(step_5).ipynb`
- `codes/rigorous_reconstructed_mars(step_5-1).ipynb`
- `codes/fit_models(step6).ipynb`
- `codes/noise_comparison(step7).ipynb`
- `codes/LLM_reasoning(step8).ipynb`
- `codes/dicussion_model_fitting(step9).ipynb`

Data

- `data/synthetic_mars_1arcmin.csv`
- `data/synthetic_mars_2arcmin.csv`
- `data/synthetic_mars_5arcmin.csv`
- `data/orbit_verification_summary.csv`
- `data/earth_orbit_verification.csv`
- `data/mars_orbit_verification.csv`
- `data/period_search_1arcmin.csv`

- `data/period_search_2arcmin.csv`
- `data/period_search_5arcmin.csv`
- `data/mars_reconstructed_1arcmin.csv`
- `data/mars_reconstructed_2arcmin.csv`
- `data/mars_reconstructed_5arcmin.csv`
- `data/model_fit_summary.csv`
- `data/step7_noise_comparison.csv`

Figures

All generated figures are stored in the `result images` directory.