

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

CJC-1295 (No DAC) + Ipamorelin 5mg/5mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your CJC/Ipamorelin Vial

Time: 5 minutes | Location: Clean, flat surface

This is a pre-blended vial containing 5mg CJC-1295 (No DAC) and 5mg Ipamorelin (10mg total peptide). Reconstitution prepares the lyophilized blend for subcutaneous injection.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
30-35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
5mg/5mg CJC/IPA blend vial	Lyophilized peptide blend, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and peptide vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.

2. Pull back to 100 units (1mL) of BAC water.
3. Insert the needle into the peptide vial. Allow the vacuum to pull the liquid in slowly.

Critical Warning

Do not push the syringe. The vial is vacuum-sealed and draws the liquid in on its own. Forcing the plunger compromises the vacuum and may cause inaccurate dosing.

4. Gently roll the vial between your fingers. Do not shake. Let fully dissolve for 30–60 seconds.
5. Place the vial in the refrigerator. After 1–2 hours refrigerated, the solution is ready for use.

Storage Requirement

Reconstituted CJC/Ipamorelin must remain refrigerated at all times. Do not freeze. Do not expose to direct sunlight or heat.

Dose Conversion

100 units total — each 1 unit = 50mcg CJC + 50mcg Ipamorelin (100mcg total)

Common Research Doses

2 units	100mcg CJC + 100mcg IPA (200mcg total)
3 units	150mcg CJC + 150mcg IPA (300mcg total)
4 units	200mcg CJC + 200mcg IPA (400mcg total)
6 units	300mcg CJC + 300mcg IPA (600mcg total)

Vial Duration

2 units daily	50 days
4 units daily	25 days
6 units daily	≈16 days

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Stomach area (preferred)

Administer your dose subcutaneously using the 30–35-unit insulin syringe. The stomach area is the preferred injection site.

Products Needed

Equipment

Peptide vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
30-35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the peptide vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to your prescribed unit mark using the conversion table above.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle. Stomach area is preferred.
2. Inject slowly and steadily.
3. After injecting, lightly rub the area in a circular motion to aid absorption.

Stage 3: Dosing Protocol

3

Timing, Frequency, and Fasting Window

Time: Ongoing | 1-3x daily

CJC-1295 (No DAC) + Ipamorelin is administered 1–3 times daily depending on protocol goals. Injection timing relative to food intake directly impacts GH pulse amplitude.

Important Reminders

Protocol

Refrigeration	Always keep the reconstituted vial refrigerated
Sterilization	Always wipe the vial stopper and injection site before each use
Fasting requirement	Inject on an empty stomach
Food window	Avoid carbohydrates and fats 60–90 minutes before and after injection
Frequency	1–3 injections daily depending on protocol

Protocol Note

Carbohydrates and fats blunt the GH pulse that CJC/Ipamorelin triggers. Maintaining a strict fasting window of at least 60 minutes before and after injection produces optimal growth hormone output. Protein-only meals do not significantly suppress the pulse.

Injection Timing Examples

Objective

Maximize natural GH pulses by injecting during fasted windows throughout the day.

Option A: Morning	Inject fasted immediately upon waking, wait 60–90 min before eating
Option B: Pre-bed	Inject at least 2–3 hours after last meal, before sleep
Option C: Multi-dose	Morning fasted + pre-bed for enhanced GH output

Stage 4: What to Expect

4

Response Timeline and Outcomes

Timeline: Weeks 1–12+

CJC-1295 (No DAC) + Ipamorelin stimulates natural growth hormone pulses from the pituitary gland. Results are progressive and compound with consistent daily use.

Common Observations

Weeks 1–12+

Sleep quality	Deeper, more restorative sleep — often the first noticeable change
Recovery	Faster recovery from training and daily activity
Muscle fullness	Increased muscle density and fullness
Fat metabolism	Enhanced lipolysis and improved body composition
Water retention	Mild and temporary as the body adjusts to elevated GH

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com